

BIRCH, STEWART, KOLASCH & BIRCH, LLP

INTELLECTUAL PROPERTY LAW
8110 GATEHOUSE ROAD
SUITE 500 EAST
FALLS CHURCH, VA 22042-1210
U S A
(703) 205-8000

FAX: (703) 205-8050
(703) 898-8590 (G IV)

e-mail: mailroom@bskb.com
web: http://www.bskb.com

CALIFORNIA OFFICE:
COSTA MESA, CALIFORNIA

THOMAS S. AUCHTERLONIE
JAMES T. BLER, JR.
SCOTT L. LOWE
MARK J. NIELL, Ph.D.
D. RICHARD ANDERSON
PAUL C. LEWIS
MARK W. MILSTEAD
RICHARD L. GALLAGHER
JAYNE M. SAYDAH

REG. PATENT AGENTS
FREDERICK R. HANDREN
MARYANNE ARMSTRONG, Ph.D.
MAKI HATSUMI
MIKE S. RYU
CRAIG A. MICROBBIE
GARTH M. DAHLEN, Ph.D.
LAURA C. LUTZ
ROBERT E. GOOZNER, Ph.D.
HYUNG M. SOHN
MATTHEW J. LATTIG
ALAN PEDERSEN-GILES
C. KEITH MONTGOMERY
TIMOTHY R. WYCKOFF
KRISTILL RUPERT, Ph.D.
LARRY J. HUME
ALBERT LEE
HRAIR A. SAYADIAN, Ph.D.

Date: October 27, 2000

Docket No.: 20-4764P

JC937 U.S. PRO
09/697719
10/27/00

Assistant Commissioner for Patents
Washington, DC 20231

Sir:

As authorized by the inventor(s), transmitted herewith for filing is a continuation-in-part application of Appl. No. 09/302,357 filed on April 30, 1999.

Inventor(s): NAKAJIMA, Hiroki and NAGASAWA, Akitsu

For: METHOD FOR GIVING RESISTANCE TO WEED CONTROL
COMPOUNDS TO PLANTS

Enclosed are:

- ☒ A specification consisting of three hundred thirteen (313) pages
- ☒ Thirteen (13) sheet(s) of formal drawings
- ☒ Executed Declaration ☒ Original ☐ Photocopy
- ☒ Preliminary Amendment
- ☒ An assignment of the invention to Sumitomo Chemical Co., Ltd.
- ☐ Certified copy of
- ☐ Priority of application No(s). 120553/1998 filed in JAPAN on April 30, 1998 is claimed under 35 U.S.C. § 119. See attached copy of the Letter claiming priority filed in the prior application on _____.

- ☒ An Information Disclosure Statement and PTO-1449 form(s) are attached hereto for the Examiner's consideration.
- ☐ A statement (☐ Original ☐ Photocopy) to establish small entity status under 37 C.F.R. § 1.9 and 37 C.F.R. § 1.27.
- ☐ Amend the specification by inserting before the first sentence

--This application is a continuation-in-part of Application No. 09/302,357 filed on April 30, 1999, the entire contents of which are hereby incorporated by reference.--

- ☐ Amend the specification by inserting before the first sentence

--This application is a continuation-in-part of PCT international application No. PCT/____ that has an international filing date of ____ that designated the United States, the entire contents of which are hereby incorporated by reference.--

- ☐ An extension of time for ____ () month(s) until ____, has been submitted in parent Application No. 09/302,357 in order to establish dependency with the present application.
- ☐ Other: _____

The filing fee has been calculated as shown below:

			LARGE ENTITY	SMALL ENTITY
BASIC FEE			\$710.00	\$355.00
	NUMBER FILED	NUMBER EXTRA	RATE FEE	RATE FEE
TOTAL CLAIMS	79-20 =	59	x 18 = \$1062.00	x 9 = \$0.00
INDEPENDENT CLAIMS	3-3 =	0	x 80 = \$0.00	x 40 = \$0.00
MULTIPLE DEPENDENT YES CLAIMS PRESENTED			+ \$270.00	+ \$135.00
TOTAL			\$2,042.00	\$0.00

- ☒ A check in the amount of \$2,082.00 to cover the filing fee and recording fee (if applicable) is enclosed.
- ☐ Please charge Deposit Account No. 02-2448 in the amount of \$0.00. A triplicate copy of this transmittal form is enclosed.
- ☒ Please send correspondence to:

BIRCH, STEWART, KOLASCH & BIRCH, LLP **or** Customer No. 2292
P.O. Box 747
Falls Church, VA 22040-0747
Telephone: (703) 205-8000

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By  #32,868

Gerald M. Murphy, Jr., #28,977

GMM/rem
20-4764P

P.O. Box 747
Falls Church, VA 22040-0747
(703) 205-8000

Attachments

(Rev. 09/29/00)

IN THE U.S. PATENT AND TRADEMARK OFFICE

Applicant: NAKAJIMA, Hiroki et al.
Appl. No.: NEW Group:
Filed: October 27, 2000 Examiner:
For: METHOD FOR GIVING RESISTANCE TO WEED
CONTROL COMPOUNDS TO PLANTS

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents October 27, 2000
Washington, DC 20231

Sir:

The following preliminary amendments and remarks are respectfully submitted in connection with the above-identified application.

AMENDMENTS

IN THE CLAIMS:

Please amend the claims as follows:

CLAIM 24: Line 1, change "claim 22 or 23" to

--Claim 22-

CLAIM 25: Line 1, change "claim 22 or 23" to

--Claim 22--

CLAIM 26: Line 1, change "claim 22 or 23" to

--Claim 22-

CLAIM 27: line 1, change "claim 22 or 23" to

--Claim 22--

REMARKS

The claims have been amended to delete improper multiple dependencies.


Entry of the above amendments is earnestly solicited. An early and favorable first action on the merits is earnestly solicited.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By

 #32, 868
Gerald M. Murphy, Jr., #28,977

GMM/rem
20-4764P

P.O. Box 747
Falls Church, VA 22040-0747
(703) 205-8000

(Rev. 04/19/2000)

METHOD FOR GIVING RESISTANCE TO
WEED CONTROL COMPOUNDS TO PLANTS

BACKGROUND OF THE INVENTION

FIELD OF THE INVENTION

The present invention relates to a method for giving resistance to weed control compounds to plants.

DISCLOSURE OF THE RELATED ART

Weed control is very important work for improving yields and quality of cultivated plants. For this purpose, weed control compounds such as herbicides are mainly used. However, for using weed control compounds, it is not always easy to distinguish cultivated plants from weeds of allied species to selectively control only weeds. Then, production of plants having resistance to weed control compounds (hereinafter referred to as weed control compound-resistance) has been attempted and some resistant plants have been put to practical use.

Recently, gene engineering techniques have been utilized for producing plants having weed control compound-resistance. As such a technique, for example, Hinchee, M.A.W. et al. disclose a method for producing a plant having resistance to a herbicide, glyphosate, wherein 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) gene

which is a target enzyme of glyphosate is mutagenized so that an affinity for glyphosate is reduced, and the gene is introduced into a plant [Hinchee, M.A.W. et al., BIO/TECHNOLOGY, 6: p 915 (1988)].

5

OBJECTS OF THE INVENTION

Varieties of known methods for giving weed control compound-resistance to plants are not necessarily sufficient and it has been desired to develop further various kinds of methods for giving weed control compound-resistance to plants.

10

The main object of the present invention is to provide a new kind of a method for giving weed control compound-resistance to plants.

15

This object as well as other objects and advantages of the present invention will become apparent to those skilled in the art from the following description with reference to the accompanying drawings.

20

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is the restriction map of plasmid pETBCH. bchH is magnesium chelatase protoporphyrin IX binding subunit gene of a photosynthetic bacterium *Rhodobacter sphaeroides*. T7 pro represents the promoter sequence of T7 phage, and T7 ter represents the terminator sequence of T7

25

phage. Amp^r is an ampicillin resistant gene, lacI^r is a repressor protein gene of a lactose operon, and ori is the replication origin.

Fig. 2 is the restriction map of plasmid pACYCSP.

5 PPO is protoporphyrinogen IX oxidase gene of soybean and lac pro represents the promoter sequence of a lactose operon. Cm^r is a chloramphenicol resistant gene and ori is the replication origin.

Fig. 3 is the restriction map of plasmid pTVBCH.

10 bchH is magnesium chelatase protoporphyrin IX binding subunit gene of the photosynthetic bacterium *Rhodobacter sphaeroides*. lac pro represents the promoter sequence of a lactose operon. Amp^r is an ampicillin resistant gene and ori is the replication origin.

15 Fig. 4 is the restriction map of plasmid pBIBCH. bchH is magnesium chelatase protoporphyrin IX binding subunit gene of the photosynthetic bacterium *Rhodobacter sphaeroides*. NP is the promoter sequence of a nopaline synthase gene, NT is the terminator sequence of the
20 nopaline synthase gene, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII represents a kanamycin resistant gene, and RB and LB represent right and left border sequences of T-DNA, respectively.

Fig. 5 is the restriction map of plasmid pNO. NP
25 is the promoter sequence of a nopaline synthase gene, NT is

the terminator sequence of the nopaline synthase gene, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII represents a kanamycin resistant gene, and RB and LB represent right and left border sequences of T-DNA, respectively.

Fig. 6 is the restriction map of plasmid pTCHLH. TCHLH is protoporphyrin IX binding subunit gene of tobacco magnesium chelatase whose chloroplast transit signal has been deleted. lac pro represents the promoter sequence of a lactose operon. Amp^r is an ampicillin resistant gene, Km^r is a kanamycin resistant gene and ori is the replication origin.

Fig. 7 is the restriction map of plasmid pBITCHLH. TCHLH is protoporphyrin IX binding subunit gene of tobacco magnesium chelatase whose chloroplast transit signal has been deleted. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of the nopaline synthase and 35S is the 35S promoter of cauliflower mosaic virus. NPTII represents a kanamycin resistant gene, and RB and LB represent right and left border sequences of T-DNA, respectively.

Fig. 8 is the restriction map of plasmid pTVGMP. GMP is soybean protoporphyrinogen IX oxidase gene whose chloroplast transit signal and FAD binding sequence have been deleted. lac pro represents the promoter sequence of

a lactose operon. Amp^r represents an ampicillin resistant gene and ori is the replication origin.

Fig. 9 is the restriction map of plasmid pBIGMP. GMP is soybean protoporphyrinogen oxidase gene whose chloroplast transit signal and FAD binding sequence have been deleted. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 10 is the restriction map of plasmid pTVCRP. CRP is protoporphyrinogen oxidase gene of *Chlamydomonas reinhardtii* whose chloroplast transit signal and FAD binding sequence have been deleted. lac pro represents the promoter sequence of a lactose operon. Amp^r is an ampicillin resistant gene and ori is the replication origin.

Fig. 11 is the restriction map of plasmid, pBICRP. CRP is protoporphyrinogen oxidase gene of *Chlamydomonas reinhardtii* whose chloroplast transit signal and FAD binding sequence have been deleted. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border

sequences of T-DNA, respectively.

Fig. 12 is the restriction map of plasmid pTVHVF1. HVF is barley ferrochelataase gene whose signal sequence has been deleted. lac pro represents the promoter sequence of a lactose operon. Amp^r represents an ampicillin resistant gene and ori is the replication origin.

Fig. 13 is the restriction map of plasmid pBIHVF. HVF is barley ferrochelataase gene whose signal sequence has been deleted. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 14 is the restriction map of plasmid pTVCSF. CSF is cucumber ferrochelataase gene whose signal sequence has been deleted. lac pro represents the promoter sequence of a lactose operon. Amp^r is an ampicillin resistant gene, and ori is the replication origin.

Fig. 15 is the restriction map of plasmid pBICSF. CSF is cucumber ferrochelataase gene whose signal sequence has been deleted. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant

gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 16 is the restriction map of plasmid pHEMF. HEMF is coproporphyrinogen III oxidase gene (hemF) of *Escherichia coli*. lac pro is the promoter sequence of a lactose operon. Amp^r is an ampicillin resistant gene, and ori is the replication origin.

Fig. 17 is the restriction map of plasmid pBIHEMF. HEMF is coproporphyrinogen III oxidase gene (hemF) of *Escherichia coli*. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 18 is the restriction map of plasmid pBIHASYS8. HASYS8 is a gene encoding MG(HASYS)₈ protein. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 19 is the restriction map of plasmid pBIRASSL8. RASSL8 is MG(RASSL)₈ protein. NP is the promoter sequence of a nopaline synthase, NT is the

terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 20 is the restriction map of plasmid pNATP. PPO(A220V) is a PPO gene having a herbicidal compound-resistant mutation (A220V). NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 21 is the restriction map of plasmid pBIAPTCH. PPO(A220V) is a PPO gene having a herbicidal compound-resistant mutation (A220V) and TCHLH is a tobacco magnesium chelatase subunit gene whose chloroplast transit signal has been deleted. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 22 is the restriction map of plasmid pCRATF. ATF is a chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana*. lac pro represents the promoter

sequence of a lactose operon. Amp^r is an ampicillin resistant gene, Km^r is kanamycin resistant gene and ori is the replication origin.

Fig. 23 is the restriction map of plasmid pBIATF.

ATF is a chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana*. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 24 is the restriction map of plasmid

pBIAPATF. PPO(A220V) is PPO gene having a herbicidal compound-resistant mutation (A220V). ATF is a chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana*. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 25 is the restriction map of plasmid

PCRSCPOX. SCPOX is soybean coproporphyrinogen III oxidase gene and lac pro represents the promoter sequence of a lactose operon. Amp^r is an ampicillin resistant gene, Km^r is a kanamycin resistant gene and ori is the replication

origin.

Fig. 26 is the restriction map of plasmid pBISCPOX. SCPOX is soybean coproporphyrinogen III oxidase gene. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 27 is the restriction map of plasmid pBIAPSCP. PPO(A220V) is PPO gene having a herbicidal compound-resistant mutation (A220V). SCPOX is soybean coproporphyrinogen III oxidase gene. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 28 is the restriction map of plasmid pCREPSFS. CTP-EPSFS is a variant gene in which EPSFS gene derived from *Agrobacterium* is ligated to the downstream of a nucleotide sequence encoding a chloroplast transit peptide of EPSFS derived from petunia. lac pro represents the promoter sequence of a lactose operon. Amp^r is an ampicillin resistant gene, Km^r is kanamycin resistant gene and ori is the replication origin.

Fig. 29 is the restriction map of plasmid pNG01. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, GUS is β -glucuronidase gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 30 is the restriction map of plasmid pNG04. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, GUS is β -glucuronidase gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 31 is the restriction map of plasmid pNT35S. NT is the terminator sequence of a nopaline synthase, 35S is the 35S promoter of cauliflower mosaic virus, and lac pro is the promoter sequence of a lactose operon. Amp^r is an ampicillin resistant gene and ori is the replication origin.

Fig. 32 is the restriction map of plasmid pCENS. CTP-EPSPS is a variant gene in which EPSPS gene derived from *Agrobacterium* is ligated to the downstream of a nucleotide sequence encoding a chloroplast transit peptide of EPSPS derived from petunia. NT is the terminator

sequence of a nopaline synthase, 35S is the 35S promoter of cauliflower mosaic virus, and lac pro is the promoter sequence of a lactose operon. Amp^r is an ampicillin resistant gene, ori is the replication origin.

5 Fig. 33 is the restriction map of plasmid pCENSK. CTP-EPSPS is a variant gene in which EPSPS gene derived from *Agrobacterium* is ligated to the downstream of a nucleotide sequence encoding a chloroplast transit peptide of EPSPS derived from petunia. NT is the terminator sequence of a nopaline synthase, 35S is the 35S promoter of cauliflower mosaic virus, and lac pro is the promoter sequence of a lactose operon. Amp^r is an ampicillin resistant gene, ori is the replication origin.

10 Fig. 34 is the restriction map of plasmid pBICE. CTP-EPSPS is a variant gene in which EPSPS gene derived from *Agrobacterium* is ligated to the downstream of a nucleotide sequence encoding a chloroplast transit peptide of EPSPS derived from petunia. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and PB and LB are the right and left border sequences of T-DNA, respectively.

20 Fig. 35 is the restriction map of plasmid pBICETCH. CTP-EPSPS is a variant gene in which EPSPS gene

derived from *Agrobacterium* is ligated to the downstream of a nucleotide sequence encoding a chloroplast transit peptide of EPSPS derived from petunia. TCHLH is protoporphyrin IX binding subunit gene of tobacco magnesium chelatase whose chloroplast transit signal has been deleted. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 36 is the restriction map of plasmid pBIGMP. GMP is soybean PPO gene whose chloroplast transit signal and FAD binding sequence have been deleted. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 37 is the restriction map of plasmid pBICEGMP. CTP-EPSPS is a chimera gene in which EPSPS gene derived from *Agrobacterium* is ligated to the downstream of a nucleotide sequence encoding a chloroplast transit peptide of EPSPS derived from petunia. GMP is soybean PPO gene whose chloroplast transit signal and FAD binding gene have been deleted. NP is the promoter sequence of a

nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 38 is the restriction map of plasmid pBICRP. CRP is PPO gene of *Chlamydomonas reinhardtii* whose chloroplast transit signal and FAD binding sequence have been deleted. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 39 is the restriction map of plasmid pBICECRP. CTP-EPSPS is a variant gene in which EPSPS gene derived from *Agrobacterium* is ligated to the downstream of a nucleotide sequence encoding a chloroplast transit peptide of EPSPS derived from petunia. CRP is PPO gene of *Chlamydomonas reinhardtii* whose chloroplast transit signal and FAD binding sequence have been deleted. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and

left border sequences of T-DNA, respectively.

Fig. 40 is the restriction map of plasmid pBICEATF. CTP-EPSPS is a variant gene in which EPSPS gene derived from *Agrobacterium* is ligated to the downstream of a nucleotide sequence encoding a chloroplast transit peptide of EPSPS originated from petunia. ATF is chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana*. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 41 is the restriction map of plasmid pBISCPOX. SCPOX is soybean coproporphyrinogen III oxydase gene. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

Fig. 42 is the restriction map of plasmid pBICESCPOX. CTP-EPSPS is a variant gene in which EPSPS gene derived from *Agrobacterium* is ligated to the downstream of a nucleotide sequence encoding a chloroplast transit peptide of EPSPS derived from petunia. SCPOX is

soybean coproporphyrinogen III oxydase gene. NP is the promoter sequence of a nopaline synthase, NT is the terminator sequence of a nopaline synthase, and 35S is the 35S promoter of cauliflower mosaic virus. NPTII is a kanamycin resistant gene, and RB and LB are the right and left border sequences of T-DNA, respectively.

SUMMARY OF THE INVENTION

Under these circumstances, the present inventors have studied intensively so as to develop a new kind of a method for giving weed control compound-resistance to plants. As a result, it has been found that weed control compound-resistance can be given to plants by allowing the plants to produce a certain protein in the plant cells. Thus, the present invention has been completed.

That is, the present invention provides:

1. A method for giving weed control compound-resistance to a plant which comprises the steps of:

introducing a gene encoding a protein having the following characteristics (a) to (c):

(a) having a specific affinity for a substance which is concerned with the weed control activity of a weed control compound,

(b) having substantially no capability of modifying a substance for which said protein has a specific

affinity, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin, into a plant cell; and

5 expressing the gene (hereinafter referred to as the first aspect of the method of the present invention).

2. The method according to the above 1, wherein the gene is introduced into the plant cell in the form that it is operably ligated to a promoter and a terminator both
10 of which are functional in the plant cell.

3. The method according to the above 1 or 2, wherein the substance which is concerned with the weed control activity of the weed control compound is the weed control compound itself.

15 4. The method according to the above 1 or 2, wherein the substance which is concerned with the weed control activity of a weed control compound is an endogenous substance in a plant.

20 5. The method according to the above 1 or 2, wherein the weed control compound is that inhibiting porphyrin biosynthesis of a plant.

6. The method according to the above 1 or 2, wherein the weed control compound is a protoporphyrinogen IX oxidase inhibitory-type herbicidal compound.

25 7. The method according to the above 5 or 6,

wherein the substance which is concerned with the weed control activity of the weed control compound is protoporphyrin IX.

8. The method according to the above 5 or 6,
5 wherein the protein is protoporphyrin IX binding subunit protein of magnesium chelatase, or a variant of said protein having a specific affinity for protoporphyrin IX.

9. The method according to the above 8, wherein
10 the protein is magnesium chelatase derived from a photosynthetic microorganism.

10. The method according to the above 8, wherein
the protein is magnesium chelatase derived from a plant.

11. The method according to the above 8, wherein
the protein is magnesium chelatase derived from tobacco.

12. The method according to the above 5 or 6,
15 wherein the protein comprises the amino acid sequence of SEQ ID NO: 53.

13. The method according to the above 5 or 6,
20 wherein the protein has the amino acid sequence of SEQ ID NO: 54.

14. The method according to the above 5 or 6,
wherein the protein comprises the amino acid sequence of SEQ ID NO: 55.

15. The method according to the above 5 or 6,
25 wherein the protein has the amino acid sequence of SEQ ID

NO: 56.

16. The method according to the above 5 or 6,
wherein the protein comprises the amino acid sequence of SEQ
ID NO: 57.

5 17. The method according to the above 5 or 6,
wherein the protein has the amino acid sequence of SEQ ID
NO: 58.

10 18. The method according to the above 5 or 6,
wherein the protein comprises the amino acid sequence of SEQ
ID NO: 59.

19. The method according to the above 5 or 6,
wherein the protein has the amino acid sequence of SEQ ID
NO: 60.

15 20. The method according to the above 5 or 6,
wherein the protein is composed of 4 to 100 amino acids.

21. The method according to the above 5 or 6,
wherein the substance which is concerned with the weed
control activity of the weed control compound is
protoporphyrinogen IX.

20 22. The method according to the above 5 or 6,
wherein the protein is a variant of protoporphyrinogen IX
oxidase having no capability of oxidizing
protoporphyrinogen IX and having a specific affinity for a
protoporphyrinogen IX.

25 23. The method according to the above 5 or 6,

wherein the protein is a variant of protoporphyrinogen IX
oxidase having no capability of oxidizing
protoporphyrinogen IX and having a specific affinity for a
protoporphyrin IX oxidase inhibitory-type herbicidal
5 compound.

24. The method according to the above 22 or 23,
wherein the protein is a variant of protoporphyrinogen IX
oxidase derived from a plant.

25. The method according to the above 22 or 23,
10 wherein the protein is a variant of protoporphyrinogen IX
oxidase derived from soybean.

26. The method according to the above 22 or 23,
wherein the protein is a variant of protoporphyrinogen IX
oxidase derived from an algae.

27. The method according to the above 22 or 23,
15 wherein the protein is a variant of protoporphyrinogen IX
oxidase derived from *Chlamydomonas*.

28. A method for giving weed control compound-
resistance to a plant which comprises the steps of:

20 introducing a gene encoding a protein having the
following characteristics (a) to (c):

(a) having a specific affinity for
protoporphyrin IX,

(b) having substantially no capability of
25 modifying protoporphyrinogen IX, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin, into a plant cell; and

expressing the gene (hereinafter referred to as the second aspect of the method of the present invention).

29. The method according to the above 28, wherein the gene is introduced in the plant cell in the form that it is operably ligated to a promoter and a terminator both of which are functional in the plant cell.

30. The method according to the above 28 or 29, wherein the weed control compound is that inhibiting porphyrin biosynthesis of a plant.

31. The method according to the above 28 or 29, wherein the weed control compound is a protoporphyrinogen IX oxidase inhibitory-type herbicidal compound.

32. The method according to the above 30 or 31, wherein the protein is magnesium chelatase or a variant of said protein having a specific affinity for protoporphyrin IX.

33. The method according to the above 30 or 31, wherein the protein is ferrochelatase or a variant of said protein having an specific affinity for protoporphyrin IX.

34. The method according to the above 30 or 31, wherein the protein is ferrochelatase derived from a plant.

35. The method according to the above 30 or 31, wherein the protein is ferrochelatase derived from barley.

36. The method according to the above 30 or 31, wherein the protein is ferrochelatase derived from cucumber.

5 37. The method according to the above 30 or 31, wherein the protein is a peptide composed of 4 to 100 amino acids.

38. A method for giving weed control compound-resistance to a plant which comprises the steps of:

10 introducing a gene encoding a protein having the following characteristics (a) to (c):

(a) having a specific affinity for protoporphyrinogen IX,

15 (b) having the capability for modifying coproporphyrinogen III, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin, into a plant cell; and

20 expressing the gene (hereinafter referred to as the third aspect of the method of the present invention).

39. The method according to the above 38, wherein the gene is introduced into the plant cell in the form that it is operably ligated to a promoter and a terminator both of which are functional in the plant cell.

25 40. The method according to the above 38 or 39,

wherein the protein is coproporphyrinogen III oxidase or a variant of said protein having a specific affinity for protoporphyrinogen IX.

41. The method according to the above 38 or 39,
5 wherein the protein is coproporphyrinogen III oxidase derived from a microorganism.

42 The method according to the above 38 or 39,
wherein the protein is coproporphyrinogen III oxidase derived from *Escherichia coli*.

43. A weed control compound-resistant plant
10 whose resistance is given by the method of the above 1, 2, 28 or 29.

44. A weed control compound-resistant plant
whose resistance is given by the method of the above 38 or
15 39.

45. A method for protecting a plant which comprises applying the weed control compound to a growth area of the plant of the above 43.

46. A method for protecting a plant which
20 comprises applying the weed control compound to a growth area of the plant of the above 44.

47. A method for selecting a plant which comprises applying a weed control compound to which the plant of the above 43 is resistant to a growth area of the
25 plant of the above 43 and other plants, and selecting

either plant on the basis of difference in growth between the plants.

48. A method for selecting a plant which comprises applying a weed control compound to which the
 5 plant of the above 44 is resistant to a growth area of the plant of the above 44 and other plants, and selecting either plant on the basis of difference in growth between the plants.

49. The method according to the above 47,
 10 wherein the plants are plant cells.


50. The method according to the above 48, wherein the plants are plant cells.

51. The method according to the above 1 or 2, wherein the weed control compound is a protoporphyrinogen
 15 IX oxidase inhibitory-type herbicidal compound selected from the compounds of (1) to (3) below, and the substance which is concerned with the weed control activity of the weed control compound is protoporphyrin IX, protoporphyrinogen IX or a protoporphyrinogen IX oxidase
 20 inhibitory-type herbicidal compound:


(1) chlormethoxynil, bifenox, chlornitrofen (CNP),
 acifluorfen (5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoic acid) and its ethyl ester, acifluorfen-sodium,
 oxyfluorfen (2-chloro-1-(3-ethoxy-4-nitrophenoxy)-4-
 25 trifluoromethylbenzene), oxadiazon (3-[2,4-dichloro-5-(1-

methylethoxy)phenyl]-5-(1,1-dimethylethyl)-1,3,4-oxadiazol-
2(3H)-one), 2-[4-chloro-2-fluoro-5-(prop-2-ynyloxy)phenyl]-
2,3,4,5,6,7-hexahydro-1H-isoindol-1,3-dione, chlorphthalim
(N-(4-chlorophenyl)-3,4,5,6-tetrahydrophtalimide), TNPP-
ethyl (ethyl 2-[1-(2,3,4-trichlorophenyl)-4-nitropyrzoly]-
5-oxy]propionate), or N3-(1-phenylethyl)-2,6-dimethyl-5-
propyonylnicotinamide;


(2) a compound represented by the general
formula: J-G (I), wherein G is a group represented by
any one of the following general formulas G-1 to G-9 and J
is a group represented by any one of the following general
formulas of J-1 to J-30:




G-1




G-2




G-3




G-4




G-5




G-6




G-7




G-8




G-9




J-1




J-2




J-3




J-4




J-5




J-6




J-7




J-8




J-9




J-10




J-11




J-12




J-13




J-14




J-15




J-16




J-17




J-18




J-19




J-20




J-21




J-22




J-23




J-24




J-25




J-26




J-27



J-28



J-29



J-30

wherein the dotted lines in the formulas J-5, J-6, J-12 and J-24 represent that the left hand ring contains only single bonds, or one bond in the ring is a double bond between carbon atoms;

X is oxygen atom or sulfur atom;

Y is oxygen atom or sulfur atom;

R¹ is hydrogen atom or halogen atom;

R² is hydrogen atom, C₁-C₈alkyl group, C₁-C₈ haloalkyl group, halogen atom, OH group, OR²⁷ group, SH group, S(O)_pR²⁷ group, COR²⁷ group, CO₂R²⁷ group, C(O)SR²⁷ group, C(O)NR²⁹R³⁰ group, CHO group, CR²⁷=NOR³⁸ group, CH=CR³⁷CO₂R²⁷ group, CH₂CHR³⁷CO₂R²⁷ group, CO₂N=CR³¹R³² group;

nitro group, cyano group, $\text{NHSO}_2\text{R}^{33}$ group, $\text{NHSO}_2\text{NHR}^{33}$ group, $\text{NR}^{27}\text{R}^{38}$ group, NH_2 group or phenyl group optionally substituted with one or more and the same or different C_1 - C_6 alkyl groups;

5 p is 0, 1 or 2;

R^3 is C_1 - C_2 alkyl group, C_1 - C_2 haloalkyl group, OCH_3 group, SCH_3 group, OCHF_2 group, halogen atom, cyano group or nitro group;

10 R^4 is hydrogen atom, C_1 - C_3 alkyl group, C_1 - C_3 haloalkyl group or halogen atom;

R^5 is hydrogen atom, C_1 - C_3 alkyl group, halogen atom, C_1 - C_3 haloalkyl group, cyclopropyl group, vinyl group, C_2 alkynyl group, cyano group, $\text{C}(\text{O})\text{R}^{38}$ group, CO_2R^{38} group, $\text{C}(\text{O})\text{NR}^{38}\text{R}^{39}$ group, $\text{CR}^{34}\text{R}^{35}\text{CN}$ group, $\text{CR}^{34}\text{R}^{35}\text{C}(\text{O})\text{R}^{38}$ group, 15 $\text{CR}^{34}\text{R}^{35}\text{CO}_2\text{R}^{38}$ group, $\text{CR}^{34}\text{R}^{35}\text{C}(\text{O})\text{NR}^{38}\text{R}^{39}$ group, CHR^{34}OH group, $\text{CHR}^{34}\text{OC}(\text{O})\text{R}^{38}$ group or $\text{OCHR}^{34}\text{OC}(\text{O})\text{NR}^{38}\text{R}^{39}$ group, or, when G is G-2 or G-6, R^4 and R^5 may form $\text{C}=\text{O}$ group together with the carbon atom to which they are attached;

20 R^6 is C_1 - C_6 alkyl group, C_1 - C_6 haloalkyl group, C_2 - C_6 alkoxyalkyl group, C_3 - C_6 alkenyl group or C_3 - C_6 alkynyl group;

X^1 is single bond, oxygen atom, sulfur atom, NH group, $\text{N}(\text{C}_1$ - C_3 alkyl) group, $\text{N}(\text{C}_1$ - C_3 haloalkyl) group or $\text{N}(\text{allyl})$ group;

25 R^7 is hydrogen atom, C_1 - C_6 alkyl group, C_1 - C_6

haloalkyl group, halogen atom, $S(O)_2(C_1-C_6\text{alkyl})$ group or $C(=O)R^{40}$ group;

R^8 is hydrogen atom, C_1-C_8 alkyl group, C_3-C_8 cycloalkyl group, C_3-C_8 alkenyl group, C_3-C_8 alkynyl group, C_1-C_8 haloalkyl group, C_2-C_8 alkoxyalkyl group, C_3-C_8 alkoxyalkoxyalkyl group, C_3-C_8 haloalkynyl group, C_3-C_8 haloalkenyl group, C_1-C_8 alkylsulfonyl group, C_1-C_8 haloalkylsulfonyl group, C_3-C_8 alkoxy-carbonylalkyl group, $S(O)_2NH(C_1-C_8\text{ alkyl})$ group, $C(O)R^{41}$ group or benzyl group whose phenyl ring may be substituted with R^{42} ;

n and m are independently 0, 1, 2 or 3 and $m + n$ is 2 or 3;

Z is CR^9R^{10} group, oxygen atom, sulfur atom, $S(O)$ group, $S(O)_2$ group or $N(C_1-C_4\text{ alkyl})$ group;

each R^9 is independently hydrogen atom, C_1-C_3 alkyl group, halogen atom, hydroxyl group, C_1-C_6 alkoxy group, C_1-C_6 haloalkyl group, C_1-C_6 haloalkoxy group, C_2-C_6 alkylcarbonyloxy group or C_2-C_6 haloalkylcarbonyloxy group;

each R^{10} is independently hydrogen atom, C_1-C_3 alkyl group, hydroxyl group or halogen atom;

R^{11} and R^{12} are independently hydrogen atom, halogen atom, C_1-C_6 alkyl group, C_3-C_6 alkenyl group or C_1-C_6 haloalkyl group;

R^{13} is hydrogen atom, C_1-C_6 alkyl group, C_1-C_6 haloalkyl group, C_3-C_6 alkenyl group, C_3-C_6 haloalkenyl group,

C₃-C₆ alkynyl group, C₃-C₆ haloalkynyl group, HC(=O) group, (C₁-C₄ alkyl)C(=O) group or NH₂ group;

R¹⁴ is C₁-C₆ alkyl group, C₁-C₆ alkylthio group, C₁-C₆ haloalkyl group or N(CH₃)₂ group;

5 W is nitrogen atom or CR¹⁵;

R¹⁵ is hydrogen atom, C₁-C₆ alkyl group, halogen atom, or phenyl group optionally substituted with C₁-C₆ alkyl group, one or two halogen atoms, C₁-C₆ alkoxy group or CF₃ group;

10 each Q is independently oxygen atom or sulfur atom;

Q¹ is oxygen atom or sulfur atom;

Z¹ is CR¹⁶R¹⁷ group, oxygen atom, sulfur atom, S(O) group, S(O)₂ group or N(C₁-C₄alkyl) group;

15 each R¹⁶ is independently hydrogen atom, halogen atom, hydroxyl group, C₁-C₆ alkoxy group, C₁-C₆ haloalkyl group, C₁-C₆ haloalkoxy group, C₂-C₆ alkylcarbonyloxy group or C₂-C₆ haloalkylcarbonyloxy group;

20 each R¹⁷ is independently hydrogen atom, hydroxyl group or halogen atom;

R¹⁸ is C₁-C₆ alkyl group, halogen atom or C₁-C₆ haloalkyl group;

R¹⁹ and R²⁰ are independently hydrogen atom, C₁-C₆ alkyl group, or C₁-C₆ haloalkyl group;

25 Z² is oxygen atom, sulfur atom, NR⁹ group or CR⁹R¹⁰

group;

R^{21} and R^{22} are independently C_1 - C_6 alkyl group, C_1 - C_6 haloalkyl group, C_3 - C_6 alkenyl group, C_3 - C_6 haloalkenyl group, C_3 - C_6 alkynyl group or C_3 - C_6 haloalkynyl group;

5 R^{23} is hydrogen atom, halogen atom or cyano group;

R^{24} is C_1 - C_6 alkylsulfonyl group, C_1 - C_6 alkyl group, C_1 - C_6 haloalkyl group, C_3 - C_6 alkenyl group, C_3 - C_6 alkynyl group, C_1 - C_6 alkoxy group, C_1 - C_6 haloalkoxy group or halogen atom;


R^{25} is C_1 - C_6 alkyl group, C_1 - C_6 haloalkyl group, C_3 - C_6 alkenyl group or C_3 - C_6 alkynyl group;

15 R^{26} is C_1 - C_6 alkyl group, C_1 - C_6 haloalkyl group or phenyl group optionally substituted with C_1 - C_6 alkyl, one or two halogen atoms, one or two nitro groups, C_1 - C_6 alkoxy group or CF_3 group;


W^1 is nitrogen atom or CH group;

T is a group represented by any one of the following general formulas T-1, T-2 and T-3;

20



T-1



T-2



T-3

(wherein E^1 , E^2 , E^3 , E^4 , E^5 , E^6 , E^7 , E^8 , E^9 , E^{10} , E^{11} and E^{12} are

independently hydrogen atom or C₁-C₃ alkyl group);

R²⁷ is C₁-C₈ alkyl group, C₃-C₈ cycloalkyl group, C₃-C₈ alkenyl group, C₃-C₈ alkynyl group, C₁-C₈ haloalkyl group, C₂-C₈ alkoxyalkyl group, C₂-C₈ alkylthioalkyl group, C₂-C₈ alkylsulfanylalkyl group, C₂-C₈ alkylsulfonylalkyl group, C₁-C₈ alkylsulfonyl group, phenylsulfonyl group whose phenyl ring may be substituted with at least one substituent selected from the group consisting of halogen atom and C₁-C₄ alkyl group, C₄-C₈ alkoxyalkoxyalkyl group, C₄-C₈ cycloalkylalkyl group, C₆-C₈ cycloalkoxyalkyl group, C₄-C₈ alkenyloxyalkyl group, C₄-C₈ alkynyloxyalkyl group, C₃-C₈ haloalkoxyalkyl group, C₄-C₈ haloalkenyloxyalkyl group, C₄-C₈ haloalkynyloxyalkyl group, C₆-C₈ cycloalkylthioalkyl group, C₄-C₈ alkenylthioalkyl group, C₄-C₈ alkynylthioalkyl group, C₁-C₄ alkyl group substituted with phenoxy group whose ring is substituted with at least one substituent selected from the group consisting of halogen atom, C₁-C₃ alkyl group and C₁-C₃ haloalkyl group, benzyloxy group whose ring is substituted with at least one substituent selected from the group consisting of halogen atom, C₁-C₃ alkyl group and C₁-C₃ haloalkyl group, C₄-C₈ trialkylsilylalkyl group, C₃-C₈ cyanoalkyl group, C₃-C₈ halocycloalkyl group, C₃-C₈ haloalkenyl group, C₃-C₈ alkoxyalkenyl group, C₅-C₈ haloalkoxyalkenyl group, C₅-C₈ alkylthioalkenyl group, C₃-C₈ haloalkynyl group, C₅-C₈ alkoxyalkynyl group, C₅-C₈

haloalkoxyalkynyl group, C₃-C₈ alkylthioalkynyl group, C₂-C₈ alkylcarbonyl group, benzyl group whose ring is substituted with at least one substituent selected from the group consisting of halogen atom, C₁-C₃ alkyl group and C₁-C₃ haloalkyl group, CHR³⁴COR²⁸ group, CHR³⁴COOR²⁸ group, CHR³⁴P(O)(OR²⁸)₂ group, CHR³⁴P(S)(OR²⁸)₂ group, CHR³⁴C(O)NR²⁹R³⁰ group or CHR³⁴C(O)NH₂ group;

R²⁸ is C₁-C₆ alkyl group, C₂-C₆ alkenyl group, C₃-C₆ alkynyl group or tetrahydrofuranlyl group;

R²⁹ and R³¹ are independently hydrogen atom or C₁-C₄ alkyl group;

R³⁰ and R³² are independently C₁-C₄ alkyl group or phenyl group whose ring may be substituted with at least one substituent selected from the group consisting of halogen atom, C₁-C₃ alkyl group and C₁-C₃ haloalkyl group; or,

R²⁹ and R³⁰ together may form -(CH₂)₅-, -(CH₂)₄- or -CH₂CH₂OCH₂CH₂-, or the ring thus formed may be substituted with at least one substituent selected from the group consisting of C₁-C₃ alkyl group, phenyl group and benzyl group; or,

R³¹ and R³² may form C₃-C₈ cycloalkyl group together with the carbon atom to which they are attached;

R³³ is C₁-C₄ alkyl group, C₁-C₄ haloalkyl group or C₃-C₆ alkenyl group;

R³⁴ and R³⁵ are independently hydrogen atom or C₁-

C₄ alkyl group;

R³⁶ is hydrogen atom, C₁-C₆ alkyl group, C₃-C₆ alkenyl group or C₃-C₆ alkynyl group;

R³⁷ is hydrogen atom, C₁-C₄ alkyl group or halogen atom;

R³⁸ is hydrogen atom, C₁-C₆ alkyl group, C₃-C₆ cycloalkyl group, C₃-C₆ alkenyl group, C₃-C₆ alkynyl group, C₂-C₆ alkoxyalkyl group, C₁-C₆ haloalkyl group, phenyl group whose ring may be substituted with at least one substituent selected from the group consisting of halogen atom, C₁-C₄ alkyl group and C₁-C₄ alkoxy group, -CH₂CO₂(C₁-C₄ alkyl) group or -CH(CH₃)CO₂(C₁-C₄ alkyl) group;


R³⁹ is hydrogen atom, C₁-C₂ alkyl group or C(O)O(C₁-C₄ alkyl) group;

R⁴⁰ is hydrogen atom, C₁-C₆ alkyl group, C₁-C₆ alkoxy group or NH(C₁-C₆ alkyl) group;

R⁴¹ is C₁-C₆ alkyl group, C₁-C₆ haloalkyl group, C₁-C₆ alkoxy group, NH(C₁-C₆ alkyl) group, phenyl group whose ring may be substituted with one substituent selected from the group consisting of R⁴² group, benzyl group and C₂-C₆ dialkylamino group; and

R⁴² is C₁-C₆ alkyl group, one or two halogen atoms, C₁-C₆ alkoxy group or CF₃ group;

(3) a compound of the formula (II):



or nipilacrofen,

wherein R^{43} is C_1 - C_4 alkyl group;

5 R^{44} is C_1 - C_4 alkyl group, C_1 - C_4 alkylthio group, C_1 - C_4 alkoxy group, C_1 - C_4 haloalkyl group, C_1 - C_4 haloalkylthio group or C_1 - C_4 haloalkoxy group;

R^{43} and R^{44} together may form $-(CH_2)_3-$ or $-(CH_2)_4-$;

R^{45} is hydrogen atom or halogen atom;












R^{46} is hydrogen atom or C_1 - C_4 alkyl group;

10 R^{47} is hydrogen atom, nitro group, cyano group, $-COOR^{49}$ group, $-C(=X)NR^{50}R^{51}$ group or $-C(=X^2)R^{52}$ group;

R^{48} is hydrogen atom, halogen atom, cyano group, C_1 - C_4 alkyl group optionally substituted with at least one substituent selected from the group consisting of halogen
15 atom and hydroxyl group, C_1 - C_4 alkoxy group, phenyl group optionally substituted with at least one substituent selected from the group consisting of halogen atom, nitro group, cyano group, C_1 - C_4 alkyl group, C_1 - C_4 alkoxy group and halo- C_1 - C_4 alkyl group, pyrrolyl group, C_2 - C_8 alkyl group,
20 C_3 - C_8 alkenyl group, C_3 - C_8 alkynyl group, C_3 - C_8 alkoxy group, a group selected from the group consisting of C_2 - C_8 alkyl

group, C₃-C₈ alkenyl group, C₃-C₈ alkynyl group and C₃-C₈ alkoxy group into which at least one oxygen atom is inserted, or any one of groups represented by the following formulas:

5



wherein R^{49} , R^{50} and R^{52} are, the same or different, hydrogen atom or C_1-C_4 alkyl group;

R^{50} and R^{51} may form saturated alicyclic 5 or 6 membered ring together with the nitrogen atom to which they are attached;

R^{52} is hydrogen atom, C_1-C_4 alkyl group or C_1-C_4 alkyl group substituted with at least one halogen atom;

R^{53} is hydrogen atom, C_1-C_4 alkyl group optionally substituted with at least one halogen atom, C_2-C_6 alkenyl group optionally substituted with at least one halogen atom, C_3-C_6 alkynyl group optionally substituted with at least one halogen atom, phenyl group optionally substituted with at least one halogen atom, C_3-C_6 cycloalkyl group, cyanomethyl group, or $R^{63}CO-$ group;

R^{54} is hydrogen atom, C_1-C_6 alkyl group optionally substituted with at least one halogen atom, C_2-C_6 alkenyl group optionally substituted with at least one halogen atom, C_3-C_6 alkynyl group optionally substituted with at least one halogen atom, phenyl group optionally substituted with halogen atom, C_3-C_6 cycloalkyl group, cyanomethyl group, C_1-C_4 alkoxy- C_1-C_6 alkyl group, di- C_1-C_4 alkylamino- C_1-C_4 alkyl group, tetrahydrofurfurylmethyl group, C_3-C_6 alkynyloxy- C_1-C_4 alkyl group, benzyl whose ring may be substituted with substituent selected from the group consisting of halogen atom, nitro group, cyano group, C_1-C_4 alkyl group, C_1-C_4

alkoxy group and halo-C₁-C₄ alkyl group, -C(=X²)R⁶³ group, -
(CH₂)_a-(O)_d-R⁷⁰ group, -(CH₂)_a-O-(CH₂)_b-R⁷⁰ group, -(CH₂)_a-X²-R⁷⁶
group;

R⁵³ and R⁵⁴ together with the nitrogen atom to
5 which they are attached may form saturated alicyclic 3, 5
or 6 membered ring or aromatic 5 or 6 membered ring in
which a carbon atom may be optionally replaced with oxygen
atom;

R⁵⁵ is hydrogen atom, C₁-C₄ alkyl group, C₂-C₆
10 alkenyl group or C₃-C₆ alkynyl group, or R⁵⁵ and R⁵⁶ together
may form -(CH₂)_e-;

R⁵⁶ and R⁵⁷ are independently C₁-C₄ alkyl group
optionally substituted with at least one halogen atom, C₂-
C₆ alkenyl group optionally substituted with at least one
15 halogen atom, C₃-C₆ alkynyl optionally substituted with at
least one halogen atom or phenyl group optionally
substituted with at least one halogen atom, hydrogen atom,
C₃-C₆ cycloalkyl group, -XR⁶⁰ group or -NR⁶¹R⁶² group;

R⁵⁸ is hydrogen atom, C₁-C₆ alkyl group, C₂-C₆
20 alkenyl group, C₃-C₆ alkynyl group, C₁-C₄ alkylcarbonyl group,
cyano-C₁-C₃ alkyl group, C₁-C₄ alkoxy carbonyl-C₁-C₄ alkyl
group, di-C₁-C₄ alkoxy carbonyl-C₁-C₄ alkyl group, benzyl
group, C₁-C₄ alkoxy-C₁-C₄ alkynyl group, -(CH₂)_a-R⁷⁵ group, -
(CH₂)_a-X²-R⁷² group, -(CH₂)_a-X²-(CH₂)_b-R⁷² group or -(CH₂)_a-X²-
25 (CH₂)_b-X²-(CH₂)_c-R⁷² group;

R^{59} is hydrogen atom, C_1-C_4 alkyl group, C_2-C_6 alkenyl group, C_3-C_6 alkynyl group, cyano- C_1-C_3 alkyl group, C_1-C_4 alkylcarbonyl- C_1-C_3 alkyl group or phenyl group;

R^{60} is C_1-C_4 alkyl group optionally substituted
5 with at least one halogen atom;

R^{61} and R^{62} are, the same or different, hydrogen atom or C_1-C_4 alkyl group;

R^{63} is C_1-C_4 alkyl group optionally substituted
10 with at least one halogen atom, C_1-C_4 alkoxy- C_1-C_4 alkyl group, C_1-C_4 alkylthio- C_1-C_4 alkyl group, C_3-C_6 cycloalkyl group, phenyl group whose ring may be substituted with one substituent selected from the group consisting of halogen atom, nitro group, cyano group, C_1-C_4 alkyl group, C_1-C_4 alkoxy group and halo- C_1-C_4 alkyl group, $-NR^{73}R^{74}$ group or -
15 $(CH_2)_a-(O)_d-R^{75}$ group;

R^{64} is C_1-C_4 alkoxy carbonyl group or carboxyl group;

R^{65} is chloromethyl group, cyanomethyl group, C_3-C_6 cycloalkyl group into which at least one oxygen atom may
20 be inserted, or C_1-C_4 alkoxy carbonyl- C_1-C_4 alkyl group;

R^{66} is hydroxyl group or $-NR^{67}R^{68}$ group;

A is $-NR^{67}R^{68}$ group or $-S(O)_r-R^{69}$ group;

R^{67} and R^{68} are, the same or different, hydrogen atom or C_1-C_4 alkyl group;

R^{69} is C_1-C_4 alkyl group or C_1-C_4 haloalkyl group;

R^{70} is hydrogen atom, hydroxyl group, halogen atom, C_1-C_4 alkyl group optionally substituted with at least one C_1-C_4 alkoxy group, C_3-C_6 cycloalkyl group into which at least one oxygen atom may be inserted, C_3-C_6 cycloalkyl group optionally substituted with one or two methyl groups, furyl group, thienyl group or $-C(=O)R^{71}$ group;

R^{71} and R^{72} are, the same or different, C_1-C_4 alkyl group or C_1-C_4 alkoxy group;

R^{73} and R^{74} are, the same or different, C_1-C_4 alkyl group or phenyl group;

R^{75} is C_3-C_6 cycloalkyl into which at least one oxygen atom may be inserted, C_3-C_6 cycloalkyl group optionally substituted with one or two methyl groups, furyl group, thienyl group or $-C(=O)R^{71}$ group;

R^{76} is C_1-C_4 alkyl group;

a, b and c is independently 1, 2 or 3;

d is 0 or 1;

e is 2 or 3;

f is 1 or 2; and

X^2 is oxygen atom or sulfur atom.

52. The method according to the above 1, additionally comprising the steps of:

introducing into the plant cell, a second gene selected from a gene encoding a protein substantially having protoporphyrinogen oxidase activity, a gene encoding

a protein substantially having 5-enolpyruvylshikamate-3-phosphate synthase activity and a gene encoding a protein substantially having glyphosate oxidoreductase activity; and

5 expressing said second gene.

53. A plant cell having:

a gene encoding a protein having the following characteristics (a) to (c):

10 (a) having a specific affinity for a substance which is concerned with the weed control activity of a weed control compound,

 (b) having substantially no capability of modifying a substance for which said protein has a specific affinity, and

15 (c) being substantially free from framework regions of variable regions in an immunoglobulin; and

 at least one altered form of an enzymatic activity which gives a resistance to a weed control compound in an amount inhibiting a naturally occurring form of said enzymatic activity, wherein said altered form of an enzymatic activity is a form of enzymatic activity selected from a protoporphyrinogen oxidase activity, 5-enolpyruvylshikamate-3-phosphate synthase activity and glyphosate oxidoreductase activity.

25 54. A plant cell having:

a gene encoding a protein having the following characteristics (a) to (c):

(a) having a specific affinity for a substance which is concerned with the weed control activity of a weed control compound,

(b) having substantially no capability of modifying a substance for which said protein has a specific affinity, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin; and

an altered protoporphyrinogen oxidase activity which gives a resistance to a weed control compound in an amount inhibiting a natural occurring protoporphyrinogen oxidase activity.

55. A plant cell having:

a gene encoding a protein having the following characteristics (a) to (c):

(a) having a specific affinity for a substance which is concerned with the weed control activity of a weed control compound,

(b) having substantially no capability of modifying a substance for which said protein has a specific affinity, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin; and

an altered 5-enolpyruvylshikamate-3-phosphate
synthase activity which gives a resistance to a weed
control compound in an amount inhibiting a natural
occurring 5-enolpyruvylshikamate-3-phosphate synthase
activity.

56. A plant cell having:

a gene encoding a protein having the following
characteristics (a) to (c):

(a) having a specific affinity for a substance
which is concerned with the weed control activity of a weed
control compound,

(b) having substantially no capability of
modifying a substance for which said protein has a specific
affinity, and

(c) being substantially free from framework
regions of variable regions in an immunoglobulin; and

an altered glyphosate oxidoreductase activity
which gives a resistance to a weed control compound in an
amount inhibiting a natural occurring glyphosate
oxidoreductase activity.

57. The plant cell according to the above 53,
wherein said altered form of an enzymatic activity is
conferred by a second gene selected from a gene encoding a
protein substantially having a protoporphyrinogen oxidase
activity, a gene encoding a protein substantially having 5-

enolpyruvylshikamate-3-phosphate synthase activity and a gene encoding a protein substantially having glyphosate oxidoreductase activity.

58. The plant cell according to the above 57,
5 wherein the gene encoding a protein having the following characteristics (a) to (c):

(a) having a specific affinity for a substance which is concerned with the weed control activity of a weed control compound,

10 (b) having substantially no capability of modifying a substance for which said protein has a specific affinity, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin; and

15 the second gene are introduced into the plant cell in the form in that both of said genes are operably ligated to a promoter and a terminator both of which are functional in said plant cell.

59. The plant cell according to the above 57,
20 wherein the protein substantially having a proto-porphyrinogen IX oxidase activity is protoporphyrinogen IX oxidase, the protein substantially having a 5-enolpyruvylshikamate-3-phosphate synthase activity is 5-enolpyruvylshikamate-3-phosphate synthase and the protein
25 substantially having glyphosate oxidoreductase activity is

glyphosate oxidoreductase.

60. The plant cell according to the above 53, wherein the plant cell is derived from dicotyledones or monocotyledones.

5 61. A plant comprising the plant cell of the above 54.

62. A plant comprising the plant cell of the above 55.

63. A plant comprising the plant cell of the above 56.

64. A method for protecting a plant which comprises applying a protoporphyrinogen IX oxidase inhibitory-type compound to a growth area of the plant of the above 61.

65. A method for protecting a plant which comprises applying a protoporphyrinogen IX oxidase inhibitory-type compound and a compound inhibiting 5-enolpyruvylshikamate-3-phosphate synthase to a growth area of the plant of the above 62.

66. A method for protecting a plant which comprises applying a protoporphyrinogen IX oxidase inhibitory-type compound and a compound inhibiting 5-enolpyruvylshikamate-3-phosphate synthase to a growth area of the plant of the above 63.

67. A method for selecting a plant which

comprises applying a protoporphyrinogen IX oxidase inhibitory-type compound to a growth area of the plant of the above 61 and other plants, and selecting either plant on the basis of difference in growth between the plants.

5 68. A method for selecting a plant which comprises applying a protoporphyrinogen IX oxidase inhibitory-type compound and a compound inhibiting 5-enolpyruvylshikamate-3-phosphate synthase to a growth area of the plant of the above 62 and other plants, and
10 selecting either plant on the basis of difference in growth between the plants.

 69. A method for selecting a plant which comprises applying a protoporphyrinogen IX oxidase inhibitory-type compound and a compound inhibiting 5-enolpyruvylshikamate-3-phosphate synthase to a growth area
15 of the plant of the above 63 and other plants, and selecting either plant on the basis of difference in growth between the plants.

20 DETAILED DESCRIPTION OF THE INVENTION

 In the method of the present invention, substances which are concerned with weed control activities of weed control compounds (hereinafter referred to as weed control substances) are those constituting a part of
25 metabolic reaction systems in organisms which are

responsible for weed control activities upon applying the compounds to plants. Examples thereof include weed control compounds themselves, endogenous substances in plants, and the like. Specifically, as such endogenous substances in plants, for example, there are substrates of target enzymes on which weed control compounds act, or precursors or metabolites of the substrates which cause cellular dysfunction upon accumulating in plant cells; substances produced by the above substances in plant cells which cause cellular dysfunction; and the like. More specifically, it has been known that, when a compound having herbicidal activity (hereinafter referred to as herbicidal compound) which inhibits the activity of protoporphyrinogen IX oxidase (EC 1.3.3.4, hereinafter referred to as PPO) is applied to a plant, protoporphyrinogen IX which is the substrate of PPO is accumulated in the plant cells and it is metabolized to form protoporphyrin X, followed by formation of active oxygen in the presence of both protoporphyrin X and light in the cells, which damages cell functions [Junshi MIYAMOTO ed., Atarashii Noyaku no Kagaku (Chemistry of New Agrochemicals), Chapter 3, Section 3.3, p 106 (1993), Hirokawa Shoten, Tokyo]. Thus, protoporphyrinogen IX, protoporphyrin IX and active oxygen in these systems, and the like can be exemplified as these substances.

In the method of the present invention, weed control compounds include compounds having herbicidal activities, plant growth regulator activities, and the like.

Examples of the herbicidal compounds include compounds inhibiting porphyrin biosynthesis, compounds inhibiting electron transfer in photosynthesis, compounds inhibiting carotenoid biosynthesis, compounds inhibiting amino acid biosynthesis, compounds inhibiting lipid biosynthesis, compounds inhibiting cell wall biosynthesis, compounds influencing protein biosynthesis, nucleic acid biosynthesis and cell division, compounds having auxin antagonistic activity, and the like. More specifically, as the compounds inhibiting porphyrin biosynthesis, for example, there are compounds inhibiting PPO activity (PPO inhibitory-type herbicidal compound), and the like. As the compounds inhibiting electron transfer in photosynthesis, for example, there are compounds inhibiting electron transfer of photochemical system I or II, compounds inhibiting 4-hydroxyphenyl pyruvate dioxygenase (EC 1.13.11.27; hereinafter referred to as 4-HPPD) which influences biosynthesis of plastoquinone which transfers electrons, and the like. As the compounds inhibiting carotenoid biosynthesis, for example, there are compounds inhibiting phytoene desaturase (hereinafter referred to as PDS), and the like. As the compounds inhibiting amino acid

biosynthesis, for example, there are compounds inhibiting EPSPS, acetolactate synthase (EC 4.1.3.18; hereinafter referred to as ALS), glutamine synthetase (EC 6.3.1.2; hereinafter referred to as GS), dihydropteroate synthase (EC 2.5.1.15; hereinafter referred to as DHP), and the like. As the compounds inhibiting lipid biosynthesis, for example, there are compounds inhibiting acetyl CoA carboxylase (EC 6.4.1.2; hereinafter referred to as ACC), and the like. As the compounds inhibiting cell wall biosynthesis, for example, there are compounds inhibiting cellulose biosynthesis, and the like. As the compounds influencing protein biosynthesis, nucleic acid biosynthesis or cell division, for example, there are compounds inhibiting formation of microtubules, and the like.


Examples of the compounds having plant growth regulator activities include compounds having antagonistic activities against plant hormones which enhance cell elongation and differentiation, and the like. Specifically, for example, there are 2,4-D, phenoxyalkane carboxylic acid, derivatives of benzoic acid, derivatives of picolinic acid, and the like.

As the above-described PPO inhibitory-type herbicidal compounds, for example, there are the compounds disclosed in Duke, S.O., Rebeiz, C.A., ACS Symposium Series 559, Porphyric Pesticides, Chemistry, Toxicology, and


Pharmaceutical Applications, American Chemical Society, Washington DC (1994), and the like. Specifically, examples thereof include the following compounds:

(1) chlormethoxynil, bifenoxy, chlornitrofen (CNP),
 5 acifluorfen (5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitorobenzoic acid) and its ethyl ester, acifluorfen-sodium, oxyfluorfen (2-chloro-1-(3-ethoxy-4-nitrophenoxy)-4-trifluorobenzene), oxadiazon (3-[2,4-dichloro-5-(1-methylethoxy)phenyl]-5-(1,1-dimethylethyl)-1,3,4-oxdiazol-2-(3H)-one),
 10 2-[4-chloro-2-fluoro-5-(prop-2-ynyloxy)phenyl]-2,3,4,5,6,7-hexahydro-1H-isindol-1,3-dione, chlorphthalim, (N-(4-chlorophenyl)-3,4,5,6-tetrahydrophthalimide), TNPP-ethyl (ethyl 2-[1-(2,3,4-trichlorophenyl)-4-nitropyrazolyl-5-oxy]propionate), or N3-(1-phenylethyl)-2,6-dimethyl-5-propionylnicotinamide;
 15


(2) a compound represented by the general formula: J-G (I), wherein G is a group represented by any one of the following general formulas G-1 to G-9 and J is a group represented by any one of the following general
 20 formulas J-1 to J-30:




G-1




G-2




G-3




G-4




G-5




G-6




G-7




G-8




G-9




J-1




J-2




J-3




J-4




J-5




J-6




J-7




J-8




J-9




J-10




J-11




J-12




J-13




J-14




J-15




J-16




J-17




J-18




J-19




J-20




J-21




J-22




J-23




J-24




J-25




J-26




J-27



J-28



J-29



J-30

wherein the dotted lines in the formulas J-5, J-6,
J-12 and J-24 represent that the left hand ring contains
only single bonds, or one bond in the ring is a double bond
between carbon atoms;

X is oxygen atom or sulfur atom;

Y is oxygen atom or sulfur atom;

R¹ is hydrogen atom or halogen atom;

R² is hydrogen atom, C₁-C₈alkyl group, C₁-C₈
haloalkyl group, halogen atom, OH group, OR²⁷ group, SH
group, S(O)_pR²⁷ group, COR²⁷ group, CO₂R²⁷ group, C(O)SR²⁷
group, C(O)NR²³R³⁰ group, CHO group, CR²⁷=NOR³⁶ group,
CH=CR³⁷CO₂R²⁷ group, CH₂CHR³⁷CO₂R²⁷ group, CO₂N=CR³¹R³² group,
nitro group, cyano group, NHSO₂R³³ group, NHSO₂NHR³³ group,
NR²⁷R³⁸ group, NH₂ group or phenyl group optionally

substituted with one or more and the same or different C_1 - C_4 alkyl groups;

p is 0, 1 or 2;

R^2 is C_1 - C_2 alkyl group, C_1 - C_2 haloalkyl group,
 5 OCH_3 group, SCH_3 group, $OCHF_2$ group, halogen atom, cyano group or nitro group;

R^4 is hydrogen atom, C_1 - C_3 alkyl group, C_1 - C_3 haloalkyl group or halogen atom;

R^5 is hydrogen atom, C_1 - C_3 alkyl group, halogen
 10 atom, C_1 - C_3 haloalkyl group, cyclopropyl group, vinyl group, C_2 alkynyl group, cyano group, $C(O)R^{38}$ group, CO_2R^{38} group, $C(O)NR^{38}R^{39}$ group, $CR^{34}R^{35}CN$ group, $CR^{34}R^{35}C(O)R^{38}$ group, $CR^{34}R^{35}CO_2R^{38}$ group, $CR^{34}R^{35}C(O)NR^{38}R^{39}$ group, $CHR^{34}OH$ group, $CHR^{34}OC(O)R^{38}$ group or $OCHR^{34}OC(O)NR^{38}R^{39}$ group, or, when G is
 15 $G-2$ or $G-6$, R^4 and R^5 may form $C=O$ group together with the carbon atom to which they are attached;

R^6 is C_1 - C_6 alkyl group, C_1 - C_6 haloalkyl group, C_2 - C_6 alkoxyalkyl group, C_3 - C_6 alkenyl group or C_3 - C_6 alkynyl group;

20 X^1 is single bond, oxygen atom, sulfur atom, NH group, $N(C_1-C_3 \text{ alkyl})$ group, $N(C_1-C_3 \text{ haloalkyl})$ group or $N(\text{allyl})$ group;

R^7 is hydrogen atom, C_1 - C_6 alkyl group, C_1 - C_6 haloalkyl group, halogen atom, $S(O)_2(C_1-C_6 \text{ alkyl})$ group or
 25 $C(=O)R^{40}$ group;

R^8 is hydrogen atom, C_1-C_6 alkyl group, C_3-C_6 cycloalkyl group, C_3-C_6 alkenyl group, C_3-C_6 alkynyl group, C_1-C_6 haloalkyl group, C_2-C_6 alkoxyalkyl group, C_3-C_6 alkoxyalkoxyalkyl group, C_3-C_6 haloalkynyl group, C_3-C_6 haloalkenyl group, C_1-C_6 alkylsulfonyl group, C_1-C_6 haloalkylsulfonyl group, C_3-C_6 alkoxycarbonylalkyl group, $S(O)_2NH(C_1-C_6 \text{ alkyl})$ group, $C(O)R^{11}$ group or benzyl group whose phenyl ring may be substituted with R^{12} ;

n and m are independently 0, 1, 2 or 3 and $m + n$ is 2 or 3;

Z is CR^9R^{10} group, oxygen atom, sulfur atom, $S(O)$ group, $S(O)_2$ group or $N(C_1-C_4 \text{ alkyl})$ group;

each R^9 is independently hydrogen atom, C_1-C_3 alkyl group, halogen atom, hydroxyl group, C_1-C_6 alkoxy group, C_1-C_6 haloalkyl group, C_1-C_6 haloalkoxy group, C_2-C_6 alkylcarbonyloxy group or C_2-C_6 haloalkylcarbonyloxy group;

each R^{10} is independently hydrogen atom, C_1-C_3 alkyl group, hydroxyl group or halogen atom;

R^{11} and R^{12} are independently hydrogen atom, halogen atom, C_1-C_6 alkyl group, C_3-C_6 alkenyl group or C_1-C_6 haloalkyl group;

R^{13} is hydrogen atom, C_1-C_6 alkyl group, C_1-C_6 haloalkyl group, C_3-C_6 alkenyl group, C_3-C_6 haloalkenyl group, C_3-C_6 alkynyl group, C_3-C_6 haloalkynyl group, $HC(=O)$ group, $(C_1-C_4 \text{ alkyl})C(=O)$ group or NH_2 group;

R^{14} is C_1-C_6 alkyl group, C_1-C_6 alkylthio group, C_1-C_6 haloalkyl group or $N(CH_3)_2$ group;

W is nitrogen atom or CR^{15} ;

R^{15} is hydrogen atom, C_1-C_6 alkyl group, halogen atom, or phenyl group optionally substituted with C_1-C_6 alkyl group, one or two halogen atoms, C_1-C_6 alkoxy group or CF_3 group;

each Q is independently oxygen atom or sulfur atom;

Q^1 is oxygen atom or sulfur atom;

Z^1 is $CR^{16}R^{17}$ group, oxygen atom, sulfur atom, $S(O)$ group, $S(O)_2$ group or $N(C_1-C_4\text{alkyl})$ group;

each R^{16} is independently hydrogen atom, halogen atom, hydroxyl group, C_1-C_6 alkoxy group, C_1-C_6 haloalkyl group, C_1-C_6 haloalkoxy group, C_2-C_6 alkylcarbonyloxy group or C_2-C_6 haloalkylcarbonyloxy group;

each R^{17} is independently hydrogen atom, hydroxyl group or halogen atom;

R^{18} is C_1-C_6 alkyl group, halogen atom or C_1-C_6 haloalkyl group;

R^{19} and R^{20} are independently hydrogen atom, C_1-C_6 alkyl group, or C_1-C_6 haloalkyl group;

Z^2 is oxygen atom, sulfur atom, NR^9 group or CR^9R^{10} group;

R^{21} and R^{22} are independently C_1-C_6 alkyl group, C_1-

C₆ haloalkyl group, C₃-C₆ alkenyl group, C₃-C₆ haloalkenyl group, C₃-C₆ alkynyl group or C₃-C₆ haloalkynyl group;

R²³ is hydrogen atom, halogen atom or cyano group;


5 R²⁴ is C₁-C₆ alkylsulfonyl group, C₁-C₆ alkyl group, C₁-C₆ haloalkyl group, C₃-C₆ alkenyl group, C₃-C₆ alkynyl group, C₁-C₆ alkoxy group, C₁-C₆ haloalkoxy group or halogen atom;

10 R²⁵ is C₁-C₆ alkyl group, C₁-C₆ haloalkyl group, C₃-C₆ alkenyl group or C₃-C₆ alkynyl group;


R²⁶ is C₁-C₆ alkyl group, C₁-C₆ haloalkyl group or phenyl group optionally substituted with C₁-C₆ alkyl, one or two halogen atoms, one or two nitro groups, C₁-C₆ alkoxy group or CF₃ group;

15 W¹ is nitrogen atom or CH group;

T is a group represented by any one of the following general formulas T-1, T-2 and T-3;



T-1



T-2



T-3

(wherein E¹, E², E³, E⁴, E⁵, E⁶, E⁷, E⁸, E⁹, E¹⁰, E¹¹ and E¹² are
20 independently hydrogen atom or C₁-C₃ alkyl group);

R²⁷ is C₁-C₈ alkyl group, C₃-C₈ cycloalkyl group;

C_3-C_8 alkenyl group, C_3-C_8 alkynyl group, C_1-C_8 haloalkyl group,
 C_2-C_8 alkoxyalkyl group, C_2-C_8 alkylthioalkyl group, C_2-C_8
 alkylsulfinylalkyl group, C_2-C_8 alkylsulfonylalkyl group,
 C_1-C_8 alkylsulfonyl group, phenylsulfonyl group whose phenyl
 5 ring may be substituted with at least one substituent
 selected from the group consisting of halogen atom and C_1-
 C_4 alkyl group, C_4-C_8 alkoxyalkoxyalkyl group, C_4-C_8
 cycloalkylalkyl group, C_6-C_8 cycloalkoxyalkyl group, C_4-C_8
 alkenyloxyalkyl group, C_4-C_8 alkynyloxyalkyl group, C_3-C_8
 10 haloalkoxyalkyl group, C_4-C_8 haloalkenyloxyalkyl group, C_4-C_8
 haloalkynyloxyalkyl group, C_6-C_8 cycloalkylthioalkyl group,
 C_4-C_8 alkenylthioalkyl group, C_4-C_8 alkynylthioalkyl group,
 C_1-C_4 alkyl group substituted with phenoxy group whose ring
 is substituted with at least one substituent selected from
 15 the group consisting of halogen atom, C_1-C_3 alkyl group and
 C_1-C_3 haloalkyl group, benzyloxy group whose ring is
 substituted with at least one substituent selected from the
 group consisting of halogen atom, C_1-C_3 alkyl group and C_1-C_3
 haloalkyl group, C_4-C_8 trialkylsilylalkyl group, C_3-C_8
 20 cyanoalkyl group, C_3-C_8 halocycloalkyl group, C_3-C_8
 haloalkenyl group, C_5-C_8 alkoxyalkenyl group, C_5-C_8
 haloalkoxyalkenyl group, C_5-C_8 alkylthioalkenyl group, C_3-C_8
 haloalkynyl group, C_5-C_8 alkoxyalkynyl group, C_5-C_8
 haloalkoxyalkynyl group, C_5-C_8 alkylthioalkynyl group, C_2-C_8
 25 alkylcarbonyl group, benzyl group whose ring is substituted

with at least one substituent selected from the group consisting of halogen atom, C_1-C_3 alkyl group and C_1-C_3 haloalkyl group, $CHR^{34}COR^{28}$ group, $CHR^{34}COOR^{28}$ group, $CHR^{34}P(O)(OR^{28})_2$ group, $CHR^{34}P(S)(OR^{28})_2$ group, $CHR^{34}C(O)NR^{29}R^{30}$ group or $CHR^{34}C(O)NH_2$ group;

R^{28} is C_1-C_6 alkyl group, C_2-C_6 alkenyl group, C_3-C_6 alkynyl group or tetrahydrofuranlyl group;

R^{29} and R^{31} are independently hydrogen atom or C_1-C_4 alkyl group;

R^{30} and R^{32} are independently C_1-C_4 alkyl group or phenyl group whose ring may be substituted with at least one substituent selected from the group consisting of halogen atom, C_1-C_3 alkyl group and C_1-C_3 haloalkyl group; or,

R^{29} and R^{30} together may form $-(CH_2)_5-$, $-(CH_2)_4-$ or $-CH_2CH_2OCH_2CH_2-$, or the ring thus formed may be substituted with at least one substituent selected from the group consisting of C_1-C_3 alkyl group, phenyl group and benzyl group; or,

R^{31} and R^{32} may form C_3-C_6 cycloalkyl group together with the carbon atom to which they are attached;

R^{33} is C_1-C_4 alkyl group, C_1-C_4 haloalkyl group or C_3-C_6 alkenyl group;

R^{34} and R^{35} are independently hydrogen atom or C_1-C_4 alkyl group;

R^{36} is hydrogen atom, C_1-C_6 alkyl group, C_3-C_6

alkenyl group or C₃-C₆ alkynyl group;

R³⁷ is hydrogen atom, C₁-C₄ alkyl group or halogen atom;

R³⁸ is hydrogen atom, C₁-C₆ alkyl group, C₃-C₆ cycloalkyl group, C₃-C₆ alkenyl group, C₃-C₆ alkynyl group, C₂-C₆ alkoxyalkyl group, C₁-C₆ haloalkyl group, phenyl group whose ring may be substituted with at least one substituent selected from the group consisting of halogen atom, C₁-C₄ alkyl group and C₁-C₄ alkoxy group, -CH₂CO₂(C₁-C₄ alkyl) group or -CH(CH₃)CO₂(C₁-C₄ alkyl) group;


R³⁹ is hydrogen atom, C₁-C₂ alkyl group or C(O)O(C₁-C₄ alkyl) group;

R⁴⁰ is hydrogen atom, C₁-C₆ alkyl group, C₁-C₆ alkoxy group or NH(C₁-C₆ alkyl) group;

R⁴¹ is C₁-C₆ alkyl group, C₁-C₆ haloalkyl group, C₁-C₆ alkoxy group, NH(C₁-C₆ alkyl) group, phenyl group whose ring may be substituted with one substituent selected from the group consisting of R⁴² group, benzyl group and C₂-C₈ dialkylamino group; and

R⁴² is C₁-C₆ alkyl group, one or two halogen atoms, C₁-C₆ alkoxy group or CF₃ group;

(3) a compound of the formula (II):



or nipilacrofen,

wherein R^{43} is C_1-C_4 alkyl group;

R^{44} is C_1-C_4 alkyl group, C_1-C_4 alkylthio group, C_1-C_4 alkoxy group, C_1-C_4 haloalkyl group, C_1-C_4 haloalkylthio group or C_1-C_4 haloalkoxy group;






R^{43} and R^{44} together may form $-(CH_2)_3-$ or $-(CH_2)_4-$;

R^{45} is hydrogen atom or halogen atom;

R^{46} is hydrogen atom or C_1-C_4 alkyl group;

R^{47} is hydrogen atom, nitro group, cyano group, $-COOR^{49}$ group, $-C(=X)NR^{50}R^{51}$ group or $-C(=X^2)R^{52}$ group;

R^{48} is hydrogen atom, halogen atom, cyano group, C_1-C_4 alkyl group optionally substituted with at least one substituent selected from the group consisting of halogen atom and hydroxyl group, C_1-C_4 alkoxy group, phenyl group optionally substituted with at least one substituent selected from the group consisting of halogen atom, nitro group, cyano group, C_1-C_4 alkyl group, C_1-C_4 alkoxy group and halo- C_1-C_4 alkyl group, pyrrolyl group, C_2-C_8 alkyl group, C_3-C_8 alkenyl group, C_3-C_8 alkynyl group, C_3-C_8 alkoxy group, a group selected from the group consisting of C_2-C_8 alkyl group, C_3-C_8 alkenyl group, C_3-C_8 alkynyl group and C_3-C_8 alkoxy group into which at least one oxygen atom is inserted, or any one of groups represented by the following formulas:



wherein R^{49} , R^{50} and R^{52} are, the same or different, hydrogen atom or $\text{C}_1\text{-C}_4$ alkyl group;

R^{50} and R^{51} may form saturated alicyclic 5 or 6 membered ring together with the nitrogen atom to which they are attached;

R^{52} is hydrogen atom, $\text{C}_1\text{-C}_4$ alkyl group or $\text{C}_1\text{-C}_4$ alkyl group substituted with at least one halogen atom;

R^{53} is hydrogen atom, $\text{C}_1\text{-C}_4$ alkyl group optionally

substituted with at least one halogen atom, C₂-C₆ alkenyl group optionally substituted with at least one halogen atom, C₃-C₆ alkynyl group optionally substituted with at least one halogen atom, phenyl group optionally substituted with at least one halogen atom, C₃-C₈ cycloalkyl group, cyanomethyl group, or R⁶³CO- group;

R⁵⁴ is hydrogen atom, C₁-C₆ alkyl group optionally substituted with at least one halogen atom, C₂-C₆ alkenyl group optionally substituted with at least one halogen atom, C₃-C₆ alkynyl group optionally substituted with at least one halogen atom, phenyl group optionally substituted with halogen atom, C₃-C₈ cycloalkyl group, cyanomethyl group, C₁-C₄ alkoxy-C₁-C₆ alkyl group, di-C₁-C₄ alkylamino-C₁-C₄ alkyl group, tetrahydrofurfurylmethyl group, C₃-C₆ alkynyloxy-C₁-C₄ alkyl group, benzyl whose ring may be substituted with substituent selected from the group consisting of halogen atom, nitro group, cyano group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group and halo-C₁-C₄ alkyl group, -C(=X²)R⁶³ group, -(CH₂)_a-(O)_d-R⁷⁰ group, -(CH₂)_a-O-(CH₂)_b-R⁷⁰ group, -(CH₂)_a-X²-R⁷⁶ group;

R⁵³ and R⁵⁴ together with the nitrogen atom to which they are attached may form saturated alicyclic 3, 5 or 6 membered ring or aromatic 5 or 6 membered ring in which a carbon atom may be optionally replaced with oxygen atom;

R^{55} is hydrogen atom, C_1-C_4 alkyl group, C_2-C_6 alkenyl group or C_3-C_6 alkynyl group, or R^{55} and R^{56} together may form $-(CH_2)_a-$;

R^{56} and R^{57} are independently C_1-C_4 alkyl group
 5 optionally substituted with at least one halogen atom, C_2-C_6 alkenyl group optionally substituted with at least one halogen atom, C_3-C_6 alkynyl optionally substituted with at least one halogen atom or phenyl group optionally substituted with at least one halogen atom, hydrogen atom,
 10 C_3-C_6 cycloalkyl group, $-XR^{60}$ group or $-NR^{61}R^{62}$ group;

R^{58} is hydrogen atom, C_1-C_6 alkyl group, C_2-C_6 alkenyl group, C_3-C_6 alkynyl group, C_1-C_4 alkylcarbonyl group, cyano- C_1-C_3 alkyl group, C_1-C_4 alkoxy carbonyl- C_1-C_4 alkyl group, di- C_1-C_4 alkoxy carbonyl- C_1-C_4 alkyl group, benzyl
 15 group, C_1-C_4 alkoxy- C_1-C_4 alkynyl group, $-(CH_2)_a-R^{75}$ group, $-(CH_2)_a-X^2-R^{72}$ group, $-(CH_2)_a-X^2-(CH_2)_b-R^{72}$ group or $-(CH_2)_a-X^2-(CH_2)_b-X^2-(CH_2)_c-R^{72}$ group;

R^{59} is hydrogen atom, C_1-C_4 alkyl group, C_2-C_6 alkenyl group, C_3-C_6 alkynyl group, cyano- C_1-C_3 alkyl group,
 20 C_1-C_4 alkylcarbonyl- C_1-C_3 alkyl group or phenyl group;

R^{60} is C_1-C_4 alkyl group optionally substituted with at least one halogen atom;

R^{61} and R^{62} are, the same or different, hydrogen atom or C_1-C_4 alkyl group;

25 R^{63} is C_1-C_4 alkyl group optionally substituted

with at least one halogen atom, C₁-C₄ alkoxy-C₁-C₄ alkyl group, C₁-C₄ alkylthio-C₁-C₄ alkyl group, C₃-C₆ cycloalkyl group, phenyl group whose ring may be substituted with one substituent selected from the group consisting of halogen atom, nitro group, cyano group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group and halo-C₁-C₄ alkyl group, -NR⁷³R⁷⁴ group or - (CH₂)_a-(O)_d-R⁷⁵ group;

R⁶⁴ is C₁-C₄ alkoxycarbonyl group or carboxyl group;

R⁶⁵ is chloromethyl group, cyanomethyl group, C₃-C₆ cycloalkyl group into which at least one oxygen atom may be inserted, or C₁-C₄ alkoxycarbonyl-C₁-C₄ alkyl group;

R⁶⁶ is hydroxyl group or -NR⁶⁷R⁶⁸ group;

A is -NR⁶⁷R⁶⁸ group or -S(O)_z-R⁶⁹ group;

R⁶⁷ and R⁶⁸ are, the same or different, hydrogen atom or C₁-C₄ alkyl group;

R⁶⁹ is C₁-C₄ alkyl group or C₁-C₄ haloalkyl group;

R⁷⁰ is hydrogen atom, hydroxyl group, halogen atom, C₁-C₄ alkyl group optionally substituted with at least one C₁-C₄ alkoxy group, C₃-C₆ cycloalkyl group into which at least one oxygen atom may be inserted, C₃-C₆ cycloalkyl group optionally substituted with one or two methyl groups, furyl group, thienyl group or -C(=O)R⁷¹ group;

R⁷¹ and R⁷² are, the same or different, C₁-C₄ alkyl group or C₁-C₄ alkoxy group;

R^{73} and R^{74} are, the same or different, C_1 - C_4 alkyl group or phenyl group;

R^{75} is C_3 - C_6 cycloalkyl into which at least one oxygen atom may be inserted, C_3 - C_6 cycloalkyl group
5 optionally substituted with one or two methyl groups, furyl group, thienyl group or $-C(=O)R^{71}$ group;

R^{76} is C_1 - C_4 alkyl group;

a, b and c is independently 1, 2 or 3;

d is 0 or 1;

10 e is 2 or 3;

f is 1 or 2; and

X^2 is oxygen atom or sulfur atom.


In addition, as other N-substituted pyrazoles,
there are the 3-substituted-2-aryl-4,5,6,7-tetrahydro-
15 indazoles described in Lyga et al., Pesticide Sci., 42: p
29 (1994), and the like.

As specific examples of the compounds inhibiting
electron transfer in photochemical system I, for example,
there are paraquat, diquat, and the like. As specific
20 examples of the compounds inhibiting electron transfer in
photochemical system II, for example, there are triazine
compounds (e.g., atrazine, etc.), urea compounds (e.g.,
diuron, etc.), nitrile compounds (e.g., bromoxynil and
ioxynil) and the like. As specific examples of the
25 compounds inhibiting 4-HPPD, for example, there are


isoxazoles (e.g., isoxaflutole), pyrazoles, triketones, and the like. As specific examples of the compounds inhibiting PDS, for example, there are norflurazon, flurochloridone, fluridone, flurtamone, diflufenican, and the like. As
5 specific examples of the compounds inhibiting EPSPS, for example, there are glyphosate, and the like. As specific examples of the compounds inhibiting ALS, for example, there are sulfonylureas, imidazolinones, pyrimidinylthiobenzoates, triazolopyrimidines, and the like.
10 As specific examples of the compounds inhibiting GS, for example, there are bialaphos, glufosinate, and the like. As specific examples of the compounds inhibiting DHP, for example, there are asulam, and the like. As specific examples of the compounds inhibiting ACC, for example,
15 there are cyclohexanediones, aryloxyphenoxypropionates, and the like. As specific examples of the compounds inhibiting cellulose, for example, there are dichlobenil, and the like.

Various examples of the weed control compounds useful in the present invention are shown by the following
20 chemical structures:


Structure 1




Structure 2




Structure 3




Structure 4




Structure 5




Structure 6




Structure 7




Structure 8




Structure 9




Structure 10




Structure 11




Structure 12




Structure 13




Structure 14




Structure 15




Structure 16




Structure 17




Structure 18




Structure 19




Structure 20




Structure 21




Structure 22




Structure 23




Structure 24




Structure 25




Structure 26




Structure 27




Structure 28




Structure 29




Structure 30




Structure 31




Structure 32




Structure 33




Structure 34




Structure 35



Structure 36



Structure 37



In the first aspect of the method of the present invention, the genes to be used are those encoding proteins having the following characteristics (a) to (c) (hereinafter sometimes referred to as the objective proteins):

(a) having a specific affinity for weed control substances;

(b) having substantially no capability of modifying substances for which said protein has a specific affinity; and

(c) being substantially free from framework regions of variable regions of an immunoglobulin.

The term "a specific affinity" for weed control substances of the above characteristic (a) means that an enzyme (the objective protein) and a substrate (the weed control substance), or an enzyme (the objective protein) and an inhibitor or a regulator of an activity of the enzyme (the weed control substance) bind to each other, enzymatically; or that the objective protein and the weed control substance bind to each other on the basis of affinity and specificity, such as those shown in a receptor-chemical bond, for example, a bond between a receptor and a ligand, and the like. The objective proteins may be naturally occurring proteins; variants thereof obtained by introduction of amino acid substitution,

addition, deletion, modification and the like into naturally occurring proteins; and artificially synthesized proteins having random amino acid sequences selected with the guidance of their affinity for weed control substances, in so far as they have structures specifically binding to weed control substances.

The term "having substantially no capability of modifying" in the characteristic (b) means that enzymatic reactivity with substances for which said protein has a specific affinity is substantially inactive or not existed (except the specific affinity for weed control substances in the characteristic (a)). Examples of this include a case that the objective protein does not have any capability of converting a substance for which said protein has a specific affinity such as a certain weed control substance or a substance having an essential part of the structure of the substrates on the basis of a specific affinity for said protein, and the like to a substance having a chemical structure different from that of the substance for which said protein has a specific affinity. The protein "having substantially no capability of modifying" can be, for example, identified by checking non-recovery of the growth of a microorganism whose gene encoding the said protein is deleted and thus cannot grow under a usual condition in a case where the gene encoding

the said protein is introduced into the microorganism in such a state that the introduced gene is expressed in the microorganism.

The term "substantially free from the framework regions of variable regions of an immunoglobulin" in the characteristic (c) mean that the objective protein does not form a stereostructure specific for the variable regions of an immunoglobulin. The term "framework regions of variable regions of an immunoglobulin" mean regions remaining after removing the hypervariable regions from the variable regions of H chain and L chain which are the constituents of the immunoglobulin molecule. In these regions, conservation of the amino acid sequences is relatively high and these regions function for maintaining the highly conserved stereostructure of the variable regions. Due to formation of the above stereostructure, the hypervariable regions separately located at three sites on respective H chain and L chain are collected to one site on the stereostructure to form an antigen binding site [Alberts, B., et al. ed. (1983), Molecular Biology of the Cell, p 979, Garland Publishing, Inc., New York].

The objective protein having the above characteristic (c) can be selected on the basis of, for example, the amino acid sequences of the proteins. As specific examples of the protein, there are a protein which

does not contain any amino acid sequence composed of about 30 amino acids or more and having about 60% or more homology with the known amino acid sequences of the framework regions of the variable regions of an immunoglobulin, and the like. For example, the presence or absence of the above framework regions can be confirmed by PCR using a gene encoding the protein as a template and DNAs having nucleotide sequences encoding the variable regions derived from H chain or L chain of the immunoglobulin as amplification primers, for example, the primers VH1BACK and VH1FOR-2, or VK2BACK and VK4FOR described by Clackson, T. et al., Nature 352; p 624 (1991), or primers contained in a commercially available kit for cloning recombinant antibody genes, for example, Heavy primer mix or Light primer mix of Recombinant Phage Antibody System (Pharmacia Biotech) to analyze presence or absence of amplification of DNA having a given length. Examples of the binding proteins having a specific affinity for weed control substances also include peptides having an affinity for the weed control substances.

Specific examples of the objective proteins having the above characteristics of (a) to (c) include inactive-type binding proteins having an affinity for protoporphyrin IX [e.g., inactive-type magnesium chelatase whose substrate is protoporphyrin IX (the weed control

substance), inactive-type ferrochelatase (protoheme ferrolyase; EC 4.9.9.1), inactive-type cobalt chelatase which catalyzes a chelating reaction of a cobalt ion with a compound having tetrapyrrole ring as a substrate, peptides
5 having an affinity for protoporphyrin IX, i.e., proteins composed of 4 to 100 amino acids (for example, peptide HASYS having an affinity for protoporphyrin IX, e.g., a protein comprising the amino acid sequence of SEQ ID NO: 53 and a protein having the amino acid sequence of SEQ ID NO:
10 54; peptide RASSL having an affinity for protoporphyrin IX, e.g., a protein comprising the amino acid sequence of SEQ ID NO: 55 and a protein having the amino acid sequence of SEQ ID NO: 56; peptide YAGY having an affinity for porphyrin compounds, e.g., a protein comprising the amino
15 acid sequence of SEQ ID NO: 57 and a protein having the amino acid sequence of SEQ ID NO: 58; peptide YAGF having affinity for porphyrin compounds, e.g., a protein comprising the amino acid sequence of SEQ ID NO: 59 and a protein having the amino acid sequence of SEQ ID NO: 60; and the like)], inactive-type binding proteins having an
20 affinity for protoporphyrinogen IX (e.g., inactive-type PPO, inactive-type coproporphyrinogen III oxidase), and the like.

The above inactive-type binding proteins include variants thereof whose activities have been lost by amino
25 acid substitution, addition, deletion, modification and the

like of naturally occurring active proteins under natural or artificial conditions.

Cellular dysfunction caused by weed control substances can be prevented by binding of these binding proteins to the weed control substances in plant cells to exhibit the desired weed control compound-resistance.

The inactive-type magnesium chelatase is protoporphyrin IX binding subunit protein of magnesium chelatase, or its variant having a specific affinity for protoporphyrin IX, and specific examples thereof include the subunit protein from which its organelle transit signal sequence has been deleted, and the like.

The inactive-type ferrochelatase is its variant having no capability of modifying protoporphyrin IX and having a specific affinity for protoporphyrin IX, and specific examples thereof include a ferrochelatase variant in which a region presumed to be a Fe ion binding site of ferrochelatase has been modified, and the like.

The inactive-type cobalt chelatase is a substrate binding subunit protein of cobalt chelatase, or its variant having no capability of modifying protoporphyrin IX and having a specific affinity for protoporphyrin IX.

The inactive-type PPO is its variant having no capability of oxidizing protoporphyrinogen IX and having a specific affinity for protoporphyrinogen IX, and specific

examples thereof include a PPO variant in which a region presumed to be FAD binding site of PPO (a region having the amino acid sequence GXGXXG wherein X is any amino acid, e.g., a region comprising the 63rd to 68th amino acids from the N-terminus of chloroplast localized PPO of mouse-ear cress (*Arabidopsis thaliana*) and having the amino acid sequence of GGGISG) has been deleted, and the like.

The inactive-type coproporphyrinogen III oxidase is its variant having no capability of oxidizing protoporphyrinogen IX and having a specific affinity for protoporphyrinogen IX.

The genes encoding the above proteins can be obtained by, for example, as follows.

As the genes encoding protoporphyrin IX binding subunit protein of magnesium chelatase, for example, those derived from the photosynthetic bacterium, *Rhodobacter capsulatus* (Genebank accession M74001), mouse-ear cress (Genebank accession Z68495), barley (Genebank accession U96216), snapdragon (*Antirrhinum majus*) (Genebank accession U26916), *Synechocystis* P.C.C. 6803 (Genebank accession U29131) and the like have been known. For isolating such a known gene (its nucleotide sequence has been known), PCR can be carried out by using genomic DNA or cDNA of an organism having the desired gene as a template and primers produced on the basis of nucleotide sequences corresponding

to those about the N- and C-termini of the protein encoded by the gene to amplify the desired gene. Further, genes encoding protoporphyrin IX binding subunit protein of magnesium chelatase can be obtained from photosynthetic organisms other than the above. For example, first, a cDNA library is constructed by obtaining mRNA from the desired photosynthetic organism, synthesizing cDNA by using the mRNA as a template with a reverse transcriptase, and integrating the cDNA into a phage vector such as ZAPII, etc. or a plasmid vector such as pUC, etc. For amplifying a DNA fragment containing at least a part of the gene encoding protoporphyrin IX binding subunit protein of magnesium chelatase, PCR can be carried out by using the above-constructed cDNA library as a template and primers designed and synthesized on the basis of nucleotide sequences well conserved among known genes such as the above-described genes. Screening of the cDNA library can be carried out by using the DNA fragment thus obtained as a probe to select positive clones. The desired gene of protoporphyrin IX binding subunit protein of magnesium chelatase can be confirmed by sequence determination of the nucleotide sequence of the selected clone.

For obtaining the gene encoding a variant of protoporphyrin IX binding subunit protein of magnesium chelatase having a specific affinity for protoporphyrin IX,

for example, the gene encoding the subunit protein is mutagenized by introduction of nucleotide substitution, addition, deletion, modification and the like, followed by introducing the resultant gene into *Escherichia coli* BL21(DE3) strain according to the method described by Gibson, L.C. D. et al., Proc. Natl. Acad. Sci. USA, 92; p 1941 (1995) and the like to obtain transformants, and culturing the transformants under conditions that high expression of the gene thus introduced occurs. The desired gene encoding a variant of the subunit protein having a specific affinity for protoporphyrin IX can be obtained by selecting a strain whose cultured cells have turned red and have the fluorescence absorption showing accumulation of protoporphyrin IX (excitation wavelength 405 nm, emission wavelength 630 nm).

As the genes encoding ferrochelatase, for example, those derived from *Escherichia coli* (Genebank accession D90259), *Bacillus subtilis* (Genebank accession M97208), *Bradyrhizobium japonicum* (Genebank accession M92427), yeast *Saccharomyces cerevisiae* (Genebank accession J05395), mouse (Genebank accession J05697), human being (Genebank accession D00726), barley (Genebank accession D26105), cucumber (Genebank accession D26106), and the like have been known. For isolating such a known gene (its nucleotide sequence has been known), PCR can be carried out

by using genomic DNA or cDNA of an organism having the desired gene as a template and primers produced on the basis of nucleotide sequences corresponding to those about the N- and C-termini of the protein encoded by the gene to amplify the desired gene. Further, for obtaining other genes encoding ferrochelatase, for example, first, a cDNA library is constructed by obtaining mRNA from the desired organism, synthesizing cDNA by using the mRNA as a template with a reverse transcriptase, and integrating the cDNA into a phage vector such as ZAPII, etc. or a plasmid vector such as pUC, etc. The cDNA library can be introduced into ferrochelatase deficient mutant strain of *Escherichia coli* VS200 described by Miyamoto, K, et al., Plant Physiol., 105; p 769 (1994), followed by subjecting a complementation test to select clones containing ferrochelatase gene derived from the desired organism. Further, for amplifying a DNA fragment, PCR can be carried out by using the above-constructed cDNA library as a template and primers prepared on the basis of nucleotide sequences well conserved among known genes such as the above-described genes. Screening of the cDNA library can be carried out by using the DNA fragment thus obtained as a probe to select positive clones. The desired ferrochelatase gene can be confirmed by sequence determination of the nucleotide sequence of the selected clone.

For obtaining the gene encoding a variant of ferrochelatase having no capability of modifying protoporphyrin IX and having a specific affinity for protoporphyrin IX (for example, the gene encoding a ferrochelatase variant in which the region presumed to be a Fe ion binding site of ferrochelatase is modified), PCR can be carried out by preparing a mutagenesis primer for introduction of mutation into the region on the basis of nucleotide sequence encoding the amino acid sequence about the region, and using a commercially available site-directed mutagenesis kit (Mutan-Super Express, Takara Shuzo) to obtain the gene encoding the above variant. Specifically, a wild type ferrochelatase gene is inserted into the cloning site of plasmid vector pKF19K and PCR is carried out by using the resultant plasmid DNA as a template, the above-described mutagenesis primer and a selection primer for restoration of amber mutation located on kanamycin resistant gene of pKF19K. The gene amplified by PCR is introduced into *Escherichia coli* MV1184 (suppressor free strain) and the transformants are screened according to kanamycin resistance to isolate *Escherichia coli* having ferrochelatase gene in which the nucleotide sequence corresponding to the amino acid sequence which constitutes the desired region has been modified. The isolated gene can be confirmed as the gene encoding the

desired protein by analyzing the nucleotide sequence of the plasmid DNA of the *Escherichia coli*.

The genes encoding the peptides having an affinity for protoporphyrin IX, i.e., the proteins composed of 4 to 100 amino acids can be obtained by synthesizing a peptide library according to, for example, the combinatorial chemistry method as described by Sugimoto, N., Nakano, S., Chem., Lett., p 939 (1997) and the like, selecting a peptide having an affinity for the weed control substance, analyzing the amino acid sequence of the peptide thus selected with a peptide sequencer, designing a gene containing a nucleotide sequence encoding the amino acid sequence, and synthesizing the nucleotide sequence with a DNA synthesizer or the like.

Further, a phase clone displaying a peptide having an affinity for the weed control substance can be selected from a phage library according to phage display method. Specifically, for example, a phage library displaying a protein having a random amino acid sequence on the surface of M13 phage particles is constructed by inserting a nucleotide sequence encoding the protein having the random amino acid sequence into the upstream from the region encoding the coat protein pIII of M13 phage gene. On the other hand, the weed control substance labeled with biotin is bound to a plate coated with avidin or

streptoavidin to prepare a support coated with the weed control substance. A phage displaying the desired protein having an affinity for the weed control substance can be isolated by screening the above phage library on the plate coated with the weed control substance and the gene of the desired protein can be obtained from the isolated phage.

The gene encoding a protein containing the repetition of the amino acid sequence represented by SEQ ID NO: 53, 55, 57 or 59 four times or eight times can be produced by, for example, selecting a nucleotide sequence in which the nucleotide sequence encoding the above amino acid sequence is repeated the given times after the initiation codon ATG, synthesizing an oligonucleotide comprising the selected nucleotide sequence and an oligonucleotide comprising a nucleotide sequence complementary to the selected nucleotide sequence by a DNA synthesizer, and then subjecting them to annealing. Further, the genes encoding the amino acid sequence represented by SEQ ID NO: 54, 56, 58 or 60 can be produced by selecting a nucleotide sequence encoding the amino acid sequence, synthesizing an oligonucleotide comprising the selected nucleotide sequence and another oligonucleotide comprising a nucleotide sequence complementary to the selected nucleotide sequence by a DNA synthesizer, and then subjecting them to annealing. In this respect, for

selecting the nucleotide sequence encoding the given amino acid sequence, for example, it is preferred to select codons frequently used in genes derived from plants.

As PPO genes, for example, those derived from

5 *Escherichia coli* (Genebank accession X68660), *Bacillus subtilis* (Genebank accession M97208), *Haemophilus influenzae* (Genebank accession L42023), mouse (Genebank accession D45185), human being (Genebank accession D38537), mouse-ear cross (Genebank accession D83139), tobacco

10 (Genebank accession Y13465, Y13466) and the like have been known. For isolating such a known gene (its nucleotide sequence has been known), PCR is carried out by using genomic DNA or cDNA of an organism having the desired gene as a template and primers produced on the basis of

15 nucleotide sequences corresponding to those about the N- and C-termini of the protein encoded by the gene to amplify the desired gene. Further, for obtaining other PPO genes, for example, first, a cDNA library is constructed from an organism having the desired gene according to the above-

20 described method. The cDNA library can be introduced into *Escherichia coli* PPO deficient mutant strain VSR800 described by Narita, S., et al., Gene, 182; p 169 (1996), followed by subjecting a complementation test to select clones containing PPO gene derived from the desired

25 organism. Further, for amplifying a DNA fragment, PCR can

be carried out by using the above-constructed cDNA library as a template and primers prepared on the basis of nucleotide sequences well conserved among known genes such as the above-described genes. Screening of the cDNA library can be carried out by using the DNA fragment thus obtained as a probe to select positive clones. The desired PPO gene can be confirmed by sequence determination of the nucleotide sequence of the selected clone.

For obtaining the gene encoding a variant of PPO having no capability of oxidizing protoporphyrinogen IX and having a specific affinity for protoporphyrinogen IX, for example, PPO gene is mutagenized by introducing nucleotide substitution, addition, deletion, modification, etc. and the resultant modified gene is introduced into the above *Escherichia coli* whose growth is inhibited light-dependently by treatment with a PPO inhibitory-type herbicidal compound. A gene encoding a protein having protoporphyrinogen IX binding capability can be selected by culturing the *Escherichia coli* thus obtained in the presence of hemin, aminolevulinic acid and a PPO inhibitory-type herbicidal compound to select a clone which can grow even in the light. A gene encoding a protein having no capability of oxidizing protoporphyrinogen IX can be selected by expressing the modified gene thus selected in a host such as *Escherichia coli*, etc. to prepare a

protein encoded by the gene, and measuring its capability of oxidizing protoporphyrinogen IX according to the method described by Jacobs, N.J. and Jacobs, J.M. (1982) *Enzyme*, 28, 206-219 and the like. More specifically, the above

5 modified gene is inserted into an expression vector for *Escherichia coli* and introduced into PPO gene (hemG locus) deficient mutant of *Escherichia coli* such as *Escherichia coli* BT3 strain described by Yamamoto, F., et al., *Japanese J. Genet.*, 63; p 237 (1988) and the like. The *Escherichia coli*

10 *coli* is cultured in a culture medium containing hemin and aminolevulinic acid in addition to the cell growth inhibitor corresponding to the selection marker of the vector introduced into the *Escherichia coli* to obtain transformants. The protein encoded by the modified gene

15 can be produced from the transformant. Further, a gene which does not complement PPO gene deficiency of its host cell can be obtained by culturing the transformant in a culture medium substantially free from hemin and aminolevulinic acid to identify a strain which does not

20 grow. This latter method can also be used for selection of the gene encoding a protein having no capability of oxidizing protoporphyrinogen IX.

Further, for obtaining the gene encoding a variant of PPO in which the region presumed to be a FAD

25 binding site of PPO (the region having the amino acid

sequence GXGXXG, wherein X is any amino acid) is deleted, first, a mutagenesis primer for introduction of deletion mutation of the region is prepared on the basis of the nucleotide sequence encoding the amino acid sequence about the region. Then, PCR is carried out by using the mutagenesis primer and a commercially available site-directed mutagenesis kit (Mutan-Super Express, Takara Shuzo) as described above to obtain the gene encoding the above variant protein in which the region has been deleted.

The genes encoding peptide proteins such as the peptides HASYS and RASSL having an affinity for protoporphyrin IX, and the peptides YAGA and YAGF having an affinity for prophyrin compounds, and the like can be obtained by subjecting oligonucleotides synthesized by a DNA synthesizer to annealing.

Furthermore, genes encoding unknown peptide proteins having affinities for other weed control substances can be produced by the following methods and the like. For example, various peptide libraries can be constructed according to, for example, the combinatorial chemistry method as described by Sugimoto, N., Nakano, S., Chem., Lett., p 939 (1997), and the like. Peptides are selected from the peptide libraries thus constructed with the guidance of affinities for weed control substances, followed by analyzing the amino acid sequences of the

peptides with a peptide sequencer. Thus, genes encoding the peptides can be synthesized by a DNA synthesizer. Alternatively, phase clones displaying peptides having affinities for weed control substances can be obtained by selecting phage libraries according to phage display method. Specifically, for example, a phage library displaying a protein having a random amino acid sequence on the surface of M13 phage particles is constructed by inserting a nucleotide sequence encoding the protein having the random amino acid sequence into the upstream from the region encoding the coat protein pIII of M13 phage gene. On the other hand, a weed control substance labeled with biotin is bound to a plate coated with avidin or streptoavidin to prepare a support coated with the weed control substance. A phage displaying the desired protein having an affinity for the weed control substance can be isolated by screening the above phage library on the plate coated with the weed control substance and the gene of the desired protein can be obtained from the isolated phage.

As the genes encoding coproporphyrinogen III oxidase, for example, those derived from *Escherichia coli* (Genebank accession X75413), *Salmonella typhimurium* (Genebank accession L19503), yeast *Saccharomyces cerevisiae* (Genebank accession J03873), mouse (Genebank accession D1633), human being (Genebank accession D16333), soybean

(Genebank accession X71083), barley (Genebank accession X82830), tobacco (Genebank accession X82831) and the like have been known. For isolating such a known gene (its nucleotide sequence has been known), PCR is carried out by using genomic DNA or cDNA of an organism having the desired gene as a template and primers produced on the basis of nucleotide sequences corresponding to those about the N- and C-termini of the protein encoded by the gene to amplify the desired gene. Further, for obtaining other coproporphyrinogen III oxidase genes, for example, first, a cDNA library is constructed from an organism having the desired gene by preparing mRNA from the desired organism, synthesizing cDNA using the mRNA as a template with a reverse transcriptase and integrating this into a plasmid vector such as pRS313 described by Sikorski, R.S., et al., Genetics, 122; p 19 (1989), and the like. The cDNA library can be introduced into yeast coproporphyrinogen III oxidase deficient mutant strain HEM13 described by Troup, B., et al., Bacteriol., 176; p 673 (1994), followed by subjecting a complementation test to select clones containing coproporphyrinogen III oxidase derived from the desired organism. Further, for amplifying a DNA fragment, PCR can be carried out by using the above-constructed cDNA library as a template and primers prepared on the basis of nucleotide sequences well conserved among known genes such

as the above-described genes. Screening of the cDNA library can be carried out by using the DNA fragment thus obtained as a probe to select positive clones. The desired coproporphyrinogen III oxidase gene can be confirmed by
5 sequence determination of the nucleotide sequence of the selected clone.

For obtaining the gene encoding a variant of coporphyrinogen III oxidase having no capability of oxidizing protoporphyrinogen IX and having a specific
10 affinity for protoporphyrinogen IX, for example, coproporphyrinogen III oxidase gene is mutagenized by introducing nucleotide substitution, addition, deletion, modification, etc. and the resultant gene is introduced into the above *Escherichia coli* whose growth is inhibited
15 light-dependently by treatment with a PPO inhibitory-type herbicidal compound. A gene encoding a protein having protoporphyrinogen IX binding capability can be selected by culturing the *Escherichia coli* thus obtained in the presence of hemin, aminolevulinic acid and a PPO
20 inhibitory-type herbicide to select a clone which can grow even in the light. A gene encoding a protein having no capability of oxidizing protoporphyrinogen IX can be selected by expressing the modified gene thus selected in a host such as *Escherichia coli*, etc. to prepare a protein
25 encoded by the gene, and measuring its capability of

oxidizing protoporphyrinogen IX according to the method described by Jacobs, N.J. and Jacobs, J.M. (1982) Enzyme, 28, 206-219 and the like.

The genes which is used in the second aspect of the method of the present invention are those encoding proteins having the following characteristics (a) to (c):

(a) having a specific affinity for protoporphyrin IX;

(b) having substantially no capability of modifying protoporphyrinogen IX; and

(c) being substantially free from framework regions of variable regions of immunoglobulins.

The term "a specific affinity" for protoporphyrin IX in the characteristic (a) is substantially the same as that in the above first aspect of the method of the present invention and means that the protein and protoporphyrin IX bind to each other, enzymatically or the protein and protoporphyrin IX bind to each other on the basis of affinity and specificity as those shown in receptor chemical bond such as a bond between a receptor and a ligand and the like. The proteins may be naturally occurring proteins; variants thereof in which amino acid substitution, addition, deletion, modification and the like are introduced into naturally occurring proteins; and artificially synthesized proteins having random amino acid

sequences which are selected with the guidance of an affinity for protoporphyrin IX in so far as they have structures specifically binding to protoporphyrin IX.

The term "having substantially no capability of modifying" protoporphyrinogen IX in the characteristic (b) means that enzymatic reactivity with protoporphyrinogen IX of the protein is substantially inactive or not existed. For example, this means that the protein does not have capability of converting protoporphyrinogen IX into a substance having a chemical structure different from that of protoporphyrinogen IX.

The term "substantially free from framework regions of variable regions of immunoglobulins" means the same as that in the above first aspect of the method of the present invention and the protein does not form the stereostructure specific for the variable regions in the immunoglobulin as is described hereinabove.

As specific examples of the proteins having the above characteristics (a) to (c), there are active or inactive-type binding proteins having an affinity for protoporphyrin IX [e.g., active or inactive-type magnesium chelatase whose substrate is protoporphyrin IX, active or inactive-type ferrochelatase, active or inactive-type cobalt chelatase which catalyzes a chelating reaction of a cobalt ion with a compound having tetrapyrrole ring as a

substrate, peptides, i.e., proteins composed of 4 to 100 amino acids, having an affinity for protoporphyrin IX (for example, proteins containing at least one peptide selected from peptide HASYS having an affinity for protoporphyrin IX, e.g., a protein comprising the amino acid sequence of SEQ ID NO: 53 and a protein having the amino acid sequence of SEQ ID NO: 54; peptide RASSL having an affinity for protoporphyrin IX, i.e., a protein comprising the amino acid sequence of SEQ ID NO: 55 and a protein having the amino acid sequence of SEQ ID NO: 56; peptide YAGY having an affinity for porphyrin compounds, e.g., a protein comprising the amino acid sequence of SEQ ID NO: 57 and a protein having the amino acid sequence of SEQ ID NO: 58; peptide YAGF having affinity for porphyrin compounds, i.e., a protein comprising the amino acid sequence of SEQ ID NO: 59 and a protein having the amino acid sequence of SEQ ID NO: 60; and the like)], and the like.

The genes encoding the above proteins can be obtained by, for example, as follows.

Active-type magnesium chelataes are composed of three heterogenous subunit proteins, i.e., protoporphyrin IX binding subunit protein (H subunit protein), I subunit protein and D subunit protein, all of them are essential for catalytic acitivity. Three independent subunit proteins are encoded by different genes. The genes of

protoporphyrin IX binding subunit protein can be obtained by PCR or screening of cDNA library as described hereinabove.

As the gene encoding I subunit protein of a magnesium chelatase, for example, those derived from photosynthetic bacterium, *Rhodobacter sphaeroides* (Genebank accession AF017642), *Rhodobacter capsulatus* (Genebank accession Z11165), *Arabidopsis* (Genebank accession D49426), barley (Genebank accession U26545), soybean (Genebank accession D45857), tobacco (Genebank accession AF14053), *Synechocystis* P.C.C.6803 (Genebank accession U35144) and the like have been known. For isolating such a known gene (its nucleotide sequence has been known), PCR can be carried out by using genomic DNA or cDNA of an organism having the desired gene as a template and primers produced on the basis of nucleotide sequences corresponding to those about the N- and C-termini of the protein encoded by the desired gene. Further, genes encoding I subunit protein of a magnesium chelatase can be obtained from photosynthetic organisms other than the above. For example, first, a cDNA library is constructed by obtaining mRNA from the desired photosynthetic organisms, synthesizing cDNA by using the mRNA as a template with a reverse transcriptase, and integrating the cDNA into a phage vector such as ZAPII, etc. or plasmid vector such as pUC, etc. For amplifying a DNA

fragment containing at least a part of the gene encoding I subunit protein of a magnesium chelatase, PCR can be carried out by using the above-constructed cDNA library as a template and primers designed and synthesized on the basis of nucleotide sequences well conserved among known genes such as the above described genes. Screening of the cDNA library can be carried out by using the DNA fragment thus obtained as a probe to select positive clones. The desired gene of I subunit protein of a magnesium chelatase can be confirmed by determination of the nucleotide sequence of the selected clone.

As the gene encoding D subunit protein of a magnesium chelatase, for example, those derived from photosynthetic bacterium, *Rhodobacter sphaeroides* (Genebank accession AJ001690), *Rhodobacter capsulatus* (Geneband accession Z11165), pea (Genebank accession AF014399), tobacco (Genebank accession Y10022), *Synechocystis* P.C.C.6803 (Genebank accession X96599) and the like have been known. The isolation of such a known gene (its nucleotide sequence has been known) or genes other than the above can be carried out in the same manner as described in that of the gene encoding I subunit protein of magnesium chelatase.

The genes used in the third aspect of the method of the present invention are those encoding proteins having

the following characteristics (a) to (c):

(a) having a specific affinity for protoporphyrinogen IX;

5 (b) having the capability of modifying coproporphyrinogen III; and

(c) being substantially free from framework regions of variable regions of immunoglobulins.

10 The term "a specific affinity" for protoporphyrinogen IX in the characteristic (a) is substantially the same as that in the above first or second aspect of the method of the present invention and means that the protein and protoporphyrinogen IX bind to each other, enzymatically or the protein and protoporphyrinogen IX are bound to each other on the basis of affinity and
15 specificity as those shown in receptor-chemical bond such as a bond between a receptor and a ligand and the like. The proteins may be naturally occurring proteins; variants thereof in which amino acid substitution, addition, deletion, modification and the like are introduced into
20 naturally occurring proteins; and artificially synthesized proteins having random amino acid sequences which are selected with the guidance of an affinity for protoporphyrinogen IX in so far as they have structures specifically binding to protoporphyrinogen IX.

25 The term "having the capability of modifying"

coproporphyrinogen III in the characteristic (b) means that enzymatical reactivity with coproporphyrinogen III of the proteins is active. For example, this means that the protein has the capability of converting coproporphyrinogen
 5 III into a substance having a chemical structure different from that of coproporphyrinogen III.

The term "substantially free from framework regions of variable regions of immunoglobulins" means the same as that in the above first or second aspect of the
 10 method of the present invention and the protein does not form the stereostructure specific for the variable regions in the immunoglobulin as is described hereinabove.

As specific examples of the proteins having the above characteristics (a) to (c), there are active or
 15 inactive-type binding proteins having an affinity for proporphyrinogen IX, for example, active-type coproporphyrinogen III oxidase whose substrate is proporphyrinogen IX, and the like.

As a reference, the activity of a magnesium
 20 chelatase, a ferrochelatase or a coproporphyrinogen III oxidase is, for example, measured by using the following method.

(1) A magnesium chelatase:

The genes encoding independent three subunit
 25 proteins are used to detect a magnesium chelatase activity

according to the method by Gibson, L.C.D., et al. (Proc. Natl. Acad. Sci. USA, 92; p 1941 (1995)) and the like.

(2) A ferrochelatase:

A ferrochelatase activity can, for example, be
5 detected according to the method by Porra, R.J. (Anal. Biochem., 68; p 289 (1975)) and the like.

(3) A coproporphyrinogen III oxidase:

A coproporphyrinogen III oxidase activity can,
for example, be detected according to the method by
10 Yoshinaga, T., Sano, S., et al. (J. Biol. Chem., 255; p 4722 (1980)) and the like.

In the fourth aspect of the method of the present invention, there may be used in addition to the gene encoding the protein having the characteristics (a) to (c)
15 (as described in the first to third aspects of the present invention), at least one altered form of an enzymatic activity selected from an altered PPO activity, an altered EPSPS activity and an altered glyphosate oxidoreductase (GOX) activity. Said altered form of the enzymatic
20 activity in the plant cell can give a resistance to a weed control compound in an amount inhibiting a naturally occurring form of said enzymatic activity. Typically, such an amount of the weed control compound is an amount which can set forth a herbicidal control over the growth of a
25 plant cell, which by inhibiting a naturally occurring form

of the enzymatic activity. In this regard, to give the resistance to a PPO inhibitory-type herbicidal compound, it is preferable that the plant cell additionally comprises the altered PPO activity. Likewise, to give the resistance to glyphosate, it is preferable that the plant cell additionally comprises the altered EPSPS activity or the altered GOX activity.

Glyphosate is the common name given to the weed control compound, N-(phosponomethyl)glycine. In this regard, glyphosate includes the ammonium salt, sodium salt, isopropylamine salt, trimethylsulfonium salt, potassium salt or the like salt. Further, glyphosate is a compound encompassed by said compound inhibiting EPSPS, as described above.

The term "altered form of an enzymatic activity" means that the enzymatic activity is different from that which naturally occurs in a plant cell, which altered form of an enzymatic activity provides a resistance to a weed control compound that inhibits the naturally occurring activity thereof. Said naturally occurring enzymatic activity in the plant cell is the enzymatic activity which occurs naturally in the absence of direct or indirect manipulation by man of such naturally occurring enzymatic activity..

A second gene in the plant cell is useful to

confer said altered form of the enzymatic activity therein. As such, the second gene typically provides in the plant cell, a gene providing for the altered PPO activity, for the altered EPSPS activity or for the altered GOX activity.

5 Various proteins can be encoded by the second gene, so that the second gene can provide for said altered form of enzymatic activity when expressed in the plant cell.

10 In utilizing the second gene for the altered PPO activity, the second gene can encode a naturally occurring protein substantially having PPO activity. In the plant cell, such a protein substantially having PPO activity can be a protein having a capability of oxidizing protoporphyrinogen IX and which has a specific affinity for protoporphyrinogen IX.

15 In its amino acid sequence, the protein substantially having PPO activity preferably contains said region presumed to be FAD binding site of PPO. Such a protein substantially having PPO activity may be PPO. As genes encoding PPO, there can be utilized the known "PPO genes" as described above. Further, there can be utilized
20 a naturally occurring protein substantially having PPO activity which activity is resistant to the PPO inhibitory-type herbicidal compound (as described in EP 0770682 or WO 9833927)

25 Further, the protein substantially having PPO

activity can have substituted, deleted or added thereto amino acids, such that the resulting variant protein has substantially the PPO activity. Conventional methods well known in the art can be used to substitute, delete or add the amino acids thereto. USP 5939602 and WO 9704089 describe variant PPO substantially having PPO activity which activity is uninhibited by a PPO inhibitory-type herbicidal compound. The second gene may encode such a variant PPO.

In utilizing the second gene for the altered EPSPS activity, the second gene can encode a naturally occurring protein substantially having EPSPS activity. Such a protein substantially having EPSPS activity is a protein having a capability to modify in the plant cell, phosphoenolpyruvic acid (PEP) with 3-phosphoshikimic acid to 5-enolpyruvyl-3-phosphoshikimic acid.

Such a protein substantially having EPSPS activity may be EPSPS. As a polynucleotide encoding EPSPS, there can be utilized a known polynucleotide encoding EPSPS.

Examples of such a polynucleotide encoding EPSPS include those derived from *Petunia hybrida* (Genebank accession M37029), Mitchell diploid petunia (as described in EP218571), *Salmonella typhimurium* (as described in EP508909), Tomato (strain VF36) pistil (Genebank accession M21071), *Arabidopsis thaliana* (Genebank accession X06613),

soy beans, *Zea mays* (Genebank accession X63374),
Escherichia coli (Genebank accession X00557), *Agrobacterium*
tumefaciens sp.strain CP4 (class II) and the like.

Additionally, the second gene can also encode a naturally
5 occurring protein substantially having EPSPS activity which
activity is resistant to glyphosate, such as a bacterial
EPSPS which activity is resistant to glyphosate.

Further, the second gene when providing for the
altered EPSPS activity in the plant cell can also encode a
10 variant protein substantially having EPSPS activity. As
such, the protein substantially having EPSPS activity can
have substituted, deleted or added thereto amino acids such
that the resulting protein substantially has the EPSPS
activity. Examples of such a variant protein substantially
15 having EPSPS activity include a variant EPSPS which
activity is resistant to glyphosate, a variant EPSPS in
which a chloroplast transit peptide is added thereto and
the like.

The variant EPSPS which activity is resistant to
20 glyphosate can be produced by substituting, deleting or
adding nucleotides to a gene encoding EPSPS. For example,
a substitutive mutation can be introduced to a
polynucleotide encoding EPSPS to produce a variant
polynucleotide. The protein encoded by the resulting
25 variant gene can then be confirmed for a resistance to

glyphosate and for the EPSPS activity.

The resistance to glyphosate may be confirmed by introducing the variant gene to a particular *Escherichia coli* mutant and by culturing the resulting particular *Escherichia coli* mutant in a specified minimal nutrient MOPS medium which has glyphosate added thereto. As the particular variant *Escherichia coli* in this case, there is utilized an *Escherichia coli* mutant which is deficient in its endogenous EPSPS gene (aroA locus) and which has the growth thereof inhibited in the specified minimal nutrient MOPS medium (in which there is no glyphosate therein). Further, in this case, the specified minimal nutrient MOPS medium is specified in that there is no aromatic amino acids present therein. When glyphosate is added to the minimal nutrient MOPS medium, the glyphosate is in an amount which would typically inhibit in normal growing conditions, the growth of an *Escherichia coli* mutant which is deficient in its endogenous EPSPS gene but which has introduced thereto a naturally occurring gene encoding herbicidally sensitive EPSPS. By selecting the resulting clones growable in such specified minimal nutrient MOPS medium containing glyphosate, there can be obtained a variant polynucleotide encoding a variant EPSPS having an activity which is resistant to glyphosate. The EPSPS activity can be confirmed by introducing said variant gene

to a host cell and then by according to the method described in EP 409815. In this regard, there can be obtained the second gene encoding the variant EPSPS substantially having EPSPS activity which activity is
5 resistant to glyphosate.

In utilizing the second gene for the altered GOX activity, the second gene can encode a naturally occurring protein substantially having GOX activity. Such a protein substantially having the GOX activity is a protein having a
10 capability to degrade glyphosate to less herbicidal products, such as aminomethyl phosphonate (AMPA) and glyoxylate. In cases in which glyphosate is degraded into AMPA and glyoxylate, for example, the protein substantially having GOX activity may cleave the C-N bond
15 of glyphosate.

Such a protein substantially having GOX activity may be GOX. As a polynucleotide encoding GOX, there can be utilized a known polynucleotide encoding GOX. Examples of naturally occurring GOX genes include those derived from
20 *Pseudomonas* sp.strains LBAA, *Pseudomonas* sp.strains LBr, *Agrobacterium* sp.strain T10 and the like.

Further, the second gene when providing for the altered GOX activity in the plant cell can encode a variant protein substantially having GOX activity. As such, the
25 protein substantially having GOX activity can have

substituted, deleted or added thereto amino acids such that the resulting protein substantially has the GOX activity. As an example of such a variant protein substantially having GOX activity, there is mentioned a variant GOX in which a chloroplast transit peptide is added thereto. Conventional methods well known in the art can be used to substitute, delete or add the amino acids thereto.

The GOX activity of a protein substantially having GOX activity can be confirmed by introducing a gene encoding the protein substantially having GOX activity into a specified *Escherichia coli* mutant and by culturing the resulting specified *Escherichia coli* mutant in a minimal nutrient MOPS medium containing glyphosate as the sole nitrogenous source therein. As the specified *Escherichia coli* mutant in this case, there is utilized an *Escherichia coli* mutant which can grow in a minimal nutrient MOPS medium having a non-herbicidal aminophosphate compound as the sole nitrogenous source therein, such as *E.coli* SR2000 Mpu+ . The glyphosate therein is in an amount which would typically inhibit the growth of the specified *Escherichia coli* mutant having no said gene encoding the protein substantially having GOX activity introduced thereto. By selecting the resulting clones growable in such minimal nutrient MOPS medium containing glyphosate as the sole nitrogenous source therein, there can be obtained a

polynucleotide encoding a protein substantially having GOX activity. Such results suggest that in cases in which the protein substantially having GOX activity degrades glyphosate into AMPA, the growable clones use to grow, AMPA as a nitrogenous source. In practice, a 3-¹⁴C labeled glyphosate may be used to confirm that said growable clone consumes and degrades glyphosate. For example, the growable clone may be cultured with the 3-¹⁴C labeled glyphosate and the cell extract thereof may then be analyzed with HPLC.

The second gene encoding an above protein substantially having PPO activity, EPSPS activity or GOX activity can be obtained, for example, as follows.

For isolating a known gene encoding a protein substantially having PPO activity, EPSPS activity or GOX activity, PCR can be carried out by using genomic DNA or cDNA of an organism having the desired gene as a template and primers produced on the basis of nucleotide sequences corresponding to those about the N- and C-termini of the protein to amplify the desired gene. Further, genes encoding a protein substantially having PPO activity, EPSPS activity or GOX activity can be obtained from organisms other than the above. For example, first, a cDNA library is constructed by obtaining mRNA from an organism and synthesizing cDNA by using the mRNA as template with

reverse transcriptase and integrating the cDNA into a phage vector such as ZAP II, etc. or a plasmid vector such as pUC, etc. For the protein substantially having PPO activity, the cDNA library may be introduced into *Escherichia coli* PPO deficient mutant strain VSR800 described by Narita, S., et al., *Gene*, 182; p 169 (1996), followed by subjecting a complementation test to select clones containing PPO gene derived from the desired organism. Further, for amplifying a DNA fragment containing at least a part of the desired gene, PCR can be carried out by using the above-constructed cDNA library as a template and primers designed and synthesized on the basis of nucleotide sequences well conserved among known genes such as the above-described genes. Screening of the cDNA library can be carried out by using the DNA fragment thus obtained as a probe to select positive clones. The desired gene, i.e., a gene encoding the protein substantially having the PPO activity, EPSPS activity or GOX activity, can be confirmed by determination of the nucleotide sequence of the selected clone.

Examples of methods used to confer the altered EPSPS activity or altered GOX activity include the following. An example may include a method of introducing into a cultivated plant a gene having a polynucleotide sequence encoding a petunia (Mitchell diploid petunia)

EPSPS downstream of a high expression promoter such as a cauliflower mosaic virus 35S promoter (EP 218571). A further example may include a method of introducing into a cultivated plant a gene having a 35S promoter upstream of a polynucleotide sequence encoding an *Agrobacterium* (*Agrobacterium tumefaciens* sp. strain CP4) EPSPS fused with a chloroplast transit peptide of a petunia (*Petunia hybrida*) EPSPS (WO 9204449, USP 5633435). A furthermore example may include a method of introducing into a cultivated plant a gene having 2 continuous 35S promoters upstream a polynucleotide encoding a sunflower chloroplast transit peptide of small subunit of ribulose-1,5-bisphosphate carboxylase (ssRUBISCO), the 22 amino acids from the N-terminus of maize ssRUBISCO, maize chloroplast transit peptide of ssRUBISCO and a *Salmonella* (*Salmonella typhimurium*) EPSPS (EP 508909). Even furthermore, an example may include a method of introducing into a cultivated plant, a gene having downstream from a promoter of *Arabidopsis thaliana* alcohol dehydrogenase A, a polynucleotide encoding an *Arabidopsis thaliana* chloroplast transit peptide and GOX (WO 9706269). Yet even furthermore, an example may include a method of introducing into a cultivated plant, the above gene encoding GOX as well as a gene having downstream from a 35S promoter possessing an enhanced promoter activity with the omega sequence of tobacco mosaic virus, a

polynucleotide sequence which encodes an *Agrobacterium* (*Agrobacterium tumefaciens* sp. strain CP4) EPSPS (class II) downstream a chloroplast transit peptide of *Petunia* (*Petunia hybrida*) EPSPS (WO 9706269). Still yet even
5 furthermore, an example may include a method of introducing into a cultivated plant, a gene encoding a variant EPSPS having amino acid substitutions therein which augment the resistant to glyphosate [Hinchee, M.A.W. et al., BIO/TECHNOLOGY, 6: p915 (1988), EP 389066, EP 409815, WO
10 9206201 and USP 5312910].

Examples of methods used to confer the altered PPO activity include the following. An example may include a method of over-expressing in a plant cell, a gene encoding wild-type, naturally occurring PPO (USP 5767373).
15 A further example may include a method of expressing in a plant cell, a variant protein substantially having PPO activity which activity is not inhibited by a PPO inhibitory-type herbicidal compound (USP 5939602). A
furthermore example may include a method of expressing a
20 PPO substantially having PPO activity which is not inhibited by a PPO inhibitory-type herbicidal compound, wherein the PPO is derived from bacteria (EP 0770682 or WO 9833927).

In the method (including the above first to third
25 aspects) of the present invention, for introducing the gene

encoding the protein having the characteristics of (a) to (c) into a plant cell, a gene encoding one protein can be introduced. Further, plural genes encoding different proteins can be introduced into a plant cell. When said altered form of enzymatic activity is given to the plant cell, the second gene encoding one protein may also be introduced. Further, plural genes of the second gene can be introduced into the plant cell to provide for said altered form of enzymatic activity therein. In introducing the gene encoding the protein having the characteristics of (a) to (c) and second gene thereto, the gene encoding the protein having the characteristics of (a) to (c) may be introduced into the plant cell with the second gene, or may be introduced before or after the second gene is introduced to the plant cell. Such gene introduction into plant cells can be carried out by conventional gene engineering techniques, for example, *Agrobacterium* infection (JP-B 2-58917 and JP-A 60-70070), electroporation into protoplasts (JP-A 60-251887 and JP-A 5-68575), particle gun methods (JP-A 5-508316 and JP-A 63-258525), and the like.

Preferably, the gene to be introduced into a plant cell is integrated into a vector having a selection marker gene such as a gene which can give cell growth inhibitor resistance to the plant cell. For example, the gene encoding the protein having the characteristics of (a)

to (c) and the second gene, when utilized for the altered form of enzymatic activity, can be integrated into one of such vectors. Further, the gene encoding the protein having the characteristics of (a) to (c) and the second gene may also each be integrated, respectively, into such vectors having a selection marker gene. In integrating the gene encoding the protein having the characteristics (a) to (c) and the second gene into such respective vectors, the selection marker gene utilized for the vector for the second gene is typically different from the selection marker gene utilized for the vector for the gene encoding the protein having the characteristics (a) to (c).

For expression of the gene encoding the protein having the characteristics (a) to (c) in the plant cell, the gene can be introduced into a chromosome of a plant cell by homologous recombination [Fraley, R.T. et al., Proc. Natl. Acad. Sci. USA, 80; p 4803 (1983)] to select the plant cell expressing the gene. Alternatively, the gene can be introduced into a plant cell in the form that it is operably ligated to a promoter and a terminator both of which can function in the plant cell.

The term "operably ligated" used herein means that the above promoter and terminator are joined in such a state that the introduced gene is expressed in the plant cell under control of the promoter and the terminator.

To provide for the altered form of enzymatic activity, the second gene is expressed in a plant cell. For expression of the second gene in the plant cell, the second gene can likewise be introduced into a chromosome of a plant cell by homologous recombination to select the plant cell expressing the second gene. Alternatively, the second gene can be introduced into a plant cell in the form that it is operably ligated to a promoter and a terminator both of which can function in the plant cell. When utilized, the second gene is typically expressed at a level such that the amount of the protein encoded by the second gene provides for the altered form of enzymatic activity and further confer the resistance of the plant cell. It is preferable when the second gene encodes PPO or EPSPS, that the second gene provide for the altered form of enzymatic activity through over-expression. If so desired, a transcriptionally strong promoter which can function in the plant cell can be utilized with the second gene.

As the promoter which can function in a plant cell, for example, there are constitutive promoters derived from T-DNA such as nopaline synthase gene promoter, octopine synthase gene promoter, etc., promoters derived from plant viruses such as 19S and 35S promoters derived from cauliflower mosaic virus, etc., inductive promoters such as phenylalanine ammonia-lyase gene promoter, chalcone

synthase gene promoter, pathogenesis-related protein gene promoter, etc., and the like. The promoter is not limited these promoters and other plant promoters can be used.

As the terminator which can function in a plant cell, for example, there are terminators derived from T-DNA such as nopaline synthase terminator, etc., terminators derived from plant viruses such as terminators derived from garlic viruses GV1, GV2, etc., and the like. The terminator is not limited to these terminators and other plant terminators can be used.

As the plant cells into which the gene encoding the protein having the characteristics of (a) to (c) are introduced, for example, there are plant tissues, whole plants, cultured cells, seeds and the like. Examples of the plant species into which the genes are introduced include dicotyledones such as tobacco, cotton, rapeseed, sugar beet, mouse-ear cress, canola, flax, sunflower, potato, alfalfa, lettuce, banana, soybean, pea, legume, pine, poplar, apple, grape, citrus fruits, nuts, etc.; and monocotyledones such as corn, rice, wheat, barley, rye, oat, sorghum, sugar cane, lawn, etc. The second gene may also be introduced into such plant cells.

The transformant plant cells expressing the gene encoding the protein having the characteristics of (a) to (c) can be obtained by culturing cells into which the gene

is transferred in a selection culture medium corresponding to a selection marker joined to the locus on the gene, for example, a culture medium containing a cell growth inhibitor, or the like, and isolating a clone capable of growing in the culture medium. Further, the selection culture medium should also correspond to a selection marker joined to the locus of the second gene when the altered form of enzymatic activity is also present in the transformant plant cells. Alternatively, the above transformant plant cells can be selected by culturing plant cells into which the gene is introduced in a culture medium containing the weed control compound to which the resistance is given, and isolating clones capable of growing in the culture medium.

The desired weed control compound-resistant plant can be obtained from the transformant cells thus obtained by regenerating the whole plant according to a conventional plant cell culture method, for example, that described in Plant Gene Manipulation Manual, Method for Producing Transgenic Plants, UCHIMIYA, Kodansha Scientific (1996). Thus, the transformed plants such as plant tissues, whole plants, cultured cells, seeds and the like can be obtained.

For example, rice and mouse-ear cress expressing the gene encoding the protein having the characteristics of

(a) to (c) can be obtained according to the method

described Experimental Protocol of Model Plants, Rice and Mouse-Ear Cress Edition, (Supervisors: Koh SHIMAMOTO and Kiyotaka OKADA, Shujun-sha, 1996), Chapter 4. Further, according to the method described in JP-A 3-291501, soybean expressing the gene encoding the binding protein by introducing the gene into soybean adventitious embryo with a particle gun. Likewise, according to the method described by Fromm, M.E., et al., Bio/Technology, 8; p 838 (1990), corn expressing the gene encoding the above protein can be obtained by introducing the gene into adventitious embryo with a particle gun. Wheat expressing the gene encoding the above protein can be obtained by introducing the gene into sterile-cultured wheat immature scutellum with a particle gun according to a conventional method described by TAKUMI et al., Journal of Breeding Society (1995), 44: Extra Vol. 1, p 57. Likewise, according to a conventional method described by HAGIO, et al., Journal of Breeding Society (1995), 44; Extra Vol. 1, p 67, barley expressing the gene encoding the above protein can be obtained by introducing the gene into sterile-cultured barley immature scutellum with a particle gun.

For confirmation of weed control compound-resistance of the plant expressing the gene encoding the above protein, preferably, the plant is reproduced with applying the weed control compound to which resistance is

given to evaluate the degree of reproduction of the plant. For more quantitative confirmation, for example, in case of resistance to a compound having PPO inhibitory-type herbicidal activity, preferably, pieces of leaves of the
5 plant are dipped in aqueous solutions containing the compound having PPO inhibitory-type herbicidal activity at various concentrations, or the aqueous solutions containing the compound having herbicidal activity are sprayed on pieces of leaves of the plant, followed by allowing to
10 stand on an agar medium in the light at room temperature. After several days, chlorophyll is extracted from the plant leaves according to the method described by Mackenney, G., J. Biol. Chem., 140; p 315 (1941) to determine the content of chlorophyll.

15 Since the weed control compound-resistant plants (e.g., plant tissues, whole plants, cultured cells, seeds, etc.) obtained by the method of the present invention (including the first to fourth aspects) show resistance to weed control compounds, even in case that a weed control
20 compound is applied to a growth area (e.g., cultivation area, proliferation area, etc.), the plant can grow. Therefore, when a weed control compound is applied to a growth area of the desired weed control compound resistant-plant, the desired plant can be protected from plants
25 without resistance to the weed control plant. For example,

weeds can be controlled efficiently by applying a weed control compound on a growth area of the plant having resistance to the weed control compound.

Further, by applying a weed control compound to a growth area of the weed control compound-resistant plant obtained by the method of the present invention (including the first to third aspects) and other plants (e.g., those having no or weak resistance to the weed control compound), one of the plants can be selected on the basis of the difference in growth between the plants. For example, by applying (adding) a weed control compound to a cultivation area (culture medium) of the weed control compound-resistant plant cells obtained by the method of the present invention and other plant cells (e.g., those having no or weak resistance to the weed control compound), one of the plant cells can be selected efficiently on the basis of the difference in growth between the plants.

The following Examples further illustrate the present invention in detail but are not to be construed to limit the scope thereof.

Example 1

Isolation of Protoporphyrin IX Binding Subunit Protein Gene of Magnesium Chelatase.

Genomic DNA of photosynthetic bacterium *Rhodobacter sphaeroides* ATCC17023 was prepared using

ISOPLANT kit for genomic DNA preparation (manufactured by Nippon Gene). Then, according to the description of Gibson, L.C.D. et al., Proc. Natl. Acad. Sci. USA, 92; p 1941 (1995), PCR was carried out by using about 1 µg of said genomic DNA as a template, and 10 pmol of an oligonucleotide composed of nucleotide sequence represented by SEQ ID NO: 1 and 10 pmol of an oligonucleotide composed of nucleotide sequence represented by SEQ ID NO: 2 as primers to amplify the DNA fragment containing protoporphyrin IX binding subunit protein gene *bchH* of magnesium chelatase. The oligonucleotides were prepared with a DNA synthesizer (PE Applied Biosystems: Model 394 DNA/RNA Synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems: OPC Cartridge). The PCR was carried out by maintaining at 94°C for 2 minutes, at 96°C for 40 seconds and then at 68°C for 7 minutes, repeating a cycle for maintaining at 96°C for 40 seconds and then at 68°C for 7 minutes 28 times, and finally maintaining at 96°C for 40 seconds, at 68°C for 7 minutes and then at 72°C for 10 minutes.

Example 2

Expression of Protoporphyrin IX Binding Subunit Protein Gene of Magnesium Chelatase in *Escherichia Coli* (hereinafter abbreviated to *E. coli*)

According to the description of Gibson, L.C.D. et

al., Proc. Natl. Acad. Sci. USA, 92; p 1941 (1995), the DNA fragment containing *bchH* gene prepared in Example 1 was digested with the restriction enzymes NdeI and BglII. The resultant DNA fragment was inserted between NdeI restriction site and BamHI restriction site of expression vector pET11a (manufactured by Stratagene) to obtain plasmid pETBCH (Fig. 1). This plasmid pETBCH was introduced into *E. coli* BL21(DE3) strain competent cells (manufactured by Stratagene) according to the manual attached to the competent cells to obtain *E. coli* BL21(DE3)/pETBCH strain. The strain was inoculated into 1.5 ml LB liquid culture medium (1% tryptone, 0.5% yeast extract, 0.5% NaCl) containing 100 µg/ml ampicillin in a tube (14 x 10 mm), and the tube was covered with aluminum foil (hereinafter referred to as dark conditions), cultured with shaking at 37°C under light of fluorescent lamp (about 8000 lux). When the absorbance at 600 nm of the liquid culture medium became about 0.6, isopropyl β-D-thiogalactopyranoside (IPTG) was added to the liquid culture medium so that the final concentration was 0.4 mM, and the culture was continued for about additional 20 hours. At that time, the *Escherichia coli* turned red and fluorescent absorbance (excitation wavelength 405 nm, emission wavelength 630 nm) which showed the accumulation of protoporphyrin IX in *E. coli* was observed. When *E. coli* BL21(DE3)/pETBCH strain

was cultured according to the same manner except that IPTG was not added, *E. coli* did not turned red and the above fluorescent absorbance did not detected. In contrast to this, when *E. coli* BL21(DE3)/pETBCH strain was cultured according to the same manner (with IPTG) except that the tube was not covered with aluminum foil (hereinafter referred to as light conditions), *E. coli* grew and turned red as above.

Example 3

Expression of PPO Gene Derived from Soybeans in hemG Gene Deficient *E. coli*

Soybeans (Glycine max var. Williams82) were seeded and cultivated at 25°C for 20 days and green leaves were collected. The collected green leaves were frozen with liquid nitrogen and the frozen leaves were ground in a mortar with a pestle. From the ground leaves, RNA were extracted by using RNA extracting reagent ISOGEN (manufactured by Nippon Gene) according to the manual attached thereto. The resultant RNA liquid extract was subjected to ethanol precipitation to collect total RNA, then the total RNA was fractionated by using poly (A) RNA fractionating kit BIOMAG mRNA Purification Kit (manufactured by Perceptive Bio System) according to the manual attached thereto to collect poly (A) RNA fraction. Using 1 µg of this poly (A) RNA fraction as a template, cDNA was

synthesized with the cDNA synthetic reagent contained in Marathon cDNA amplification kit (manufactured by Clontech) according to the manual attached thereto. PCR was carried out by using the resultant cDNA as a template, and an oligonucleotide composed of nucleotide sequence of SEQ ID NO: 3 and an oligonucleotide composed of nucleotide sequence of SEQ ID NO: 4 as primers to amplify the DNA fragment containing chloroplast-type protoporphyrinogen IX oxidase gene. The above oligonucleotides were prepared with a DNA synthesizer (PE Applied Biosystems: Model 394 DNA/RNA Synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems: OPC Cartridge). The PCR was carried out by maintaining at 94°C for 1 minutes and then at 65°C for 5 minutes, repeating a cycle for maintaining at 94°C for 15 seconds and then at 65°C for 5 minutes 29 times. After the PCR, the amplified DNA fragment was purified by filtering the reaction mixture with MicroSpin S-400HR (manufactured by Pharmacia Biotech), and the DNA fragment was ligated to plasmid pCR2.1 (manufactured by Invitrogen) cleaved by restriction enzyme SalI to obtain plasmid pSPPO-P. Then, the plasmid was introduced into competent cells of *E. coli* INVαF' strain (manufactured by Invitrogen) and ampicillin resistant strains were selected. Then, the plasmid contained in selected ampicillin resistant strains was sequenced by using

Dye terminator cycle sequencing kit (manufactured by PE Applied Biosystems) and DNA sequencer 373S (manufactured by PE Applied Biosystems). As a result, the nucleotide sequence of SEQ ID NO: 5 was revealed, thereby confirming that plasmid pSPPO-P contained chloroplast-type protoporphyrinogen IX oxidase gene of soybean.

The plasmid pSPPO-P was digested with restriction enzyme PshBI, the resultant DNA fragment was blunted by using T4 DNA polymerase and further digested with SphI to isolate the DNA fragment containing chloroplast-type PPO gene of soybean and lac promoter. Then, the plasmid pACYC184 (manufactured by Nippon Gene) was digested with the restriction enzymes NruI and SphI to remove a fragment of 410 bp and the above DNA fragment was inserted instead to obtain plasmid pACYCSP (Fig. 2). Then, the plasmid pACYCSP was introduced into PPO gene (hemG gene locus) deficient mutant *E. coli* BT3 strain (described in Yamamoto, F. et al., Japanese J. Genet., 63; p 237 (1988) etc.) according to the method described in Hanahan, D.J., Mol. Biol., 166; p 557 (1983). The resultant *E. coli* were cultured in YPT medium (5 g/liter yeast extract, 5 g/liter tryptone, 5 g/liter peptone, 10 g/liter NaCl, pH 7.0) containing 15 µg/ml chloramphenicol and 10 µg/ml kanamycin to select *E. coli* BT3/pACYCSP strain resistant to chloramphenicol and kanamycin whose hemG gene deficiency

was complemented by PPO gene derived from soybean.

Example 4

Test of Protoporphyrin IX Binding Subunit Protein
of Magnesium Chelatase for Capability of Giving Weed
Control Compound-Resistance

E. coli BT3/pACYCSP strain prepared in Example 3
was inoculated into YPT medium containing 10 or 1 ppm of
PPO inhibitory type herbicidal compound represented by the
above Structure 8, 10 µg/ml hemin, 50 µg/ml aminolevulinic
acid, 15 µg/ml chloramphenicol and 10 µg/ml kanamycin,
cultured under dark conditions or light conditions
according to the same manner as in Example 2. As a control,
E. coli BT3/pACYCSP strain was cultured in the same medium
as above without the herbicidal compounds under the same
conditions. Then, 18 hours after initiation of culture,
the absorbance of the liquid culture medium was measured at
600 nm. By taking the absorbance of the medium without the
herbicidal compound as 1, the relative value of the
absorbance of the medium containing the herbicidal compound
was calculated. The results are shown in Table 1.

Table 1

<i>E. coli</i> strain	Culture conditions	Relative absorbance		
		Concentration of test compound		
		10 ppm	1 ppm	0 ppm
BT3/pACYCSP	in the light	0.10	0.25	1.0
BT3/pACYCSP	in the dark	0.73	0.95	1.0

Plasmid pTVBCH (Fig. 3) was constructed by amplification of the DNA fragment containing *bchH* gene derived from photosynthetic bacterium *Rhodobacter* *sphaeroides* using the oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 1 and the oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 2 according to the same manner as in Example 1, digestion of the resultant DNA fragment with the restriction enzymes NcoI and BglII and introducing the digested DNA fragment between NcoI restriction site and BamHI restriction site of plasmid pTV118N (manufactured by Takara Shuzo Co., Ltd.).

Plasmids pTVBCH and pTV118N respectively were introduced into *E. coli* BT3/pACYCSP strain prepared in Example 3 according to the method described in Hanahan, D.J., Mol. Biol., 166; p 557 (1983). The resultant *E. coli* were cultured in YPT medium containing 100 µg/ml ampicillin, 15 µg/ml chloramphenicol and 10 µg/ml kanamycin to obtain *E. coli* BT3/pACYCSP+pTVBCH strain bearing plasmids pACYCSP and pTVBCH, and *E. coli* BT3/pACYCSP+pTV118N strain bearing plasmids pACYCSP and pTV118N.

These strains were inoculated into YPT medium containing 10 or 1 ppm of the PPO inhibitory-type herbicidal compound represented by the above Structure 8, 100 µg/ml ampicillin, 15 µg/ml chloramphenicol, 10 µg/ml

kanamycin, 10 µg/ml hemin and 50 µg/ml aminolevulinic acid, cultured under dark conditions or light conditions according to the same manner as in Example 2. Then, 18 hours after initiation of culture, the absorbance of the liquid culture medium was measured at 600 nm. By taking the absorbance of the medium without the herbicidal compound as 1, the relative value of the absorbance of the medium containing the herbicidal compound was calculated. The results are shown in Table 2.

Table 2

<i>E. coli</i> strain	Culture conditions	Relative absorbance		
		Concentration of test compound		
		10 ppm	1 ppm	0 ppm
BT3/pACYCSP+pTVBCH	in the light	0.80	0.77	1.0
BT3/pACYCSP+pTVBCH	in the dark	0.90	1.06	1.0
BT3/pACYCSP+pTV118N	in the light	0.18	0.31	1.0
BT3/pACYCSP+pTV118N	in the dark	0.68	0.77	1.0

Further, these strains were inoculated into YPT medium containing PPO inhibitory-type herbicidal compounds represented by the above Structures 1, 14, 15, 18-22, 29, 32, 33, 34 and 36, respectively, 100 µg/ml ampicillin, 15 µg/ml chloramphenicol, 10 µg/ml kanamycin, 10 µg/ml hemin and 50 µg/ml aminolevulinic acid, cultured under dark conditions or light conditions similar to the Example 2. Then, 18 hours after initiation of culture, the absorbance

of liquid culture medium was measured at 600 nm. By taking the absorbance of the medium without the herbicidal compound as 1, the relative value of the absorbance of the medium containing the herbicidal compound was calculated.

5 The results are shown in Table 3.

Table 3

Test compound Structure No.	Test concent- ration	Relative absorbance			
		BT3/ pACYCSP+pTVBCH in the light	BT3/ pACYCSP+pTVBCH in the dark	BT3/ pACYCSP+pTV118N in the light	BT3/ pACYCSP+pTV118N in the dark
Structure 1	5.0	0.88	0.88	0.31	0.87
Structure 14	10	0.47	0.93	0.12	0.81
Structure 15	0.5	0.94	0.94	0.38	0.82
Structure 18	2.0	0.68	1.0	0.33	0.91
Structure 19	5.0	0.78	0.89	0.40	0.71
Structure 20	5.0	0.57	0.88	0.11	0.75
Structure 21	10	0.88	0.91	0.25	0.85
Structure 22	10	0.55	0.93	0.29	0.94
Structure 29	0.5	0.64	0.90	0.22	0.77
Structure 32	2.0	0.70	0.94	0.37	0.87
Structure 33	2.0	0.81	0.92	0.41	0.91
Structure 34	1.0	0.41	0.94	0.19	0.86
Structure 36	0.5	0.55	0.95	0.28	0.96

0067719, 106700

Example 5

Introduction of Gene Encoding Protoporphyrin IX Binding Subunit Protein of Magnesium Chelatase into Tobacco

A plasmid was constructed for introducing bchH gene into a plant by *Agrobacterium* infection method. First, binary vector pBI121 (manufactured by Clontech) was digested with restriction enzyme SacI, and KpnI linker (manufactured by Takara Shuzo Co., Ltd.) was inserted to prepare plasmid pBIK wherein SacI recognition site of pBI121 was removed and KpnI recognition site was added. On the other hand, according to the same manner as described in Example 1, PCR was carried out by using the genomic DNA of photosynthetic bacterium *Rhodobacter sphaeroides* as a template, and the oligonucleotide primer composed of the nucleotide sequence of SEQ ID NO: 7 and the oligonucleotide primer composed of the nucleotide sequence of SEQ ID NO: 8 to amplify the DNA fragment containing bchH gene. Then, the above plasmid pBIK was digested with restriction enzymes XbaI and KpnI to remove β -glucuronidase gene, and instead thereof, a DNA fragment which was obtained by digesting the above DNA fragment containing bchH gene with restriction enzymes XbaI and KpnI was inserted to produce plasmid pBIBCH (Fig. 4) in which bchH gene was joined downstream from 35S promoter. Binary vector pBI121 (manufactured by Clontech) was also digested with

restriction enzymes BamHI and SacI to remove β -glucuronidase gene, the resultant DNA fragment was blunted by using T4 DNA polymerase, followed by self-cyclization with T4 DNA ligase to construct plasmid pNO (Fig. 5). The plasmid was used as a vector control of bchH expression plasmid pBIBCH.

The plasmid pBIBCH and pNO were introduced into *Agrobacterium tumefaciens* LBA4404, respectively. *Abrobacterium* strain bearing pBIBCH and that bearing pNO were isolated by culturing the resultant transformants in a medium containing 300 μ g/ml streptomycin, 100 μ g/ml rifampicin and 25 μ g/ml kanamycin and selecting the desired transformants.

Then, according to the method described in Manual for Gene Manipulation of Plant (by Hirofumi UCHIMIYA, Kodan-sha Scientific, 1992), the gene was introduced into tobacco. *Agrobacterium* strain bearing plasmid pBIBCH was cultured at 28°C overnight in LB medium and then leaf pieces of tobacco cultured sterilely were dipped in the liquid culture medium. The leaf pieces were cultured at room temperature for 2 days in Murashige-Skoog medium (MS-medium, described in Murasige T. and Skoog F., *Physiol. Plant.* (1962) 15, p 473) containing 0.8% agar, 0.1 mg/liter naphthalene acetic acid and 1.0 mg/liter benzyl aminopurine. Then, the leaf pieces were washed with sterilized water and

cultured for 7 days on MS medium containing 0.8% agar, 0.1 mg/liter naphthalene acetic acid, 1.0 mg/liter benzyl aminopurine and 500 µg/ml cefotaxime. The leaf pieces were transplanted onto MS medium containing 0.8% agar, 0.1 mg/liter naphthalene acetic acid, 1.0 mg/liter benzyl aminopurine, 500 µg/ml cefotaxime and 100 µg/ml kanamycin (hereinafter referred to as selective MS medium) and cultured on the medium continuously for 4 months with transplanting the tobacco leaf pieces onto fresh selective MS medium every 1 month. During culture, stem-leaf differentiated shoots were appeared from the tobacco leaf pieces, these shoots were transplanted to MS medium containing 0.8% agar, 300 µg/ml cefotaxime and 50 µg/ml kanamycin to induce roots to obtain regenerated plants. The resultant regenerated plant was transplanted and cultured on MS medium 0.8% agar and 50 µg/ml kanamycin to obtain tobacco plant into which bchH gene was introduced. Similarly, tobacco leaf pieces were infected with *Agrobacterium* strain bearing pNO to obtain regenerated plant from the tobacco leaf pieces and tobacco plant (hereinafter referred to as control recombinant tobacco).

Example 6

Test of Tobacco Bearing Introduced Gene Encoding Protoporphyrin IX Binding Subunit Protein of Magnesium Chelatase for Resistance to Herbicidal Compounds

The tobacco leaves into which bchH gene was introduced and control recombinant tobacco leaves obtained in Example 5 were collected and each leaf was divided into the right and left equivalent pieces along the main vein, respectively. To one piece was applied an aqueous solution containing 0.3 ppm PPO inhibitory-type herbicidal compound of Structure 8, while, to the other piece was not applied the compound. These leaf pieces were placed on MS medium containing 0.8% agar and allowed to stand at room temperature for 7 days in light place. Then, each leaf piece was ground with pestle and mortar in 5 ml of 80% aqueous acetone solution to extract chlorophyll. The extract liquid was diluted with 80% aqueous acetone solution and the absorbance was measured at 750 nm, 663nm and 645 nm to calculate total chlorophyll content according to the method described by Macknney G., J. Biol. Chem. (1941) 140, p 315. The results obtained from 4 clones of tobacco into which bchH gene was introduced (BCH1 to 4) and control recombinant tobacco is shown in Table 4. In the table, the resistant level to the herbicidal compound was represented by percentages of the total chlorophyll content of leaf pieces treated with herbicidal compound to that of untreated leaf pieces.

Table 4

Recombinant tobacco	Total chlorophyll content (mg/ g-fresh weight)		Resistant level to test compound(%)
	untreated-leaf	treated-leaf	
control	2.49	0.19	7.63
BCH-1	1.35	1.70	126
BCH-2	2.06	2.14	104
BCH-3	1.93	1.57	81.3
BCH-4	1.51	1.06	70.2

The tobacco clone into which *bchH* gene was introduced and control recombinant tobacco were also treated in the same manner with the solution containing PPO inhibitory-type herbicidal compound represented by the above Structure 3, 7, 10, 11, 13, 17, 23, 24, 25, 27, 28, 30 or 35, and the resistant level to each herbicidal compound was measured. The results are shown in Table 5. In the table, the resistant levels to the herbicidal compound were represented by percentages of the total chlorophyll content of leaf pieces treated with the herbicidal compound to that of untreated leaf pieces.

Table 5

Test compound Structure No.	Test concentration (ppm)	Resistant level to test compound (%)	
		<i>bchH</i> recombinant tobacco	control recombinant tobacco
Structure 3	10	114	9.94
Structure 7	30	89.3	8.62
Structure 10	10	84.0	14.9

Structure 11	0.30	78.1	5.51
Structure 13	30	95.2	14.8
Structure 17	0.30	80.4	14.3
Structure 23	3.0	106	5.58
Structure 24	10	129	5.18
Structure 25	10	104	16.0
Structure 27	10	86.8	16.8
Structure 28	0.30	72.2	8.79
Structure 30	3.0	102	4.24
Structure 35	0.30	83.3	17.4

Example 7

Isolation of Gene Encoding Variant Protein of
Protoporphyrin IX Binding Subunit Protein of Tobacco
Magnesium Chelatase

Total RNAs were prepared from leaf tissues of tobacco (*Nicotiana tabacum* cv. SR1) by using RNeasy Plant Kit (manufactured by QIAGEN) according to the manual attached thereto. The DNA fragment containing the gene encoding protoporphyrin IX binding subunit protein of tobacco magnesium chelatase whose chloroplast transit signal had been deleted (hereinafter referred to as the variant tobacco chelatase subunit) was obtained by using RNA LA PCR Kit (AMV) Ver 1.1 (manufactured by Takara Shuzo Co., Ltd.) according to the manual attached thereto. First, 1st strand cDNA was synthesized by using tobacco total RNAs as templates and Oligo dT-Adaptor Primer contained in the

above kit as the primer with the reverse transcriptase contained in the above kit. Then, PCR was carried out by using the 1st strand cDNA as a template and LA Taq polymerase contained in the above kit to amplify the DNA fragment containing the gene encoding the variant tobacco chelatase subunit protein. In this PCR, oligonucleotide primer composed of the nucleotide sequence of SEQ ID NO: 9 and the oligonucleotide primer composed of the nucleotide sequence of SEQ ID NO: 10 were used. These oligonucleotides were synthesized by using a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA Synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems; OPC cartridge). The PCR was carried out by maintaining at 94°C for 2 minutes and then repeating a cycle for maintaining at 94°C for 30 seconds, at 50°C for 30 seconds and then at 72°C for 7 minutes 30 times. After the PCR, the DNA fragment amplified by the PCR was cloned into plasmid pCR2.1 by using TA Cloning Kit (manufactured by Invitrogen) according to the manual attached thereto. The resultant plasmid was digested with restriction enzyme KpnI and analyzed by agarose gel electrophoresis. The plasmid from which 8.0 kb DNA fragment was detected was named pTCHLH. The plasmid had the structure that the gene encoding the variant tobacco chelatase subunit has been inserted in the direction

expressible under the control of lac promoter. Plasmid pTCHLH was digested with restriction enzyme KpnI followed by self-ligaiton to obtain plasmid pTCHLH1 (Fig. 6) in which DNA fragment composed of about 60 nucleotides had been deleted from plasmid pTCHLH.

Example 8

Test of Variant Tobacco Magnesium Chelatase Subunit Protein for Capability of Giving Resistance to Herbicidal Compounds

The plasmid pTCHLH1 and pCR2.1 prepared in Example 7 were introduced into *E. coli* BT3/pACYCSP strain prepared in Example 3, respectively according to the method described in Hanahan, D.J., Mol. Biol., 166; p 557 (1983). *E. coli* BT3/pACYCSP+pTCHLH1 strain bearing plasmids pACYCSP and pTCHLH1, and *E. coli* BT3/pACYCSP+pCR2.1 strain bearing plasmids pACYCSP and pCR2.1 were obtained by culturing the above strains in YPT medium containing 100 µg/ml ampicillin, 15 µg/ml chloramphenicol and 50 µg/ml kanamycin, respectively.

These *E. coli* strains were inoculated into YPT medium containing 10 or 1 ppm of the PPO inhibitory-type herbicidal compound represented by Structure 8, 100 µg/ml ampicillin, 15 µg/ml chloramphenicol, 50 µg/ml kanamycin, 10 µg/ml hemin and 50 µg/ml aminolevulinic acid, cultured under dark conditions or light conditions according to the

same manner as in Example 2. Then, 18 hours after initiation of culture, the absorbance of the liquid culture medium was measured at 600 nm. By taking the absorbance of the medium without the herbicidal compound as 1, the relative value of the absorbance of the medium containing the herbicidal compound was calculated. The results are shown in Table 6.

Table 6

<i>E. coli</i> strain	Culture conditions	Relative absorbance		
		concentration of test compound		
		10 ppm	1 ppm	0 ppm
BT3/pACYCSP+pTCHLH1	in the light	0.69	0.89	1.0
BT3/pACYCSP+pTCHLH1	in the dark	0.92	0.93	1.0
BT3/pACYCSP+pCR2.1	in the light	0.03	0.08	1.0
BT3/pACYCSP+pCR2.1	in the dark	1.0	1.0	1.0

Example 9

Introduction of Gene Encoding Variant Tobacco Magnesium Chelatase Subunit Protein into Tobacco

A plasmid for introducing the gene encoding a variant tobacco magnesium chelatase subunit protein into tobacco by *Agrobacterium* infection method was constructed. First, the DNA fragment containing the gene encoding the variant tobacco magnesium chelatase subunit protein was prepared by digesting plasmid pTCHLH1 prepared in Example 7 with restriction enzymes KpnI and SalI. On the other hand,

binary vector pBI121 (manufactured by Clonetech) was digested with restriction enzyme SmaI and KpnI linker (manufactured by Takara Shuzo Co., Ltd.) was inserted into this portion to prepare plasmid pBI121K in which SmaI recognition site of pBI121 was removed and KpnI recognition site was added. The plasmid pBI121K was digested with restriction enzyme SacI followed by blunting the DNA by adding nucleotides to the double-stranded DNA gap with DNA Polymerase I. Then, the DNA was dephosphorylated with alkaline phosphatase derived from calf intestine and cyclized by inserting phosphorylated SalI linker (4680P, manufactured by Takara Shuzo Co., Ltd.) to construct plasmid pBI121KS. The binary vector pBI121KS was digested with restriction enzymes KpnI and SalI to remove β -glucuronidase gene and the gene encoding the variant tobacco magnesium chelatase subunit protein was inserted into this portion to prepare plasmid pBITCHLH (Fig. 7).

The plasmid pBITCHLH was introduced into *Agrobacterium tumefaciens* LBA4404. The resultant transformants were cultured in a medium containing 300 μ g/ml streptomycin, 100 μ g/ml rifampicin and 25 μ g/ml kanamycin, followed by selecting the desired transformants to isolate a *Agrobacterium* strain bearing pBITCHLH.

Leaf pieces of tobacco cultured sterilely are infected with the *Agrobacterium* strain and, according to

the same manner as in Example 5, tobacco into which the gene encoding the variant tobacco magnesium chelatase subunit protein is introduced is obtained.

Example 10

5 Confirmation of Resistance to Herbicidal Compounds of Tobacco Bearing Introduced Gene Encoding Variant Tobacco Magnesium Chelatase Subunit Protein

10 The levels of resistance to herbicidal compounds are confirmed quantitatively by testing tobacco introduced with the gene encoding the variant tobacco magnesium chelatase subunit protein prepared in Example 9 according to the same manner as in Example 6.

Example 11

15 Isolation of Gene Encoding Variant Protein of Soybean PPO Having No Capability of Oxidizing Protoporphyrinogen IX and Having Specific Affinity for Protoporphyrinogen IX

20 PCR was carried out by using plasmid pSPPO-P prepared in Example 3 as a template and an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 11 and an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 12 as primers to amplify the DNA fragment encoding soybean PPO whose chloroplast transit signal and FAD binding sequence had been deleted (hereinafter referred to as the variant soybean PPO). The oligonucleotides were

25

prepared with a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems; OPC cartridge). The PCR was carried out by
5 repeating a cycle for maintaining at 94°C for 1 minute, at 55°C for 2 minutes and the 72°C for 3 minutes 30 times. The amplified DNA fragments were digested with restriction enzymes NcoI and SalI, and introduced between NcoI restriction site and SalI restriction site of plasmid
10 pTV118N (manufactured by Takara Shuzo Co., Ltd.) to construct plasmid pTVGMP (Fig. 8).

The plasmid pTVGMP was introduced into *E. coli* PPO gene deficient mutant BT3 strain according to the method described in Hanahan, D.J., Mol. Biol., 166; p 557 (1983).
15 When the resultant *E. coli* were cultured in YPT medium containing 100 µg/ml ampicillin and 10 µg/ml kanamycin, no growth complemented clone was obtained.

Example 12

Test for Effect of Giving Resistance to
20 Herbicidal Compounds of Variant Soybean PPO

Plasmids pTVGMP and pTV118N prepared in Example 11 were introduced into *E. coli* BT3/pACYCSP strain prepared in Example 3 respectively according to the method described in Hanahan, D.J., Mol. Biol., 166; p 557 (1983). *E. coli*
25 BT3/pACYCSP+pTVGMP strain bearing plasmids pACYCSP and

pTVGMP, and *E. coli* BT3/pACYCSP+ pTV118N strain bearing plasmids pACYCSP and pTV118N were obtained by culturing the above strains in YPT medium containing 100 µg/ml ampicillin, 15 µg/ml chloramphenicol and 10 µg/ml kanamycin.

5 These *E. coli* strains were inoculated into YPT medium containing 10 or 1 ppm of PPO inhibitory-type herbicidal compound represented by Structure 8, 100 µg/ml ampicillin, 15 µg/ml chloramphenicol, 10 µg/ml kanamycin, 10 µg/ml hemin and 50 µg/ml aminolevulinic acid, cultured
10 under dark conditions or light conditions according to the same manner as in Example 2. Then, 18 hours after initiation of culture, the absorbance of liquid culture medium was measured at 600 nm. By taking the absorbance of the medium without the herbicidal compound as 1, the
15 relative value of the absorbance of the medium containing the herbicidal compound was calculated. The results are shown in Table 7.

Table 7

<i>E. coli</i> strain	Culture conditions	Relative absorbance		
		Concentration of test compound		
		10 ppm	1 ppm	0 ppm
BT3/pACYCSP+pTVGMP	in the light	0.33	0.85	1.0
BT3/pACYCSP+pTVGMP	in the dark	0.91	0.94	1.0
BT3/pACYCSP+pTV118N	in the light	0.05	0.09	1.0
BT3/pACYCSP+pTV118N	in the dark	0.89	0.91	1.0

Example 13

Introduction of the Gene Encoding Variant Soybean
PPO into Tobacco

A plasmid for introducing the gene encoding the
5 variant soybean PPO into a plant by *Agrobacterium* infection
method was constructed. PCR was carried out by using the
plasmid pSPPO-P prepared in Example 3 as a template, an
oligonucleotide primer composed of the nucleotide sequence
of SEQ ID NO: 13 and an oligonucleotide primer composed of
10 the nucleotide sequence of SEQ ID NO: 14 to amplify the DNA
fragment containing the gene encoding the variant soybean
PPO. Then, plasmid pBI121K prepared in Example 9 was
digested with the restriction enzymes KpnI and SacI to
remove β -glucuronidase gene, and the DNA fragment which was
15 obtained by digesting the DNA fragment containing the above
gene encoding the variant soybean PPO with restriction
enzymes KpnI and Sac I was inserted into this portion to
prepare plasmid pBIGMP (Fig. 9) in which the gene was
joined downstream from 35S promoter.

20 The plasmid pBIGMP was introduced into
Agrobacterium tumefaciens LBA4404. The resultant
transformants were cultured in a medium containing 300
 $\mu\text{g/ml}$ streptomycin, 100 $\mu\text{g/ml}$ rifampicin and 25 $\mu\text{g/ml}$
kanamycin, followed by selecting the desired transformants
25 to isolate *Agrobacterium* strain bearing pBIGMP.

Leaf pieces of tobacco cultured sterily were infected with the *Agrobacterium* strain and, according to the same manner as in Example 5, tobacco into which the gene encoding the variant soybean PPO was introduced was obtained.

Example 14

Confirmation of Resistance to Herbicidal Compounds of Tobacco Bearing Introduced Gene Encoding Variant Soybean PPO

The level of resistance to PPO inhibitory type herbicidal compound represented by Structure 8 was confirmed quantitatively by testing tobacco into which the gene encoding the variant soybean PPO prepared in Example 13 was introduced according to the same manner as in Example 6. The results obtained from 4 clones (GMP 1-4) of tobacco introduced with the gene encoding the variant soybean PPO and control recombinant tobacco are shown in Table 8. In the table, the resistant level to herbicidal compound is represented by percentage of the total chlorophyll content of leaf pieces treated with the herbicidal compound to that of untreated leaf pieces.

Table 8

Recombinant tobacco	Total chlorophyll content (mg/ g-fresh weight)		Resistant level to test compound(%)
	untreated-leaf	treated-leaf	
control	3.49	0.35	10.0

GMP-1	1.89	2.55	135
GMP-2	0.89	0.96	108
GMP-3	1.50	1.49	99.3
GMP-4	2.91	2.34	80.4

Example 15

Isolation of PPO Gene of *Chlamydomonas*

Chlamydomonas reinhardtii CC407 strain was obtained from Chlamydomonas Genetics Center (address: DCMB Group, Department of Botany, Box 91000, Duke University, Durham, NC 27708-1000, USA), cultured under 200 $\mu\text{E}/\text{m}^2/\text{s}$ photosynthesis active light for 5 days in TAP liquid culture medium (E. H. Harris, The Chlamydomonas Sourcebook, Academic Press, San Diego, 1989, p 576-577) containing 7 mM NH_4Cl , 0.4 mM $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.34 mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 25 mM potassium phosphate, 0.5 mM Tris (pH 7.5), 1 ml/liter Hatner miner element and 1 ml/liter glacial acetic acid to obtain 200 ml (1.0×10^6 cells/ml) liquid culture medium containing early stationary growth phase cells.

Total RNAs were prepared from these cells by using ISOGEN (manufactured by Nippon Gene) according to the manual attached thereto. Also, poly(A)RNA was fractionated using BioMag mRNA Purification Kit (manufactured by Perceptive Bio System) according to the manual attached thereto. cDNA was synthesized from the resultant poly(A)RNA by using Marathon cDNA Amplification Kit (manufactured by

Clontech) according to the manual attached thereto and the cDNA was used as a template for PCR.

As PCR primers, an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 15 and an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 16 were prepared. The oligonucleotides were prepared with a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems; OPC cartridge).

PCR was carried out by preparing a reaction liquid using Advantage cDNA PCR kit (manufactured by Clontech) according to the manual attached thereto, and then, after maintaining at 94°C for 1 minute and then at 65°C for 5 minutes, repeating a cycle for maintaining at 94°C for 15 seconds and the 65°C for 5 minutes 29 times. After the PCR, the amplified DNA fragments were purified by filtering the reaction liquid with MicroSpin S-400HR (manufactured by Pharmacia Biotech), and the DNA fragment was cloned into plasmid pCR2.1 by using TA Cloning Kit (manufactured by Invitrogen) according to the manual attached thereto to construct plasmid pCPPO.

The nucleotide sequence of DNA fragment contained in the resultant plasmid pCPPO was determined by using Dye terminator cycle sequencing kit (manufactured by PE applied

Biosystems) and DNA sequencer 373S (manufactured by PE applied Biosystems). As a result, the nucleotide sequence of SEQ ID NO: 17 was revealed, thereby confirming that plasmid pCPPO contained the full length PPO cDNA of *Chlamydomonas reinhardtii*.

Example 16

Isolation of Gene Encoding Variant Protein of *Chlamydomonas reinhardtii* PPO Having No Capability of Oxidizing Protoporphyrinogen IX and Specific Affinity for Protoporphyrinogen IX

PCR was carried out by using plasmid pCPPO prepared in Example 15 as a template, and an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 19 and an oligonucleotide composed of the nucleotide SEQ ID NO: 20 as primers to amplify the DNA fragment encoding *Chlamydomonas reinhardtii* PPO whose chloroplast transit signal and FAD binding sequence had been deleted (hereinafter referred to as the variant *Chlamydomonas reinhardtii* PPO). The oligonucleotides were prepared with a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems; OPC cartridge). The PCR was carried out by repeating a cycle for maintaining at 94°C for 1 minute, at 55°C for 2 minutes and then at 72°C for 3 minutes 30 times. The amplified DNA

fragment was digested with restriction enzymes BamHI and SacI, and inserted between BamHI restriction site and SacI restriction site of plasmid pTV119N (manufactured by Takara Shuzo Co., Ltd.) to construct plasmid pTVCRP (Fig. 10).

5 The plasmid pTVCRP was introduced into *E. coli* PPO gene deficient mutant BT3 strain according to the method described in Hanahan, D. J., Mol. Biol., 166; p 557 (1983). When the resultant *E. coli* were cultured in YPT medium containing 100 µg/ml ampicillin and 10 µg/ml kanamycin, no
10 growth complemented clone was obtained.

Example 17

Test of Variant Modified *Chlamydomonas reinhardtii* PPO for Capability of Giving Resistance to Herbicidal Compounds

15 Plasmids pTVCRP and pTV118N prepared in Example 16 were introduced into *E. coli* BT3/pACYCSP strain prepared in Example 3 respectively according to the method described in Hanahan, D. J., Mol. Biol., 166; p 557 (1983). *E. coli* BT3/pACYCSP+pTVCRP strain bearing plasmids pACYCSP and
20 pTVCRP, and *E. coli* BT3/pACYCSP+pTV118N strain bearing plasmids pACYCSP and pTV118N were obtained by culturing the above strains in YPT medium containing 100 µg/ml ampicillin, 15 µg/ml chloramphenicol and 10 µg/ml kanamycin.

25 These *E. coli* strains were inoculated into YPT medium containing 10 or 1 ppm of the PPO inhibitory-type

herbicidal compound represented by Structure 8, 100 µg/ml ampicillin, 15 µg/ml chloramphenicol, 10 µg/ml kanamycin, 10 µg/ml hemin and 50 µg/ml aminolevulinic acid, cultured under dark conditions or light conditions in the same manner as in Example 2. Then, 18 hours after initiation of culture, the absorbance of liquid culture medium was measured at 600 nm. By taking the absorbance of the medium containing no herbicidal compound as 1, the relative value of the absorbance of the medium containing the herbicidal compound was calculated. The results are shown in Table 9.

Table 9

<i>E. coli</i> strain	Culture conditions	Relative absorbance		
		Concentration of test compound		
		10 ppm	1 ppm	0 ppm
BT3/pACYCSP+pTVCRP	in the light	0.23	0.42	1.0
BT3/pACYCSP+pTVCRP	in the dark	0.81	0.82	1.0
BT3/pACYCSP+pTV118N	in the light	0.12	0.24	1.0
BT3/pACYCSP+pTV118N	in the dark	0.80	0.91	1.0

Example 18

Introduction of Gene Encoding Variant *Chlamydomonas reinhardtii* PPO into Tobacco

A plasmid for introducing the gene encoding the variant *Chlamydomonas reinhardtii* PPO into a plant by *Agrobacterium* infection method was constructed. The DNA

fragment containing the gene encoding the variant *Chlamydomonas reinhardtii* PPO was prepared by digesting plasmid pTVCRP prepared in Example 16 with restriction enzymes BamHI and SacI. Binary vector pBI121 (manufactured by Clontech) was digested with restriction enzymes BamHI and SacI to remove β -glucuronidase gene and the above gene encoding the variant *Chlamydomonas reinhardtii* PPO was inserted into this portion to prepare plasmid pBICRP (fig. 11).

The plasmid pBICRP was introduced into *Agrobacterium tumefaciens* LBA4404. The resultant transformants were cultured in a medium containing 300 μ g/ml streptomycin, 100 μ g/ml rifampicin and 25 μ g/ml kanamycin, followed by selecting the desired transformants to isolate *Agrobacterium* strain bearing pBICRP.

Leaf pieces of tobacco cultured sterily were infected with the *Agrobacterium* strain and, according to the same manner as in Example 5, tobacco into which the gene encoding the variant *Chlamydomonas reinhardtii* PPO was introduced was obtained.

Example 19

Confirmation of Resistance to Herbicidal Compounds of Tobacco Bearing Introduced Gene Encoding Variant *Chlamydomonas reinhardtii* PPO

The level of resistance to the PPO-inhibitory

type herbicidal compound represented by Structure 8 was confirmed quantitatively by testing tobacco into which the gene encoding the variant *Chlamydomonas reinhardtii* PPO prepared in Example 18 was introduced according to the same manner as in Example 6. The results obtained from 4 clones (CRP 1-4) of tobacco into which the gene encoding the variant *Chlamydomonas reinhardtii* PPO was introduced and control recombinant tobacco is shown in Table 10. In the table, the resistant levels to the herbicidal compound are represented by percentages of the total chlorophyll content of leaf pieces treated with the herbicidal compound to that of untreated leaf pieces.

Table 10

Recombinant tobacco	Total chlorophyll content (mg/ g-fresh weight)		Resistant level to test compound(%)
	untreated-leaf	treated-leaf	
control	2.28	0.42	18.4
CRP-1	1.27	1.54	121
CRP-2	1.50	1.67	111
CRP-3	1.10	1.11	101
CRP-4	1.58	1.57	99.4

Example 20

Test of Variant Protein of Barley Ferrochelatase Having Affinity for Protoporphyrin IX Specifically for Capability of Giving Resistance to Herbicidal Compounds

A plasmid bearing barley ferrochelatase gene was prepared by the method described in Miyamoto, K. et al., Plant Physiol. 105; p 769 (1994). The resultant plasmid was digested with restriction enzymes NspI and EcoRI to obtain the DNA fragment containing the gene encoding barley ferrochelatase whose signal sequence had been deleted (hereinafter referred to as the variant barley ferrochelatase). This DNA fragment was inserted between SphI restriction site and EcoRI restriction site of plasmid pTV119N (manufactured by Takara Shuzo Co., Ltd.) to construct plasmid pTVHVF1 (Fig. 12).

The plasmids pTVHVF1 and pTV118N were introduced into *E. coli* BT3/pACYCSP strains prepared in Example 3 respectively according to the method described in Hanahan, D.J., Mol. Biol., 166; p 557 (1983). *E. coli* BT3/pACYCSP+pTVHVF1 strain bearing plasmid pACYCSP and pTVHVF1, and *E. coli* BT3/pACYCSP+pTV118N strain bearing plasmid pACYCSP and pTV118N were obtained by culturing the above strains in YPT medium containing 100 µg/ml ampicillin, 15 µg/ml chloramphenicol and 10 µg/ml kanamycin.

These *E. coli* strains were inoculated into YPT medium containing 10 or 1 ppm of the PPO inhibitory-type herbicidal compound represented by Structure 8, 100 µg/ml ampicillin, 15 µg/ml chloramphenicol, 10 µg/ml kanamycin, 10 µg/ml hemin and 50 µg/ml aminolevulinic acid, cultured

under dark conditions or light conditions according to the same manner as in Example 2. Then, 18 hours after initiation of culture, the absorbance of liquid culture medium was measured at 600 nm. By taking the absorbance of the medium without the herbicidal compound as 1, the relative value of the absorbance of the medium containing the herbicidal compound was calculated. The results are shown in Table 11.

Table 11

<i>E. coli</i> strain	Culture conditions	Relative absorbance		
		Concentration of test compound		
		10 ppm	1 ppm	0 ppm
BT3/pACYCSP+pTVHVF1	in the light	0.39	0.94	1.0
BT3/pACYCSP+pTVHVF1	in the dark	0.94	0.96	1.0
BT3/pACYCSP+pTV118N	in the light	0.12	0.24	1.0
BT3/pACYCSP+pTV118N	in the dark	0.80	0.91	1.0

Example 21

Introduction of the Gene Encoding Variant Barley Ferrochelatase into Tobacco

A plasmid for introducing the gene encoding barley ferrochelatase into tobacco by *Agrobacterium* infection method was constructed. The plasmid pTVHVF1 described in Example 20 was digested with restriction enzyme Nco I followed by blunting the DNA with DNA polymerase I by

adding nucleotides to the double-stranded DNA gap. Then, the DNA was dephosphorylated with alkaline phosphatase derived from calf intestine and cyclized by inserting phosphorylated BamHI linker (4610P, manufactured by Takara Shuzo Co., Ltd.) to construct plasmid pTVHVF2. Then, pTVHVF2 was digested with restriction enzyme EcoRI followed by blunting of the DNA with DNA polymerase I by adding nucleotides to the double-stranded DNA gap. Further, the DNA was dephosphorylated with alkaline phosphatase derived from calf intestine and cyclized by inserting phosphorylated SalI linker (4680P, manufactured by Takara Shuzo Co., Ltd.) to construct plasmid pTVHVF3. Plasmid pBI121KS prepared in Example 9 was digested with restriction enzymes BamHI and SalI to remove β -glucuronidase gene. The DNA fragment containing the gene encoding the variant barley ferrochelatase was prepared by digesting the above pTVHVF3 with restriction enzymes BamHI and SalI. The resultant DNA fragment was inserted into plasmid pBI121KS with replacing β -glucuronidase gene to prepare plasmid pBIHVF (Fig. 13) in which variant barley gene joined downstream from 35S promoter.

The plasmid pBIHVF was introduced into *Agrobacterium tumefaciens* LBA4404. The resultant transformants were cultured in a medium containing 300 μ g/ml streptomycin, 100 μ g/ml rifampicin and 25 μ g/ml

kanamycin, followed by selecting the desired transformants to isolate *Agrobacterium* strain bearing pBIHVF.

Leaf pieces of tobacco cultured sterilely were infected with said *Agrobacterium* strain and, according to the same manner as in Example 5, tobacco into which the gene encoding the variant barley ferrochelatase was introduced was obtained.

Example 22

Confirmation of Resistance to Herbicidal Compounds of Tobacco Bearing Introduced Gene Encoding Variant Barley Ferrochelatase

The level of resistance to the PPO inhibitory-type herbicidal compound represented by Structure 8 was confirmed quantitatively by testing tobacco into which the gene encoding the variant barley ferrochelatase prepared in Example 21 was introduced according to the same manner as in Example 6. The results obtained from 4 clones (HVF 1-4) of tobacco introduced with the gene encoding the variant barley ferrochelatase and control recombinant tobacco are shown in table 12. In the table, the resistant levels to the herbicidal compound are represented by percentages of the total chlorophyll content of leaf pieces treated with herbicidal compound to that of untreated leaf pieces.

Table 12

Recombinant tobacco	Total chlorophyll content (mg/ g-fresh weight)		Resistant level to test compound(%)
	untreated-leaf	treated-leaf	
control	1.93	0.160	8.29
HVF-1	0.876	0.930	106
HVF-2	1.14	1.16	102
HVF-3	1.06	1.04	98.1
HVF-4	1.48	1.42	95.9

Example 23

Test of Variant Protein of Cucumber
 5 Ferrochelatase Having Specific Affinity for Protoporphyrin
 IX for Capability of Giving Resistance to Herbicidal
 Compounds

PCR was carried out by using cucumber
 ferrochelatase cDNA clone isolated by the method described
 10 in Miyamoto, K. et al., Plant Physiol., 105; p 769 (1994)
 as a template, an oligonucleotide composed of the
 nucleotide sequence of SEQ ID NO: 21 and an oligonucleotide
 composed of the nucleotide sequence of SEQ ID NO: 22 as
 primers to amplify the DNA fragment encoding cucumber
 15 ferrochelatase whose signal sequence had been deleted
 (hereinafter referred to as the variant cucumber
 ferrochelatase). The oligonucleotides were prepared with a
 DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA
 synthesizer) and purified with an oligonucleotides

purification cartridge (PE Applied Biosystems; OPC cartridge). The PCR was carried out by repeating a cycle for maintaining at 94°C for 1 minute, at 55°C for 2 minutes and then at 72°C for 3 minutes 30 times. The amplified DNA fragments were digested with restriction enzymes BamHI and SacI, and inserted between BamHI restriction site and SacI restriction site of plasmid pTV119N (manufactured by Takara Shuzo Co., Ltd.) to construct plasmid pTVCSF (Fig. 14).

The plasmids pTVCSF and pTV118N were introduced into *E. coli* BT3/pACYCSP strain prepared in Example 3 respectively according to the method described in Hanahan, D.J., Mol. Biol., 166; p 557 (1983). *E. coli* BT3/pACYCSP+ pTVCSF strain bearing plasmid pACYCSP and pTVCSF, and *E. coli* BT3/pACYCSP+pTV118N strain bearing plasmid pACYCSP and pTV118N were obtained by culturing the above strains in YPT medium containing 100 µg/ml ampicillin, 15 µg/ml chloramphenicol and 10 µg/ml kanamycin.

These *E. coli* strains were inoculated into YPT medium containing 10 or 1 ppm of the PPO inhibitory-type herbicidal compound represented by Structure 8, 100 µg/ml ampicillin, 15 µg/ml chloramphenicol, 10 µg/ml kanamycin, 10 µg/ml hemin and 50 µg/ml aminolevulinic acid, cultured under dark conditions or light conditions according to the same manner as in Example 2. Then, 18 hours after initiation of culture, the absorbance of liquid culture

medium was measured at 600 nm. By taking the absorbance of the medium without the herbicidal compound as 1, the relative value of the absorbance of the medium containing the herbicidal compound was calculated. The results are shown in Table 13.

Table 13

E. coli strain	Culture conditions	Relative absorbance		
		Concentration of test compound		
		10 ppm	1 ppm	0 ppm
BT3/pACYCSP+pTVCSF	in the light	0.73	0.78	1.0
BT3/pACYCSP+pTVCSF	in the dark	0.89	0.92	1.0
BT3/pACYCSP+pTV118N	in the light	0.06	0.08	1.0
BT3/pACYCSP+pTV118N	in the dark	0.81	0.91	1.0

Example 24

Introduction of the Gene Encoding Variant Cucumber Ferrochelataase into Tobacco

A plasmid for introducing the gene encoding the modified cucumber ferrochelataase into tobacco by *Agrobacterium* infection method was constructed. Plasmid pBI121 (manufactured by Colntech) was digested with restriction enzymes BamHI and SacI to remove β -glucuronidase gene. A DNA fragment containing the gene encoding the variant cucumber ferrochelataase was prepared by digesting plasmid pTVCSF described in Example 23 with restriction

enzymes BamHI and SacI. The resultant DNA fragment was introduced into plasmid pBI121 with replacing β -glucuronidase gene to prepare plasmid pBICSF (Fig. 15) in which variant cucumber ferrochelatase gene was joined downstream from 35S promoter.

The plasmid pBICSF was introduced into *Agrobacterium tumefaciens* LBA4404. The resultant transformants were cultured in a medium containing 300 μ g/ml streptomycin, 100 μ g/ml rifampicin and 25 μ g/ml kanamycin, followed by selecting the desired transformants to isolate *Agrobacterium* strain bearing pBICSF.

Leaf pieces of tobacco cultured sterilely were infected with said *Agrobacterium* strain to obtain tobacco introduced with the gene encoding the modified cucumber ferrochelatase according to the same manner as in Example 5.

Example 25

Confirmation of Resistance to Herbicidal Compounds of Tobacco Bearing Introduced Gene Encoding Variant Cucumber Ferrochelatase

The level of resistance to PPO inhibitory-type herbicidal compounds is confirmed quantitatively by testing tobacco introduced with the gene encoding the modified cucumber ferrochelatase prepared in Example 24 according to the same manner as in Example 6.

Example 26

Isolation of *E. coli* Coproporphyrinogen III
Oxidase (*hemF*) Gene

Genomic DNA was prepared from *E. coli* LE392 strain using Kit ISOPLANT for genome DNA preparation (manufactured by Nippon Gene). An oligonucleotide primer composed of the nucleotide sequence of SEQ ID NO: 23 and an oligonucleotide primer composed of the nucleotide sequence of SEQ ID NO: 24 were synthesized according to nucleotide sequences of its 5' and 3' regions of *E. coli* *hemF* gene registered in GenBank (Accession X75413). The oligonucleotides were prepared with a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA synthesizer) and purified with an oligonucleotides purification cartridge (PE Applied Biosystems; OPC cartridge). PCR was carried out by using about 1 µg of *E. coli* LE392 strain genomic DNA as a template and the above oligonucleotides (each 10 pmol) as primers to amplify the DNA fragment containing *E. coli* *hemF* gene. The PCR was carried out by repeating a cycle for maintaining at 96°C for 1 minute, at 55°C for 2 minutes and then at 72°C for 3 minutes 30 times.

Example 27

Test of *E. coli* *hemF* Protein for Capability of Giving Resistance to Herbicidal Compounds

The DNA fragment containing *hemF* gene amplified by the method described in Example 26 was digested with

restriction enzymes FbaI and PstI, and inserted between BamHI restriction site and PstI restriction site of commercially available plasmid pUC118N (manufactured by Takara Shuzo Co., Ltd.) to construct plasmid pHEMF (Fig. 16).

5 The plasmid pHEMF and pTV118N were introduced into *E. coli* BT3/pACYCSP strain prepared in Example 3 respectively according to the method described in Hanahan, D. J., Mol. Biol., 166; p 557 (1983). *E. coli* BT3/pACYCSP+ pHEMF strain bearing plasmid pACYCSP and pHEMF, and *E. coli* 10 BT3/pACYCSP+pTV118N strain bearing plasmid pACYCSP and pTV118N were obtained by culturing the above strains in YPT medium containing 100 µg/ml ampicillin, 15 µg/ml chloramphenicol and 10 µg/ml kanamycin.

15 These *E. coli* strains were inoculated into YPT medium containing 10 or 1 ppm of the PPO inhibitory-type herbicidal compound represented by Structure 8, 100 µg/ml ampicillin, 15 µg/ml chloramphenicol, 10 µg/ml kanamycin, 10 µg/ml hemin and 50 µg/ml aminolevulinic acid, cultured under dark conditions or light conditions according to the 20 same manner as in Example 2. Then, 18 hours after initiation of culture, the absorbance of liquid culture medium was measured at 600 nm. By taking the absorbance of the medium without the herbicidal compound as 1, the relative value of the absorbance of the medium containing 25 the herbicidal compound was calculated. The results are

shown in Table 14.

Table 14

<i>E. coli</i> strain	Culture conditions	Relative absorbance		
		Concentration of test compound		
		10 ppm	1 ppm	0 ppm
BT3/pACYCSP+pHEMF	in the light	0.48	1.0	1.0
BT3/pACYCSP+pHEMF	in the dark	0.94	0.95	1.0
BT3/pACYCSP+pTV118N	in the light	0.06	0.16	1.0
BT3/pACYCSP+pTV118N	in the dark	0.96	0.98	1.0

Example 28

Introduction of *E. coli hemF* gene into Tobacco

A plasmid for introducing *E. coli hemF* gene into a plant by *Agrobacterium* infection method was constructed. First, for obtaining *E. coli hemF* gene, an oligonucleotide primer composed of the nucleotide sequence of SEQ ID NO: 25 and an oligonucleotide primer composed of the nucleotide sequence of SEQ ID NO: 26 were synthesized with a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems; OPC cartridge). PCR was carried out by using the oligonucleotide primers according to the same manner as in Example 26 to amplify the DNA fragment containing *E. coli hemF* gene.

Plasmid pBI121 (manufactured by Clontech) was digested with restriction enzymes BamHI and SacI to remove β -glucuronidase gene. The DNA fragment containing the gene encoding the *E. coli* hemF gene was prepared by digesting the above PCR-amplified DNA fragment with restriction enzymes BamHI and SacI. The resultant DNA fragment was introduced into plasmid pBI121 with replacing β -glucuronidase gene to prepare plasmid pBIHEMF (Fig. 17) in which *E. coli* hemF gene was joined downstream from 35S promoter.

The plasmid pBIHEMF was introduced into *Agrobacterium tumefaciens* LBA4404. The resultant transformants were cultured in a medium containing 300 μ g/ml streptomycin, 100 μ g/ml rifampicin and 25 μ g/ml kanamycin, followed by selecting the desired transformants to isolate *Agrobacterium* strain bearing pBIHEMF.

Leaf pieces of tobacco cultured sterily were infected with the *Agrobacterium* strain to obtain tobacco introduced with *E. coli* hemF gene according to the same manner as in Example 5.

Example 29

Confirmation of Resistance to Herbicidal Compounds of Tobacco Introduced with the *E. coli* hemF Gene
The level of resistance to the PPO inhibitory-type herbicidal compounds is confirmed quantitatively by testing

tobacco introduced with the *E. coli* hemF gene (prepared in Example 28) according to the same manner as in Example 6.

Example 30

Binding Test of Porphyrin Compound-Binding
5 Protein to Protoporphyrin IX

A phage library presenting a protein containing an amino acid sequence composed of 5 random amino acids and a phage clone displaying a protein containing an amino acid sequence HASYS or RASSL (wherein H is histidine, A is alanine, S is serine, Y is tyrosine, R is arginine and L is leucine) which can specifically bind to porphyrin compound 5, 10, 15, 20-tetrakis (N-methylpyridinium-4-yl)-21H, 23H-porphine (H_2TMPyP) were prepared according to the method described in KITANO et al., Nihon Kagakukai (Chemical Society of Japan) 74th Spring Annual Meeting Pre-Published Abstracts of Presentation II, p 1353, 4G511 (1998).

First, the phage library displaying a protein containing an amino acid sequence composed of 5 random amino acids was constructed. Mixed oligonucleotides composed of the nucleotide sequence of SEQ ID NO: 27 and mixed oligonucleotides composed of the nucleotide sequence of SEQ ID NO: 28 were synthesized. The mixed oligonucleotides were synthesized with a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA synthesizer) and purified with an oligonucleotide purification cartridge (PE

Applied Biosystems; OPC cartridge). The above mixed oligonucleotides (each 50 pmol) were phosphorylated at 5' end by treating with T4 DNA kinase respectively. They were mixed and, after heating at 70°C for 10 minutes, subjected to annealing by cooling slowly to room temperature at rate of 0.5°C/minute. Plasmid pCANTAB5E (manufactured by Pharmacia Biotech) was digested with restriction enzymes SfiI and NotI to remove the recombinant antibody gene ScFv. The above phosphorylated and annealed oligonucleotide pair was inserted into the portion of the above recombinant antibody gene ScFv to prepare a plasmid containing a nucleotide sequence encoding a protein composed of a 5 random amino acid sequence upstream from a protein comprising an amino acid sequence of M13 phage coat protein. The plasmid was introduced into *E. coli* TG-1 strain according to the method described in Hanahan, D.J., Mol. Biol. 166; p 557 (1983) and cultured in 2 x YT medium (10 g/liter yeast extract, 15 g/liter tryptone and 5 g/liter NaCl, pH 7.2) containing 100 µg/ml ampicillin to obtain recombinant *E. coli* TG-1 strain. The recombinant *E. coli* TG-1 strain was inoculated into 2 x YT medium containing 100 µg/ml ampicillin and cultured with shaking at 37°C. Then, 1 hour after initiation of culture, 6×10^{10} pfu helper-phage M13K07 (manufactured by Pharmacia Biotech) was inoculated to the medium, and culture was continued for additional 18

hours with shaking. Then, the liquid culture medium was centrifuged at 1,000 x g for 20 minutes to collect the phage library displaying a protein containing the amino acid sequence composed of 5 random amino acids.

5 For preparing the phage clone displaying a protein containing the amino acid sequence HASYS (SEQ ID NO: 53), an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 29 and an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 30 were synthesized. And, for preparing the phage clone displaying a protein containing the amino acid sequence RASSL (SEQ ID NO: 55), an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 31 and an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 32 were synthesized. These oligonucleotides were synthesized with a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems; OPC cartridge). The phage clone displaying the protein containing the amino acid sequence HASYS or RASSL was obtained by the same operation as the above that for obtaining the phage library displaying a protein containing the amino acid sequence composed of 5 random amino acids.

25 A phage suspension containing the phage clone displaying the protein containing the amino acid sequence

HASYS, the phage clone displaying the protein containing the amino acid sequence RASSL or the phage library displaying the protein containing the amino acid sequence consisting of 5 random amino acids (titer 10^5 pfu) was respectively spotted to nitro cellulose filter (manufactured by Schleicher & Schuell), and then the nitro cellulose filter was blocked by shaking it in PBT buffer (137 mM NaCl, 8.10 mM Na_2HPO_4 , 2.68 mM KCl, 1.47 mM KH_2PO_4 , 0.05% Tween 20, pH 7.2) containing 1% bovine serum albumin. The nitro cellulose filter was washed with PBT buffer and shaken for 18 hours in 2 x SSC buffer (0.3 M NaCl, 0.03M sodium citric acid) containing 10 μM protoporphyrin IX. Further, said nitro cellulose filter was washed with 2 x SSC buffer, dried, and fluorescence derived from protoporphyrin IX was detected under ultraviolet light (365 nm).

The spots of the phage library did not show fluorescence, while the spots of both phage clones displaying the protein containing the amino acid sequence HASYS and that containing the amino acid sequence RASSL showed clear fluorescence.

Example 31

Test of Protoporphyrin IX Binding Protein for Capability of Giving Resistance to Herbicidal Compounds

First, a plasmid which could express the gene

encoding the protein containing the amino acid sequence HASYS (SEQ ID NO: 53), or the amino acid sequence RASSL (SEQ ID NO: 55) was prepared. For preparing the plasmid capable of expressing the gene encoding the protein composed of the amino acid sequence of SEQ ID NO: 54 (hereinafter referred to as the protein MGHASYS), an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 33 and an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 34 were synthesized. The oligonucleotides were synthesized with a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems; OPC cartridge). The above oligonucleotides (each 50 pmol) were phosphorylated at 5' end by treating with T4 DNA kinase, respectively. They were mixed and then, after heating for 10 minutes at 70°C, subjected to annealing by cooling slowly to room temperature at rate of 0.5°C/minute. Plasmid pTV118N was digested with restriction enzymes NcoI and EcoRI to remove the gene fragment consisting of 16 base pairs. Plasmid pHASYS capable of expressing the gene encoding protein MGHASYS was prepared by inserted the above phosphorylated and annealed oligonucleotide pairs into the position of the above 16 base pairs.

Then, for preparing the plasmid capable of

expressing the gene encoding the protein consisting of amino acid sequence of SEQ ID NO: 56 (hereinafter referred to as protein MGRASSL), an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 35 and an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 36 were synthesized. The oligonucleotides were synthesized with a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems; OPC cartridge). Plasmid pRASSL capable of expressing the gene encoding protein MGRASSL was prepared by the same procedure as that for plasmid PHASYS.

A plasmid capable of expressing the gene encoding the protein containing the amino acid sequence YAGY or YAGF (wherein Y is tyrosine, A is alanine, G is glycine, F is phenylalanine) (Sugimoto, N., Nakano, S., Chem. Lett. p 939, 1997) capable of binding to porphyrin compound H_2TMPyP was prepared. For preparing the plasmid capable of expressing the gene encoding the protein consisting of the amino acid sequence of SEQ ID NO: 58 (hereinafter referred to as protein MG_YAGY), an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 37 and an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 38 were synthesized. For preparing the plasmid capable of expressing the gene encoding the protein composed of the

amino acid sequence of SEQ ID NO: 60 (hereinafter referred to as protein MGYAGF), an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 39 and an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 40 were also synthesized. These oligonucleotides were synthesized with a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems; OPC cartridge). Plasmid pYAGY capable of expressing the gene encoding the protein MGYAGY and plasmid pYAGF capable of expressing the gene encoding protein MGYAGF were prepared by the same procedure as that for plasmid PHASYS.

The above plasmids PHASYS, PRASSL, pYAGY, pYAGF and pTV118N were introduced into *E. coli* BT3/pACYCSP strain prepared in Example 3 respectively according to the method described in Hanahan, D. J., Mol. Biol., 166; p 557 (1983). *E. coli* BT3/pACYCSP+PHASYS strain bearing plasmid pACYCSP and PHASYS, *E. coli* BT3/pACYCSP+PRASSL strain bearing plasmid pACYCSP and PRASSL, *E. coli* BT/pACYCSP+pYAGY strain bearing plasmid pACYCSP and pYAGY, *E. coli* BT3/pACYCSP+pYAGF strain bearing plasmid pACYCSP and pYAGF and *E. coli* BT3/pACYCSP+pTV118N strain bearing plasmid pACYCSP and pTV118N were obtained by culturing the above strains in YPT medium containing 100 µg/ml ampicillin, 15 µg/ml chloramphenicol and 10 µg/ml kanamycin.

These *E. coli* strains were inoculated into YPT medium containing 1 ppm of the PPO inhibitory-type herbicidal compound represented by Structure 8, 100 µg/ml ampicillin, 15 µg/ml chloramphenicol, 10 µg/ml kanamycin, 10 µg/ml hemin and 50 µg/ml aminolevulinic acid, cultured under dark conditions or light conditions according to the same manner as in Example 2. Then, 18 hours after initiation of culture, the absorbance of liquid culture medium was measured at 600 nm. By taking the absorbance of the medium without the herbicidal compound as 1, the relative value of the absorbance of the medium containing the herbicidal compound was calculated. The results are shown in Table 15.

Table 15

<i>E. coli</i> strain	Culture conditions	Relative absorbance	
		Concentration of test compound	
		1 ppm	0 ppm
BT3/pACYCSP+pHASYS	in the light	0.65	1.0
BT3/pACYCSP+pHASYS	in the dark	0.96	1.0
BT3/pACYCSP+pRASSL	in the light	0.59	1.0
BT3/pACYCSP+pRASSL	in the dark	1.0	1.0
BT3/pACYCSP+pYAGY	in the light	0.48	1.0
BT3/pACYCSP+pYAGY	in the dark	0.99	1.0
BT3/pACYCSP+pYAGF	in the light	0.62	1.0
BT3/pACYCSP+pYAGF	in the dark	0.96	1.0

BT3/pACYCSP+pTV118N	in the light	0.07	1.0
BT3/pACYCSP+pTV118N	in the dark	0.93	1.0

Further, a plasmid capable of expressing a gene encoding a protein containing an amino acid sequence in which one unit of the amino acid sequences HASYS or RASSL were repeatedly joined. For preparing the plasmid capable of expressing the gene encoding the protein composed of the amino acid sequence of SEQ ID NO: 61 (hereinafter referred to as protein MG(HASYS)₄, (HASYS)_n referred to as a sequence in which peptide HASYS was repeatedly joined to each other n times), an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43 or SEQ ID NO: 44 was synthesized. These oligonucleotides were synthesized with a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems; OPC cartridge). First, the oligonucleotide composed of the nucleotide sequence of SEQ ID NO. 42 and the oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 43 were phosphorylated respectively at 5' end by treating with T4 DNA kinase. Thereafter, the oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 41 and the oligonucleotide composed of the phosphorylated nucleotide sequence of SEQ ID NO: 42

or the oligonucleotide composed of the phosphorylated nucleotide sequence of SEQ ID NO: 43 and the oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 44 were mixed (each 300 pmol), and after heating for 5 minutes at 70°C, annealed by cooling slowly to room temperature at rate of 0.5°C/minute. The above two annealed oligonucleotide pairs were mixed and ligated with T4 DNA ligase, then the resultant DNA fragment was phosphorylated with T4 DNA kinase at 5' end. On the other hand, vector pTV118N was digested with restriction enzymes NcoI and EcoRI to remove a DNA fragment of 16 base pairs and the above phosphorylated DNA fragment was inserted into this portion to obtain plasmid pHASYS4 expressing the gene encoding protein MG(HASYS)₄.

Further, for preparing the plasmid capable of expressing the gene encoding the protein composed of the amino acid sequence of SEQ ID NO: 62 (hereinafter referred to as protein MG(HASYS)₈), an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 45 and an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 46 were synthesized. These oligonucleotides were synthesized with a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems; OPC cartridge). First, the above

oligonucleotides were phosphorylated at 5' end by treating with T4 DNA kinase. Thereafter, an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 41 and an oligonucleotide composed of the phosphorylated nucleotide sequence of SEQ ID NO: 42 were mixed (each 300 pmol), an oligonucleotide composed of the phosphorylated nucleotide sequence of SEQ ID NO: 43 and an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 44 were mixed (each 300 pmol), and further, an oligonucleotide composed of the phosphorylated nucleotide sequence of SEQ ID NO: 45 and an oligonucleotide composed of the phosphorylated nucleotide sequence of SEQ ID NO: 46 were mixed (each 600 pmol). These three mixtures were heated for 5 minutes at 70°C, and annealed by cooling slowly to room temperature at rate of 0.5°C/minute, respectively. The above three annealed oligonucleotide pairs were mixed, and ligated with T4 DNA ligase, and then the resultant DNA fragment was phosphorylated with T4 DNA kinase at 5' end. Plasmid pHASYS8 capable of expressing protein MG(HASYS)₈ were prepared in the same manner as that for the above plasmid pHASYS4.

Then, for preparing a plasmid capable of expressing the gene encoding the protein composed of the amino acid sequence of SEQ ID NO: 63 (hereinafter referred to as protein MG(RASSL)₄, (RASSL)_n referred to as a

sequence in which peptide RASSL was repeatedly joined to each other n times), an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49 or SEQ ID NO: 50 were synthesized. Also, for preparing a plasmid capable of expressing the gene encoding the protein composed of the amino acid sequence of SEQ ID NO: 64 (hereinafter referred to as protein MG(RASSL)₈), an oligonucleotide composed of the nucleotide sequence of SEQ ID NO: 51 and an oligonucleotide composed of the nucleotide sequence of SEQ ID No: 52 were synthesized. These oligonucleotides were synthesized with a DNA synthesizer (PE Applied Biosystems; Model 394 DNA/RNA synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems; OPC cartridge).

Plasmid pRASSL4 capable of expressing protein MG(RASSL)₄ were prepared according to the same manner as that for the above plasmid pHASYS4. Plasmid pRASSL8 capable of expressing protein MG(RASSL)₈ were also prepared according to the same manner as that for the above plasmid pHASYS8.

The above plasmids pHASYS4, pHASYS8, pRASSL4, pRASSL8 and pTV118N were introduced into *E. coli* BT3/pACYCSP strain prepared in Example 3 respectively according to the method described in Hanahan, D. J., Mol. Biol., 166; p 557 (1983). *E. coli* BT3/pACYCSP+pHASYS4

strain bearing plasmid pACYCSP and pHASYS4, *E. coli* BT3/pACYCSP+pHASYS8 strain bearing plasmid pACYCSP and pHASYS8, *E. coli* BT3/pACYCSP+pRASSL4 strain bearing plasmid pACYCSP and pRASSL4, *E. coli* BT3/pACYCSP+pRASSL8 strain bearing plasmid pACYCSP and pRASSL8 and *E. coli* BT3/pACYCSP+pTV118N strain bearing plasmid pACYCSP and pTV118N were obtained by culturing the above strains in YPT medium containing 100 µg/ml ampicillin, 15 µg/ml chloramphenicol and 10 µg/ml kanamycin.

These *E. coli* strains were inoculated into YPT medium containing 1 ppm of the PPO inhibitory-type herbicidal compound represented by Structure 8, 100 µg/ml ampicillin, 15 µg/ml chloramphenicol, 10 µg/ml kanamycin, 10 µg/ml hemin and 50 µg/ml aminolevulinic acid, cultured under dark conditions or light conditions according to the same manner as in Example 2. Then, 18 hours after initiation of culture, the absorbance of the liquid culture medium was measured at 600 nm. By taking the absorbance of the culture medium without the herbicidal compound as 1, the relative value of the absorbance of the culture medium containing the herbicidal compound was calculated. The results are shown in Table 16.

Table 16

E. coli strain	Culture condition	Relative absorbance	
		Concentration of test compound	
		1 ppm	0 ppm
BT3/pACYCSP+pHASYS4	in the light	0.91	1.0
BT3/pACYCSP+pHASYS4	in the dark	1.0	1.0
BT3/pACYCSP+pHASYS8	in the light	0.57	1.0
BT3/pACYCSP+pHASYS8	in the dark	1.0	1.0
BT3/pACYCSP+pRASSL4	in the light	1.1	1.0
BT3/pACYCSP+pRASSL4	in the dark	0.98	1.0
BT3/pACYCSP+pRASSL8	in the light	0.79	1.0
BT3/pACYCSP+pRASSL8	in the dark	1.0	1.0
BT3/pACYCSP+pTV118N	in the light	0.15	1.0
BT3/pACYCSP+pTV118N	in the dark	0.81	1.0

Example 32

Introduction of the Gene Encoding Protoporphyrin IX Binding Peptide into Tobacco

A plasmid for introducing the gene encoding the protoporphyrin IX binding peptide into tobacco by Agrobacterium method was constructed. The plasmid pHASYS8 prepared in Example 31 was digested with restriction enzyme NcoI followed by blunting the DNA with DNA polymerase I with addition of nucleotides to the double-stranded DNA gap. Then, the DNA was dephosphorylated with alkaline phosphatase derived from calf intestine and cyclized by inserting phosphorylated BamH I linker (4610P, manufactured by Takara Syuzo Co., Ltd.) to construct plasmid pHASYS8B. Plasmid

pBI121 (manufactured by Clontech) was digested with restriction enzymes BamHI and SacI to remove β -glucuronidase gene. On the other hand, plasmid pHASYS8B was digested with restriction enzymes BamHI and SacI to prepare the DNA
5 fragment containing the gene encoding protein MG(HASYS), the resultant DNA fragment was inserted into plasmid pBI121 with replacing β -glucuronidase gene to prepare plasmid pBIHASYS8 (Fig. 18) in which the gene encoding protoporphyrin IX binding protein MG(HASYS), was joined
10 downstream from 35S promoter.

A plasmid for introducing the gene encoding the protoporphyrin IX binding peptide MG(RASSL), into a plant by *Agrobacterium* infection method was constructed. Plasmid pBIRASSL8 (Fig. 19) in which the gene encoding
15 protoporphyrin IX binding protein MG(RASSL), was joined downstream from 35S promoter was prepared from pRASSL8 according to the same procedure as that for pBIHASYS8.

The above plasmid pBIHASYS8 and pBIRASSL8 were introduced into *Agrobacterium tumefaciens* LBA4404
20 respectively. The resultant transformants were cultured in a medium containing 300 μ g/ml streptomycin, 100 μ g/ml rifampicin and 25 μ g/ml kanamycin, followed by selecting the desired transformants to isolate *Agrobacterium* strains bearing pBIHASYS8 and pBIRASSL8, respectively.

25 Leaf pieces of tobacco cultured sterilely are

infected with said *Agrobacterium* strains to obtain tobacco introduced with the gene encoding protoporphyrin IX binding protein MG(HASYS)₈, and the tobacco introduced with the gene encoding protoporphyrin IX binding protein MG(RASSL)₈, in the same manner as in Example 5.

Example 33

Confirmation of Resistance to Herbicidal Compounds of Tobacco Bearing Introduced Gene Encoding the Protoporphyrin IX Binding Peptide

The level of resistance to herbicidal compounds is confirmed quantitatively by testing tobacco introduced with the gene encoding the protoporphyrin IX binding peptide prepared in Example 32 according to the same manner as in Example 14.

Example 34

Isolation of PPO Gene of *Arabidopsis thaliana* Having Herbicidal Compound-Resistant Mutation

A plasmid containing PPO gene of *Arabidopsis thaliana* obtained by the method described by Narita, S. et al., Gene, 182; p 169 (1996) was digested with the restriction enzyme NcoI, and nucleotides were added to the gap of the double-stranded DNA by using DNA Polymerase I to blunt the end of the DNA. Then, the 5'-end of the DNA was dephosphorylated with an alkaline phosphatase derived from calf small intestine, followed by insertion of a

phosphorylated BamHI linker (4810P manufactured by Takara Shuzo Co., Ltd.) therein and cyclization to construct plasmid pAGE17B. The plasmid pAGE17B was digested with BamHI and SacI to obtain a gene fragment containing PPO gene of *Arabidopsis thaliana*. The fragment was inserted between BamHI and SacI of a commercially available vector, pKF19k for site-directed mutagenesis (manufactured by Takara Shuzo Co., Ltd.), to construct plasmid pKFATP.

Then, for conversion of the 220th alanine into valine, a PPO inhibitory-type herbicidal compound-resistant mutation in PPO protein of *Arabidopsis thaliana* disclosed in WO 9534659, base substitution (substitution of "T" for the 659 base "C") of DNA was introduced into the above PPO gene of *Arabidopsis thaliana*. First, an oligonucleotide primer for mutagenesis represented by SEQ ID NO: 65 was synthesized. The oligonucleotide primer was synthesized with a DNA synthesizer (PE Applied Biosystems: Model 394 DNA/RNA Synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems: OPC Cartridge). The 5'-end of the oligonucleotide primer was phosphorylated with T4 DNA kinase. According to the manual attached to a commercially available site-directed mutagenesis kit, Mutan-Super Express Km (manufactured by Takara Shuzo Co., Ltd.), a reaction mixture containing 10 ng of the above plasmid pKFATP as template DNA, 5 pmol of

the attached selection primer, 5 pmol of the above phosphorylated oligonucleotide primer for mutagenesis and the like was prepared and PCR was carried out. The PCR was carried by repeating a cycle for maintaining at 94°C for 1 minute, at 55°C for 1 minute and then at 72°C for 3 minutes 30 times. The resultant reaction mixture was purified by ethanol precipitation and the precipitate was dissolved in 5 µl of sterilized-distilled water. According to the attached manual, its 2 µl portion was used for introduction into a commercially available *E. coli* competent cell, MV1184 (manufactured by Takara Shuzo Co., Ltd.), and plated on LB agar culture medium (1% tryptone, 0.5% yeast extract, 0.5% NaCl, 1% agar) containing 50 µg/ml of kanamycin. After incubation at 37°C, the resultant clone was cultured in LB liquid medium containing 50 µg/ml of kanamycin to prepare plasmid DNA. The introduction of the desired herbicidal compound-resistant mutation A220V was confirmed by analyzing the nucleotide sequence of the PPO gene of *Arabidopsis thaliana* having herbicidal compound-resistant mutation contained in the resultant plasmid pKFATP1.

Example 35

Introduction of PPO Gene of *Arabidopsis thaliana* Having Herbicidal Compound-Resistant Mutation into Tobacco

A plasmid was constructed for introducing the PPO gene of *Arabidopsis thaliana* having herbicidal compound-

resistant mutation (hereinafter referred to as *Arabidopsis thaliana* PPO(A220V) gene) into a plant by *Agrobacterium* infection method. Binary vector pBI121 (manufactured by Clontech) was digested with the restriction enzymes BamHI and SacI to remove β -glucuronidase gene. On the other hand, the plasmid pKFATP1 described in Example 34 was digested with restriction enzymes BamHI and SacI to prepare a DNA fragment containing *Arabidopsis thaliana* PPO(A220V) gene. Instead of the above β -glucuronidase gene, the resultant DNA fragment was inserted in the binary vector pBI121 to construct the plasmid pNATP (Fig. 20).

The plasmid pNATP was introduced into *Agrobacterium tumefaciens* LBA4404 and this was cultured in LB culture medium containing 300 μ g/ml of streptomycin, 100 μ g/ml of rifampicin and 25 μ g/ml of kanamycin, followed by selection of a transformant to isolate an *Agrobacterium* strain bearing pNATP.

Tobacco leaf pieces cultured sterilely were infected with the *Agrobacterium* strain and, according to the same manner as described in Example 5, tobacco bearing the introduced *Arabidopsis thaliana* PPO(A220V) gene was obtained.

Example 36

Production of Recombinant Tobacco Having *Arabidopsis thaliana* PPO(A220V) Gene and Gene Encoding

Variant Tobacco Chelatase Subunit

A plasmid was constructed for introducing both *Arabidopsis thaliana* PPO(A220V) gene and gene encoding a variant tobacco chelatase subunit into a plant by *Agrobacterium* infection method. First, oligonucleotide primers composed of the nucleotide sequence represented by SEQ ID NO: 66 and the nucleotide sequence represented by SEQ ID NO: 67, respectively, were synthesized. The oligonucleotide primers were synthesized by a DNA synthesizer (PE Applied Biosystems: Model 394 DNA/RNA Synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems: OPC Cartridge). PCR was carried out by using the primers and the plasmid pKFATP1 constructed in Example 34 as template DNA to prepare a DNA fragment containing *Arabidopsis thaliana* PPO(A220V) gene. The PCR was carried out by repeating a cycle for maintaining at 94°C for 1 minute, 55°C for 2 minutes and then 72°C for 3 minutes 30 times. The amplified DNA fragment was digested by the restriction enzymes HindIII and SalI to obtain a DNA fragment containing *Arabidopsis thaliana* PPO(A220V) gene. The fragment was inserted between HindIII and SalI restriction sites of a commercially available vector, pUC19, to construct plasmid pAP.

On the other hand, the plasmid pNG01 (Fig. 29)

described in Shiota, N. et al., Plant Physiol., 106; p 17 (1994) was digested with the restriction enzyme HindIII and nucleotides were added to the gap of the double-stranded DNA with DNA Polymerase I to blunt the end of the DNA, followed by self-cyclization to construct the plasmid pNG04 (Fig. 30). The plasmid pNG04 was digested with the restriction enzyme XbaI to isolate a DNA fragment of about 1.1 kb composed of the terminator of a nopaline synthase and 35S promoter. The DNA fragment was inserted into the XbaI restriction site of the above plasmid pAP. For selecting plasmid pAPNS wherein the terminator of the nopaline synthase was ligated to the downstream of *Arabidopsis thaliana* PPO(A220V) gene, digestion with the restriction enzymes HindIII and PstI was carried out to select a clone producing a DNA fragment of about 2.0 kb composed of *Arabidopsis thaliana* PPO(A220V) gene and the terminator of the nopaline synthase. Further, the plasmid pAPNS was digested with the restriction enzyme HindIII and nucleotides were added to the gap of the double-stranded DNA with DNA polymerase I to blunt the end of the DNA. The 5'-end of the DNA was dephosphorylated with an alkaline phosphatase derived from calf small intestine and a phosphorylated KpnI linker (4668P manufactured by Takara Shuzo Co., Ltd.) was inserted therein, followed by cyclization to construct plasmid pAPNSK.

The plasmid pAPNSK was digested with the restriction enzymes KpnI and DraI to isolate a DNA fragment of about 2.8 kb composed of *Arabidopsis thaliana* PPO(A220V) gene, the terminator of the nopaline synthase located at the downstream of the gene and 35S promoter located at the downstream of the terminator. The fragment was inserted in KpnI restriction site of the plasmid pBITCHLH constructed in Example 9. For selecting the plasmid pBIAPTCH (Fig. 21) wherein *Arabidopsis thaliana* PPO(A220V) gene and a gene encoding a variant tobacco chelatase subunit were ligated to the downstream of 35S promoter, respectively, the resultant plasmid was digested with the restriction enzyme BamHI to select a clone producing a DNA fragment of about 2.8 kb composed of *Arabidopsis thaliana* PPO(A220V) gene, the terminator of the nopaline synthase and the 35S promoter.

The plasmid pBIAPTCH was introduced into *Agrobacterium tumefaciens* LBA4404 and this was cultured in LB medium containing 300 µg/ml of streptomycin, 100 µg/ml rifampicin and 25 µg/ml kanamycin, followed by selection of a transformant to isolate an *Agrobacterium* strain bearing pBIAPTCH.

Tobacco leaf pieces cultured sterilely were infected with the *Agrobacterium* strain and, according to the same manner as in Example 5, tobacco bearing the

introduced both *Arabidopsis thaliana* PPO(A220V) gene and gene encoding the variant tobacco chelatase subunit was obtained.

Example 37

Confirmation of Resistance to Herbicidal Compound of Tobacco Bearing Introduced *Arabidopsis thaliana* PPO(A220V) Gene and Gene Encoding Variant Tobacco Chelatase Subunit

Leaves of the tobacco bearing the introduced both *Arabidopsis thaliana* PPO(A220V) gene and gene encoding the variant tobacco chelatase subunit produced in Example 36, those bearing the introduced *Arabidopsis thaliana* PPO(A220V) gene produced in Example 35, and the control recombinant tobacco leaves produced in Example 5 were collected, and each leaf was divided into the right and left equivalent pieces along the main vein. One of the pieces was treated with an aqueous solution containing 2.0 ppm PPO inhibitory-type herbicidal compound of Structure 8, while the other piece was not treated with the compound. These leaf pieces were placed on MS medium containing 0.8% agar and allowed to stand at room temperature for 7 days in a light place. Then, each leaf piece was ground in 5 ml of 80% aqueous acetone solution in a mortar with a pestle to extract chlorophyll. The extract was diluted 10 times with 80% aqueous acetone solution and the absorbance was

measured at 750 nm, 663 nm and 645 nm to calculate the total chlorophyll content according to the method described by Macknney G., J. Biol. Chem. (1941) 140, p 315. The resistant level to the herbicidal compound tested was represented by the percentage of the total chlorophyll content of the leave pieces treated with the herbicidal compound to that of untreated leaf pieces. The resistant level of the control recombinant tobacco was 2.88% and that of the tobacco bearing the introduced *Arabidopsis thaliana* PPO(A220V) gene was 12.2%. On the other hand, the resistant level of the tobacco bearing introduced both *Arabidopsis thaliana* PPO(A220V) gene and gene encoding the variant tobacco chelatase subunit was 61.6%.

Example 38

Isolation of Gene Encoding Chloroplast-Localized Type Ferrochelatase of *Arabidopsis thaliana*

Total RNAs were prepared from leaf tissues of *Arabidopsis thaliana* ecotype WS by using RNeasy Plant Kit (manufactured by QIAGEN) according to the manual attached thereto. A DNA fragment containing a chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana* was obtained by using RNA LA PCR Kit (AMV) Ver 1.1 (manufactured by Takara Shuzo Co., Ltd.) according to the manual attached thereto. First, 1st strand cDNA was synthesized by using *Arabidopsis thaliana* total RNAs as a

template and Oligo dT-Adaptor Primer contained in the above kit as the primer with the reverse transcriptase contained in the above kit. Then, PCR was carried out by using the 1st strand cDNA as a template and LA Taq polymerase
5 contained in the above kit to amplify a DNA fragment containing the gene encoding the chloroplast-localized type ferrochelatase of *Arabidopsis thaliana*. In this PCR, the primers used were the oligonucleotide primer composed of the nucleotide sequence of SEQ ID NO: 68 and the
10 oligonucleotide primer composed of the nucleotide sequence of SEQ ID NO: 69. The oligonucleotides were synthesized by a DNA synthesizer (PE Applied Biosystems: Model 394 DNA/RNA Synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems: OPC
15 Cartridge). The PCR was carried by maintaining at 94°C for 2 minutes and then repeating a cycle for maintaining at 94°C for 30 seconds, at 50°C for 30 seconds and then at 72°C for 7 minutes 30 times. After the PCR, the DNA fragment amplified by the PCR was cloned into the plasmid
20 pCR2.1 by using TA Cloning Kit (manufactured by Invitrogen) according to the manual attached thereto. The resultant plasmid was digested with the restriction enzyme BamHI and analyzed by agarose gel electrophoresis. The plasmid from which 5.3 kb DNA fragment was detected was named pCRATF
25 (Fig. 22). The plasmid has such a structure that the

chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana* is inserted in a DNA strand complementary to the lac promoter. When the nucleotide sequence of the chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana* was analyzed, it agreed with the nucleotide sequence of the chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana* described by Smith, A.G. et al., J. Biol. chem., 269; p 13405 (1994).

Example 39

Introduction of Gene Encoding Chloroplast-localized Type Ferrochelatase of *Arabidopsis thaliana*

A plasmid was constructed for introducing a gene encoding chloroplast-localized type ferrochelatase of *Arabidopsis thaliana* into a plant by *Agrobacterium* infection method. First, the plasmid pCRATF constructed in Example 38 was digested with the restriction enzymes BamHI and SacI to prepare a DNA fragment containing chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana*. Binary vector pBI121 (manufactured by Clontech) was digested with the restriction enzymes BamHI and SacI to remove β -glucuronidase gene and, instead of this gene, the above DNA fragment containing the chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana* was inserted therein to construct the plasmid pBIATF (Fig. 23) wherein the ferrochelatase gene was ligated to the

downstream of the 35S promoter.

The plasmid pBIATF was inserted into *Agrobacterium tumefaciens* LBA4404 and this was cultured in LB medium containing 300 µg/ml of streptomycin, 100 µg/ml of rifampicin and 25 µg/ml of kanamycin, followed by selection of a transformant to isolate an *Agrobacterium* strain bearing pBIATF.

Tobacco leaf pieces cultivated sterilely are infected with the *Agrobacterium* strain and tobacco bearing the introduced chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana* is obtained according to the same manner as in Example 5.

Example 40

Production of Recombinant Tobacco Bearing Both *Arabidopsis thaliana* PPO(A220V) Gene and Chloroplast-Localized Type Ferrochelatase Gene of *Arabidopsis thaliana*

A plasmid is constructed for introducing both *Arabidopsis thaliana* PPO(A220V) gene and chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana* into a plant by *Agrobacterium* infection method. First, the plasmid pAPNS constructed in Example 36 are digested with the restriction enzyme HindIII and nucleotides are added to the gap of the double-stranded DNA with DNA Polymerase I to blunt the end of the DNA. The 5'-end of the DNA is dephosphorylated with an alkaline phosphatase derived from

calf small intestine and a phosphorylated BamHI linker (4610P manufactured by Takara Shuzo Co., Ltd.) is inserted therein, followed by cyclization to construct plasmid pAPNSB.

5 The plasmid pAPNSB is digested with the restriction enzymes BamHI and DraI to isolate a DNA fragment of about 2.8 kb composed of *Arabidopsis thaliana* PPO(A220V) gene, the terminator of nopaline synthase and 35S promoter. The fragment is inserted in the BamHI
10 restriction site of the plasmid pBIATF constructed in Example 39. For selecting the plasmid pBIAPATF (Fig. 24) wherein *Arabidopsis thaliana* PP(A220V) gene and chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana* are ligated to the downstream of 35S
15 promoter, the resultant plasmid is digested with the restriction enzymes NotI and SacI to select a clone producing a DNA fragment of about 2.2 kb composed of the 35S promoter and the chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana*.

20 The plasmid pBIAPATF is introduced into *Agrobacterium tumefaciens* LBA44044 and this is cultured in LB medium containing 300 µg/ml of streptomycin, 100 µg/ml of rifampicin and 25 µg/ml of kanamycin, followed by selection of a transformant to isolate an *Agrobacterium*
25 strain bearing pBIAPATF.

Tobacco leaf pieces cultivated sterilely are infected with the *Agrobacterium* strain and tobacco bearing the introduced *Arabidopsis thaliana* PPO(A220V) gene and chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana* is obtained according to the same manner as in Example 5.

Example 41

Confirmation of Resistance to Herbicidal Compounds of Tobacco Bearing Introduced *Arabidopsis thaliana* PPO(A220V) Gene and Chloroplast-Localized Type Ferrochelatase Gene of *Arabidopsis thaliana*

The levels of resistance to herbicidal compounds are confirmed quantitatively by testing the tobacco bearing the introduced *Arabidopsis thaliana* PPO(A220V) gene and chloroplast-localized type ferrochelatase gene of *Arabidopsis thaliana* produced in Example 40 according to the same manner as in Example 37.

Example 42

Isolation of Soybean Coproporphyrinogen III Oxidase Gene

Total RNAs were prepared from leaf tissues of soybean (*Glycine max* cv. Jack) by using RNeasy Plant Kit (manufactured by QIAGEN) according to the manual attached thereto. Further, a DNA fragment containing a gene encoding soybean coproporphyrinogen III oxidase

(hereinafter referred to as the present soybean CPOX) was obtained by using RNA LA PCR Kit (AMV) Ver 1.1 (manufactured by Takara Shuzo Co., Ltd.) according to the manual attached thereto. First, 1st strand cDNA was synthesized by using the soybean total RNAs as a template and Oligo dT-Adaptor Primer contained in the above kit as a primer with the reverse transcriptase contained in the above kit. Then, PCR was carried out by using the 1st strand cDNA as a template and LA Taq polymerase contained in the above kit to amplify a DNA fragment containing the present soybean CPOX gene. In this PCR, an oligonucleotide primer composed of the nucleotide sequence represented by SEQ ID NO: 70 and an oligonucleotide primer composed of the nucleotide sequence represented by SEQ ID NO: 71 were used. These oligonucleotides were synthesized by using a DNA synthesizer (PE Applied Biosystems: Model 394 DNA/RNA Synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Biosystems: OPC Cartridge). The PCR was carried out by maintaining at 94°C for 2 minutes and then repeating a cycle for maintaining at 94°C for 30 seconds, at 50°C for 30 seconds and then at 72°C for 7 minutes 30 times. After the PCR, the DNA fragment amplified by the PCR was cloned into plasmid pCR2.1 by using TA Cloning Kit (manufactured by Invitrogen) according to the manual attached thereto. The resultant

plasmid was digested with the restriction enzyme BamHI and analyzed by agarose gel electrophoresis. The plasmid from which 1.2 kb DNA fragment was detected was named pCRSCPOX (Fig. 25). The plasmid pCRSCPOX has such a structure that the present soybean CPOX gene is inserted into a DNA strand complementary to the lac promoter. When the nucleotide sequence of the DNA fragment in the plasmid was analyzed, it was confirmed to be the present soybean CPOX gene.

Example 43

Introduction of Present Soybean CPOX Gene into Tobacco

A plasmid was constructed for introducing the present soybean CPOX gene into a plant by *Agrobacterium* infection method. First, the plasmid pCRSCPOX constructed in Example 42 was digested with the restriction enzymes BamHI and SalI to prepare a DNA fragment containing the present soybean CPOX gene. The plasmid pBI121KS constructed in Example 9 was digested with the restriction enzymes BamHI and SalI to remove β -glucuronidase gene and, instead thereof, the above DNA fragment containing the present soybean CPOX gene was inserted therein to construct the plasmid pBISCPOX (Fig. 26) wherein the gene was ligated to the downstream of 35S promoter.

The plasmid pBISCPOX was introduced into *Agrobacterium tumefaciens* LBA4404 and it was cultured in LB

medium containing 300 µg/ml of streptomycin, 100 µg/ml of rifampicin and 25 µg/ml of kanamycin, followed by selection of a transformant to isolate an *Agrobacterium* strain bearing pBISCPOX.

5 Tobacco leaf pieces cultured sterilely is infected with the *Agrobacterium* strain and, according to the same manner as in Example 5, tobacco bearing the introduced present soybean CPOX gene is obtained.

Example 44

10 Production of Recombinant Tobacco Bearing *Arabidopsis thaliana* PPO(A220V) Gene and Present Soybean CPOX Gene

A plasmid is constructed for introducing both *Arabidopsis thaliana* PPO(A220V) gene and present soybean CPOX gene into a plant by *Agrobacterium* infection method. The plasmid pAPNSB constructed in Example 40 is digested with the restriction enzymes BamHI and DraI to isolate a DNA fragment of about 2.8 kb composed of *Arabidopsis thaliana* PPO(A220V) gene, the terminator of a nopaline synthase and 35S promoter. The fragment is inserted into the BamHI restriction site of the plasmid pBISCPOX constructed in Example 43. For selecting the plasmid pBIAPSCP (Fig. 27) wherein *Arabidopsis thaliana* PPO(A220V) gene and the present soybean CPOX gene are ligated to the downstream of the 35S promoter, respectively, the resultant

25

plasmid is digested with the restriction enzymes NotI and SalI to select a clone producing a DNA fragment of about 2.0 kb composed of the 35S promoter and the present CPOX gene.

5 The plasmid pBIAPSCP is introduced into *Agrobacterium tumefaciens* LBA4404 and this is cultured in LB medium containing 300 µg/ml of streptomycin, 100 µg/ml of rifampicin and 25 µg/ml of kanamycin, followed by selection of a transformant to isolate an *Agrobacterium* strain bearing pBIAPSCP.

10 Tobacco leaf pieces cultivated sterily is infected with the *Agrobacterium* strain and, according to the same manner as in Example 5, tobacco bearing the introduced both *Arabidopsis thaliana* PPO(A220V) gene and present soybean CPOX gene is obtained.

Example 45

Confirmation of Resistance to Herbicidal Compounds of Tobacco Bearing Introduced Both *Arabidopsis thaliana* PPO(A220V) Gene and Present Soybean CPOX Gene

20 The levels of resistance to herbicidal compounds are confirmed quantitatively by testing the tobacco bearing the introduced both *Arabidopsis thaliana* PPO(A220V) gene and present soybean CPOX gene produced in Example 44 according to the same manner as in Example 37.

25 Example 46

Isolation of Glyphosate Resistant Gene

Glyphosate resistant soybean (*Glycine max*) was seeded and cultivated at 27°C for 30 days. The first leaves of germinated individuals were collected, were frozen in liquid Nitrogen and were grounded in a mortar with a pestle. Genomic DNA was extracted from the ground material with a genomic DNA extracting reagent ISOPLANT (manufactured by NIPPON GENE) according to the manual attached thereto. PCR was carried out by using the genomic DNA as a template, an oligonucleotide primer composed of the nucleotide sequence represented by SEQ ID NO: 72 and an oligonucleotide primer composed of the nucleotide sequence represented by SEQ ID NO: 73 to amplify a DNA fragment (hereinafter referred to as the present CTP-CP4 EPSPS gene) containing a nucleotide sequence encoding a chloroplast transit peptide sequence of EPSPS of petunia (*Petunia hybrida*) (hereinafter referred to as CTP) and EPSPS gene of *Agrobacterium* (*Agrobacterium* sp. Strain CP4). The oligonucleotides were synthesized by using a DNA synthesizer (PE Applied Biosystems: Model 394 DNA/RNA Synthesizer) and purified with an oligonucleotide purification cartridge (PE Applied Systems: OPS Cartridge). The PCR was carried out by maintaining at 94°C for 5 minutes, 55°C for 2 minutes and then at 72°C for 3 minutes, and further repeating a cycle for maintaining at 94°C for 1

minute, at 55°C for 2 minutes and then 72°C for 3 minutes
38 times, and, finally, further maintaining at 94°C for 1
minute, at 55°C for 2 minutes and then 72°C for 10 minutes.
The amplified DNA fraction was ligated to a PCR product
5 cloning site of plasmid pCR2.1 (manufactured Invitrogen),
to construct the plasmid pCREPSPS (Fig. 28). Then, the
plasmid was introduced in a competent cell of *E. coli* JM109
strain (manufactured by Takara Shuzo Co., Ltd.) to select
an ampicillin resistant strain. The nucleotide sequence of
10 the plasmid contained in the selected ampicillin resistant
strain was determined by using Thermo Sequence II Dye
Terminator kit (manufacture by Amersham Pharmacia Biotech)
and DNA Sequencer 373S (manufactured by PE Applied
Biosystems). As a result, the nucleotide sequence
15 represented by SEQ ID NO: 74 was revealed and it was
confirmed that the plasmid pCREPSPS contained the present
CTP-CP4 EPSPS gene.

Example 47

Introduction of Present CTP-CP4 EPSPS Gene into

Tobacco

A plasmid was constructed for introducing the
present CTP-CP4 EPSPS gene into a plant by *Agrobacterium*
infection method. First, pNG01 [Shiota et al., (1994)
Plant Physiol., 106:17-23] (Fig. 29) was digested with the
25 restriction enzyme HindIII and nucleotides were added to

the gap of the double-stranded DNA with DNA Polymerase I to blunt the end of the DNA, followed by self-cyclization with T4 DNA ligase to obtain pNG04 (Fig. 30). The plasmid pNG04 was digested with the restriction enzyme XbaI to obtain a DNA fragment containing the terminator of a nopaline synthase and 35S promoter located at the downstream thereof. The fragment was inserted in the XbaI restriction site of plasmid pUC19 (manufactured by Takara Shuzo Co., Ltd.) to obtain pNT35S (Fig. 31). Then, the plasmid pCREPPSPS constructed in Example 46 was digested with the restriction enzymes HindIII and SalI and the resultant DNA fragment containing the present CTP-CP4 EPSPS gene was inserted between HindIII and SalI restriction sites of pNT35S to obtain the plasmid pCENS (Fig. 32). The plasmid pCENS was digested with the restriction enzyme HindIII and nucleotides were added to the gap of the double-stranded DNA with DNA Polymerase I to blunt the end of the DNA. The 5'-end of the DNA was dephosphorylated with treatment of alkaline phosphatase derived from calf small intestine, followed by insertion of phosphorylated KpnI linker (4668A manufactured by Takara Shuzo Co., Ltd.) therein and cyclization to obtain the plasmid pCENSK (Fig. 33). The plasmid pBI121KS constructed in Example 9 was digested with the restriction enzymes KpnI and SalI to remove β -glucuronidase gene and, instead thereof, a DNA fragment

containing the present CTP-CP4 EPSPS gene, which was obtained by digesting the above plasmid pCENSK with the restriction enzymes KpnI and SalI, was inserted therein to construct the plasmid pBICE (Fig. 34) wherein the present CTP-CP4 EPSPS gene was ligated to the downstream of 35S promoter.

The plasmid pBICE was introduced into *Agrobacterium tumefaciens* LBA44044 (manufactured by Clontech) and this was cultured in LB medium (0.5% yeast extract, 1.0% Bacto tryptone, 0.5% NaCl) containing 300 µg/ml of streptomycin, 100 µg/ml of rifampicin and 25 µg/ml of kanamycin, followed by selection of a transformant to isolate an *Agrobacterium* strain bearing pBICE.

Tobacco leaf pieces cultivated sterilely were infected with the *Agrobacterium* strain and, according to the same manner as in Example 5, tobacco bearing the inserted present CTP-CP4 EPSPS gene was obtained.

Example 48

Introduction of Present CTP-CP4 EPSPS Gene and Gene Encoding Variant Tobacco Chelatase Subunit into Tobacco

A plasmid was constructed for introducing the present CTP-CP4 EPSPS gene and a gene encoding a variant tobacco chelatase subunit into a plant by *Agrobacterium* infection method. First, the plasmid pCENSK constructed in

Example 47 was digested with the restriction enzyme KpnI to obtain a DNA fragment containing the present CTP-CP4 EPSPS gene, a terminator of a gene encoding nopaline synthase located at the downstream thereof and 35S promoter located at the downstream of the terminator. This was inserted into the KpnI restriction site of the plasmid pBITCHLH constructed in Example 9 to construct the plasmid pBICETCH (Fig. 35) wherein the present CTP-CP4 EPSPS gene and the gene encoding the variant tobacco chelatase subunit were ligated to the downstream of 35S promoter, respectively.

The plasmid pBICETCH was introduced into *Agrobacterium tumefaciens* LBA44044 and this was cultured in LB medium containing 300 µg/ml of streptomycin, 100 µg/ml of rifampicin and 25 µg/ml of kanamycin, followed by selection of a transformant to isolate an *Agrobacterium* strain bearing pBICETCH.

Tobacco leaf pieces cultivated sterilely were infected with the *Agrobacterium* strain and, according to the same manner as described in Example 5, tobacco bearing the inserted present CTP-CP4 EPSPS gene and gene encoding the variant tobacco chelatase subunit was obtained.

Example 49

Confirmation of Resistance to Herbicidal Compounds of Tobacco Bearing Introduced Present CTP-CP4 EPSPS Gene as well as Tobacco Bearing Introduced Present

CTP-CP4 EPSPS Gene and Gene Encoding Variant Tobacco
Chelatase Subunit

Leaves of the tobacco bearing the introduced present CTP-CP4 EPSPS gene produced in Example 47, those of the control recombinant tobacco obtained in Example 5 and those bearing the introduced present CTP-CP4 gene and gene encoding the variant tobacco chelatase subunit produced in Example 48 are collected, and each leaf is divided into the right and left equivalent pieces along the main vein. One of the pieces is treated with an aqueous solution containing 0.3 ppm PPO inhibitory-type herbicidal compound of Structure 8, while to the other piece is not treated with the compound. These leaf pieces are placed on MS medium containing 0.8% agar and allowed to stand at room temperature for 7 days in a light place. Then, each leaf piece is ground in 5 ml of 80% aqueous acetone solution in a mortar with a pestle to extract chlorophyll. The extract is diluted 10 times with 80% aqueous acetone solution and the absorbance is measured at 750 nm, 663 nm and 645 nm to calculate the total chlorophyll content according to the method described by Macknney G., J. Biol. Chem. (1941) 140, p 315. The resistant level to the herbicidal compound tested is represented by the percentage of the total chlorophyll content of the leave piece treated with the herbicidal compound to that of untreated leaf piece.

Similarly, the tobacco bearing the introduced present CTP-CP4 EPSPS gene, the tobacco bearing the introduced both present CTP-CP4 EPSPS gene and gene encoding the variant tobacco chelatase subunit, and the control recombinant tobacco are treated with an aqueous solution containing 100 ppm of a glyphosate to determine the resistant level to the glyphosate. The resistant level to the glyphosate is represented by the percentage of the total chlorophyll content of the leave piece treated with the glyphosate to that of untreated leaf piece.

Example 50

Introduction of Present CTP-CP4 EPSPS Gene and Gene Encoding Variant Soybean PPO into Tobacco

A plasmid was constructed for introducing the present CTP-CP4 EPSPS gene and a gene encoding a variant soybean PPO into a plant by *Agrobacterium* infection method. According to the same manner as described in Example 11, PCR was carried out by using an oligonucleotide primer composed of the nucleotide sequence represented by SEQ ID NO: 75, an oligonucleotide primer composed of the nucleotide sequence represented by SEQ ID NO: 76, and the plasmid pSPPO-P constructed in Example 3 as a template to amplify a DNA fragment containing the gene encoding the variant soybean PPO. Then, the plasmid pBI121K constructed in Example 9 was digested with restriction enzymes KpnI and

SacI to remove β -glucuronidase gene and, instead thereof, a DNA fragment obtained by digesting the above DNA fragment containing the gene encoding the variant soybean PPO with the restriction enzymes KpnI and SacI was inserted therein to construct the plasmid pBIGMP (Fig. 36) wherein the gene was ligated to the downstream of the 35S promoter.

Then, the plasmid pCENSK constructed in Example 47 is digested with the restriction enzyme KpnI to obtain a DNA fragment containing the present CTP-CP4 EPSPS gene, the terminator of the gene encoding nopaline synthase located at the downstream of the gene and the 35S promoter located at the downstream of the terminator, followed by insertion of it in the KpnI restriction site of the above plasmid pBIGMP to construct the plasmid pBICEGMP (Fig. 37) wherein the present CTP-CP4 EPSPS gene and the gene encoding the variant soybean PPO are ligated to the downstream of the 35S promoter, respectively.

The plasmid pBICEGMP is introduced into *Agrobacterium tumefaciens* LBA44044 and this is cultured in LB medium containing 300 μ g/ml of streptomycin, 100 μ g/ml of rifampicin and 25 μ g/ml of kanamycin, followed by selection of a transformant to isolate an *Agrobacterium* strain bearing pBICEGMP.

Tobacco leaf pieces cultivated sterilely are infected with the *Agrobacterium* strain and, according to

the same manner as in Example 5, tobacco bearing the introduced present CTP-CP4 EPSPS gene and gene encoding the variant soybean PPO is obtained.

Example 51

Confirmation of Resistance to Herbicidal Compound of Tobacco Bearing Introduced Present CTP-CP4 EPSPS Gene and Gene Encoding Variant Soybean PPO

The levels of resistance to the PPO inhibitory-type herbicidal compound represented by structure 8 are confirmed quantitatively by testing the tobacco bearing the introduced present CTP-CP4 EPSPS gene and gene encoding the variant soybean PPO obtained in Example 50, and the control recombinant tobacco obtained in Example 5 according to the same manner as in Example 49.

Further the levels of resistance to glyphosate are confirmed quantitatively by testing the tobacco bearing introduced the present CTP-CP4 EPSPS gene and the gene encoding the variant soybean PPO and the control recombinant tobacco according to the same manner as in Example 49.

Example 52

Introduction of Present CTP-CP4 EPSPS Gene and Gene Encoding Variant *Chlamydomonas reinhardtii* PPO into Tobacco

A plasmid was constructed for introducing the

present CTP-CP4 EPSPS gene and a gene encoding a variant *Chlamydomonas reinhardtii* PPO into a plant by *Agrobacterium* infection method. The plasmid pTVCRP constructed in Example 16 was digested with the restriction enzymes BamHI and SacI to prepare a DNA fragment containing a gene encoding a variant *Chlamydomonas reinhardtii* PPO. Binary vector pBI121 (manufactured by Clontech) was digested with the restriction enzymes BamHI and SacI to remove β -glucuronidase gene and, instead thereof, the above DNA fragment containing the gene encoding the variant *Chlamydomonas reinhardtii* PPO was inserted therein to construct the plasmid pBICRP (Fig. 38) wherein the gene was ligated to the downstream of the 35S promoter.

Then, the plasmid pBICRP is digested with the restriction enzyme BamHI and nucleotides are added to the gap of the double-stranded DNA with DNA Polymerase I to blunt the end of the DNA. The 5'-end of the DNA is dephosphorylated by treatment with an alkaline phosphatase derived from calf small intestine, followed by inserting a phosphorylated KpnI linker (4668A manufactured by Takara Shuzo Co., Ltd.) and cyclization to obtain plasmid pBICRPK. Then, the plasmid pCENSK constructed in Example 47 is digested with the restriction enzyme KpnI to obtain a DNA fragment containing the present CTP-CP4 EPSPS gene, the terminator of the gene encoding nopaline synthase located

at the downstream of the present CTP-CP4 EPSPS gene and the 35S promoter located at the downstream of the terminator. This is inserted in the KpnI restriction site of the above plasmid pBICRPK to construct the plasmid pBICECRP (Fig. 39) wherein the present CTP-CP4 EPSPS gene and the gene encoding the variant *Chlamydomonas reinhardtii* PPO are ligated to the downstream of 35S promoter, respectively.

The plasmid pBICECRP is introduced into *Agrobacterium tumefaciens* LBA44044 and this is cultured in LB medium containing 300 µg/ml of streptomycin, 100 µg/ml of rifampicin and 25 µg/ml of kanamycin, followed by selection of a transformant to isolate an *Agrobacterium* strain bearing pBICECRP.

Tobacco leaf pieces cultivated sterilely are infected with the *Agrobacterium* strain and, according to the same manner as described in Example 5, tobacco bearing the introduced present CTP-CP4 EPSPS gene and gene encoding the variant *Chlamydomonas reinhardtii* PPO is obtained.

Example 53

Confirmation of Resistance to Herbicidal Compounds of Tobacco Bearing Introduced Present CTP-CP4 EPSPS Gene and Gene Encoding Variant *Chlamydomonas reinhardtii* PPO

The levels of resistance to the above PPO inhibitory-type herbicidal compound represented by

Structure 8 are confirmed quantitatively by testing the tobacco bearing the introduced present CTP-CP4 EPSPS gene and gene encoding the variant *Chlamydomonas reinhardtii* PPO obtained in Example 52 and the control recombinant tobacco obtained in Example 5 according to the same manner as in Example 49.

Further the levels of resistance to glyphosate are confirmed quantitatively by testing the tobacco bearing the introduced present CTP-CP4 EPSPS gene and gene encoding the variant *Chlamydomonas reinhardtii* PPO and the control recombinant tobacco according to the same manner as in Example 49.

Example 54

Introduction of Present CTP-CP4 EPSPS Gene and Gene Encoding Chloroplast-Localized Type Ferrochelatase of *Arabidopsis thaliana* into Tobacco

The plasmid pBIATF constructed in Example 39 is digested with the restriction enzyme BamHI and then nucleotides are added in the gap of the double-stranded DNA with DNA polymerase I to blunt the end of the DNA. The 5'-end of the DNA is dephosphorylated by treatment with an alkaline phosphatase derived from calf small intestine, followed by insertion of a phosphorylated KpnI linker (4668A manufactured by Takara Shuzo Co., Ltd.) and cyclization to obtain the plasmid pBIATFK. Then, the

plasmid pCENSK constructed in Example 47 is digested with the restriction enzyme KpnI to obtain a DNA fragment containing the present CTP-CP4 EPSPS gene, the terminator of the gene encoding nopaline synthase located at the downstream of the present CTP-CP4 EPSPS gene and the 35S promoter located at the downstream of the terminator. This is inserted in the KpnI restriction site of the above plasmid pBIATFK to construct the plasmid pBICEATF (Fig. 40) wherein the present CTP-CP4 EPSPS gene and the gene encoding the chloroplast-localized type ferrochelatase of *Arabidopsis thaliana* are ligated to the downstream of 35S promoter, respectively.

The plasmid pBICEATF is introduced into *Agrobacterium tumefaciens* LBA44044 and this is cultured in LB medium containing 300 µg/ml of streptomycin, 100 µg/ml of rifampicin and 25 µg/ml of kanamycin, followed by selection of a transformant to isolate an *Agrobacterium* strain bearing pBICEATF.

Tobacco leaf pieces cultivated sterily are infected with the *Agrobacterium* strain and, according to the same manner as described in Example 5, tobacco bearing the inserted present CTP-CP4 EPSPS gene and gene encoding the chloroplast-localized type ferrochelatase of *Arabidopsis thaliana* is obtained.

Confirmation of Resistance to Herbicidal Compounds of Tobacco Bearing Introduced Present CTP-CP4 EPSPS Gene and Gene Encoding Chloroplast-Localized Type Ferrochelatase of *Arabidopsis thaliana*

5 The levels of resistance to the above PPO inhibitory-type herbicidal compound represented by Structure 8 are confirmed quantitatively by testing the tobacco bearing the introduced present CTP-CP4 EPSPS gene and gene encoding the chloroplast-localized type
10 ferrochelatase of *Arabidopsis thaliana* obtained in Example 54, and the control recombinant tobacco obtained in Example 5 according to the same manner as in Example 49.

Further the levels of resistance to glyphosate are confirmed quantitatively by testing the tobacco bearing
15 the introduced present CTP-CP4 EPSPS gene and gene encoding the chloroplast-localized type ferrochelatase of *Arabidopsis thaliana* and the control recombinant tobacco according to the same manner as in Example 49.

Example 56

20 Introduction of Present CTP-CP4 EPSPS Gene and Gene Encoding Present Soybean CPOX into Tobacco

A plasmid was constructed for introducing the present CTP-CP4 EPSPS gene and a gene encoding the present soybean CPOX into a plant by *Agrobacterium* infection method.

25 First, the plasmid pCRSCPOX constructed in Example 42 was

digested with the restriction enzyme BamHI to prepare a DNA fragment containing a gene encoding the present soybean CPOX. The DNA fragment was inserted in the BamHI restriction site of the plasmid pBI121KS constructed in Example 9 to obtain the plasmid pBISCPOXGUS. This plasmid was digested with the restriction enzyme SalI to remove β -glucuronidase gene, followed by self-cyclization to construct the plasmid pBISCPOX (Fig. 41) wherein the gene was ligated to the downstream of the 35S promoter.

Then, the plasmid pBISCPOX is digested with the restriction enzyme BamHI and nucleotides are added to the gap of the double-stranded DNA with DNA polymerase I to blunt the end of the DNA. The 5'-end of the DNA is dephosphorylated by treatment with an alkaline phosphatase derived from calf small intestine, followed by inserting a phosphorylated KpnI linker (4668A manufactured by Takara Shuzo Co., Ltd.) therein and cyclization to obtain the plasmid pBISCPOXK. Then, the plasmid pCENSK constructed in Example 47 is digested with the restriction enzyme KpnI to obtain a DNA fraction containing the present CTP-CP4 EPSPS gene, the terminator of the gene encoding nopaline synthase located at the downstream of the present CTP-CP4 EPSPS gene and the 35S promoter located at the downstream of the terminator. This is inserted in the KpnI restriction site of the above plasmid pBISCPOXK to construct the plasmid

pBICESCPOX (Fig. 42) wherein the present CTP-CP4 EPSPS gene and the gene encoding the present soybean CPOX are ligated to the downstream of 35S promoter, respectively.

The plasmid pBICESCPOX is introduced into
5 *Agrobacterium tumefaciens* LBA44044 and this is cultured in LB medium containing 300 µg/ml of streptomycin, 100 µg/ml of rifampicin and 25 µg/ml of kanamycin, followed by selection of a transformant to isolate an *Agrobacterium* strain bearing pBICESCPOX.

10 Tobacco leaf pieces cultivated sterily are infected with the *Agrobacterium* strain and, according to the same manner as described in Example 45, tobacco bearing the inserted present CTP-CP4 EPSPS gene and gene encoding the present soybean CPOX is obtained.

15 Example 57

Confirmation of Resistance to Herbicidal Compounds of Tobacco Bearing Introduced Present CTP-CP4 EPSPS Gene and Gene Encoding Present Soybean CPOX

The levels of resistance to the above PPO
20 inhibitory-type herbicidal compound represented by Structure 8 are confirmed quantitatively by testing the tobacco bearing the introduced present CTP-CP4 EPSPS gene and gene encoding the present soybean CPOX obtained in Example 56, and the control recombinant tobacco according
25 to the same manner as in Example 49.

SEQUENCE LISTING FREE TEXT

SEQ ID NO: 1

Designed oligonucleotide primer to amplify bchH

5 gene

SEQ ID NO: 2

Designed oligonucleotide primer to amplify bchH

gene

SEQ ID NO: 3

Designed oligonucleotide primer to amplify

soybean PPO gene

SEQ ID NO: 4

Designed oligonucleotide primer to amplify

soybean PPO gene

SEQ ID NO: 7

Designed oligonucleotide primer to amplify bchH

gene

SEQ ID NO: 8

Designed oligonucleotide primer to amplify bchH

20 gene

SEQ ID NO: 9

Designed oligonucleotide primer to amplify DNA
fragment having partial sequence of tobacco chlH gene

SEQ ID NO: 10

25 Designed oligonucleotide primer to amplify DNA

09597719.102700

fragment having partial sequence of tobacco chlH gene

SEQ ID NO: 11

Designed oligonucleotide primer to amplify DNA

fragment having partial sequence of soybean PPO gene

5 SEQ ID NO: 12

Designed oligonucleotide primer to amplify DNA

fragment having partial sequence of soybean PPO gene

SEQ ID NO: 13

Designed oligonucleotide primer to amplify DNA

10 fragment having partial sequence of soybean PPO gene

SEQ ID NO: 14

Designed oligonucleotide primer to amplify DNA

fragment having partial sequence of soybean PPO gene

SEQ ID NO: 15

15 Designed oligonucleotide primer to amplify

Chlamydomonas PPO gene

SEQ ID NO: 16

Designed oligonucleotide primer to amplify

Chlamydomonas PPO gene

20 SEQ ID NO: 19

Designed oligonucleotide primer to amplify DNA

fragment having partial sequence of *Chlamydomonas* PPO gene

SEQ ID NO: 20

Designed oligonucleotide primer to amplify DNA

25 fragment having partial sequence of *Chlamydomonas* PPO gene

00202749-10200

Designed oligonucleotide primer to amplify DNA
having partial sequence of cucumber ferrochelatase

Designed oligonucleotide primer to amplify DNA having partial sequence of cucumber ferrochelatase

Designed oligonucleotide primer to amplify
dia coli hemF gene

Designed oligonucleotide primer to amplify
dia coli hemF gene

Designed oligonucleotide primer to amplify
dia coli hemF gene

Designed oligonucleotide primer to amplify
dia coli hemF gene

Designed oligonucleotides to synthesize genes
random peptides comprising 5 amino acids

Designed oligonucleotides to synthesize genes

encoding random peptides comprising 5 amino acids

SEQ ID NO: 29

Designed oligonucleotide to synthesize the gene
encoding the peptide HASYS

5 SEQ ID NO: 30

Designed oligonucleotide to synthesize the gene
encoding the peptide HASYS

SEQ ID NO: 31

10 Designed oligonucleotide to synthesize the gene
encoding the peptide RASSL

SEQ ID NO: 32

Designed oligonucleotide to synthesize the gene
encoding the peptide RASSL

SEQ ID NO: 33

15 Designed oligonucleotide to synthesize the gene
encoding the peptide MGHASYS

SEQ ID NO: 34

Designed oligonucleotide to synthesize the gene
encoding the peptide MGHASYS

20 SEQ ID NO: 35

Designed oligonucleotide to synthesize the gene
encoding the peptide MGRASSL

SEQ ID NO: 36

25 Designed oligonucleotide to synthesize the gene
encoding the peptide MGRASSL

SEQ ID NO: 37

Designed oligonucleotide to synthesize the gene
encoding the peptide MGYAGY

SEQ ID NO: 38

5 Designed oligonucleotide to synthesize the gene
encoding the peptide MGYAGY

SEQ ID NO: 39

Designed oligonucleotide to synthesize the gene
encoding the peptide MGYAGF

10 SEQ ID NO: 40

Designed oligonucleotide to synthesize the gene
encoding the peptide MGYAGF

SEQ ID NO: 41

15 Designed oligonucleotide to synthesize the gene
encoding the peptide MG(HASYS)4

SEQ ID NO: 42

Designed oligonucleotide to synthesize the gene
encoding the peptide MG(HASYS)4

SEQ ID NO: 43

20 Designed oligonucleotide to synthesize the gene
encoding the peptide MG(HASYS)4

SEQ ID NO: 44

Designed oligonucleotide to synthesize the gene
encoding the peptide MG(HASYS)4

25 SEQ ID NO: 45

Designed oligonucleotide to synthesize the gene
encoding the peptide MG(HASYS)8

SEQ ID NO: 46

5 Designed oligonucleotide to synthesize the gene
encoding the peptide MG(HASYS)8

SEQ ID NO: 47

Designed oligonucleotide to synthesize the gene
encoding the peptide MG(RASSL)4

SEQ ID NO: 48

10 Designed oligonucleotide to synthesize the gene
encoding the peptide MG(RASSL)4

SEQ ID NO: 49

Designed oligonucleotide to synthesize the gene
encoding the peptide MG(RASSL)4

5 SEQ ID NO: 50

Designed oligonucleotide to synthesize the gene
encoding the peptide MG(RASSL)4

SEQ ID NO: 51

20 Designed oligonucleotide to synthesize the gene
encoding the peptide MG(RASSL)8

SEQ ID NO: 52

Designed oligonucleotide to synthesize the gene
encoding the peptide MG(RASSL)8

SEQ ID NO: 53

25 Protoporphyrin IX binding protein HASYS

SEQ ID NO: 54

Protoporphyrin IX binding protein MGHASYS

SEQ ID NO: 55

Protoporphyrin IX binding protein RASSL

5 SEQ ID NO: 56

Protoporphyrin IX binding protein MGRASSL

SEQ ID NO: 57

H₂TMpyP binding protein YAGY

SEQ ID NO: 58

10 H₂TMpyP binding protein MGYAGY

SEQ ID NO: 59

H₂TMpyP binding protein YAGF

SEQ ID NO: 60

H₂TMpyP binding protein MGYAGF

15 SEQ ID NO: 61

Protoporphyrin IX binding protein MG(HASYS)₄

SEQ ID NO: 62

Protoporphyrin IX binding protein MG(HASYS)₈

SEQ ID NO: 63

20 Protoporphyrin IX binding protein MG(RASSL)₄

SEQ ID NO: 64

Protoporphyrin IX binding protein MG(RASSL)₈

SEQ ID NO: 65

25 Designed oligonucleotide primer to introduce
mutation into arabidopsis PPO gene

SEQ ID NO: 66

Designed oligonucleotide primer to amplify
herbicide resistant arabidopsis PPO(A220V) gene

SEQ ID NO: 67

5 Designed oligonucleotide primer to amplify
herbicide resistant arabidopsis PPO(A220V) gene

SEQ ID NO: 68

Designed oligonucleotide primer to amplify
arabidopsis chloroplast ferrochelatase gene

10 SEQ ID NO: 69

Designed oligonucleotide primer to amplify
arabidopsis chloroplast ferrochelatase gene

SEQ ID NO: 70

15 Designed oligonucleotide primer to amplify
soybean coproporphyrinogen III oxidase gene

SEQ ID NO: 71

Designed oligonucleotide primer to amplify
soybean coproporphyrinogen III oxidase gene

SEQ ID NO: 72

20 Designed oligonucleotide primer to amplify DNA
fragment having nucleotide sequence encoding the *Petunia*
hybrida EPSPS chloroplast transit peptide and the
Agrobacterium sp. strain CP4 EPSPS gene

SEQ ID NO: 73

25 Designed oligonucleotide primer to amplify DNA

fragment having nucleotide sequence encoding the Petunia hybrida EPSPS chloroplast transit peptide and the Agrobacterium sp. strain CP4 EPSPS gene

SEQ ID NO: 75

- 5 Designed oligonucleotide primer to amplify DNA fragment having partial sequence of soybean PPO gene

SEQ ID NO: 76

 Designed oligonucleotide primer to amplify DNA fragment having partial sequence of soybean PPO gene

006774-10200

SEQUENCE LISTING

<110> Hiroki NAKAJIMA et al.

5

<120> Method for giving resistance to weed control compounds to plants

<130>

10

<150> JP 10/120553

<151> 1998-04-30

<150> JP 10/281127

<151> 1998-10-02

15

<150> JP 10/330981

<151> 1998-11-20

<150> JP 11/054730

20

<151> 1999-03-02

<160> 76

<210> 1

25

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

5 <223> Designed oligonucleotide primer to amplify bchH gene

<400> 1

gacatctaga ggagacgacc atatgcacgg tgaagtctc 39

10 <210> 2

<211> 31

<212> DNA

<213> Artificial Sequence

15 <220>

<223> Designed oligonucleotide primer to amplify bchH gene

<400> 2

acggaagctt agatcttcac tcggcgcaa t 31

20

<210> 3

<211> 39

<212> DNA

<213> Artificial Sequence

<223> Designed oligonucleotide primer to amplify soybean PP0 gene

5 tegagetcca tggttttecgt cttcaacgag atcctattc 39

<211> 39

10 <213> Artificial Sequence

<223> Designed oligonucleotide primer to amplify soybean PP0 gene

ttgtcgacaa ctgctactat ttgtacactc tatttg 36

<211> 1632

<213> Glycine max var. Williams82

<221> CDS

25 $\langle 222 \rangle$ (1) ... (1632)

<400> 5

atg gtt tcc gtc ttc aac gag atc cta ttc ccg ccg aac caa acc ctt 48

Met Val Ser Val Phe Asn Glu Ile Leu Phe Pro Pro Asn Gln Thr Leu

1 5 10 15

5 ctt cgc ccc tcc ctc cat tcc cca acc tet ttc ttc acc tet ccc act 96

Leu Arg Pro Ser Leu His Ser Pro Thr Ser Phe Phe Thr Ser Pro Thr

20 25 30

cga aaa ttc cct cgc tct cgc cct aac cct att cta cgc tgc tcc att 144

Arg Lys Phe Pro Arg Ser Arg Pro Asn Pro Ile Leu Arg Cys Ser Ile

10 35 40 45

gcg gag gaa tcc acc gcg tct ccg ccc aaa acc aga gac tcc gcc ccc 192

Ala Glu Glu Ser Thr Ala Ser Pro Pro Lys Thr Arg Asp Ser Ala Pro

50 55 60

gtg gac tgc gtc gtc gtc ggc gga ggc gtc agc ggc ctc tgc atc gcc 240

15 Val Asp Cys Val Val Val Gly Gly Gly Val Ser Gly Leu Cys Ile Ala

65 70 75 80

cag gcc ctc gcc acc aaa cac gcc aat gcc aac gtc gtc gtc acg gag 288

Gln Ala Leu Ala Thr Lys His Ala Asn Ala Asn Val Val Val Thr Glu

85 90 95

20 gcc cga gac cgc gtc ggc ggc aac atc acc acg atg gag agg gac gga 336

Ala Arg Asp Arg Val Gly Gly Asn Ile Thr Thr Met Glu Arg Asp Gly

100 105 110

tac ctc tgg gaa gaa ggc ccc aac agc ttc cag cct tet gat cca atg 384

Tyr Leu Trp Glu Glu Gly Pro Asn Ser Phe Gln Pro Ser Asp Pro Met

25 115 120 125

0060771010200

	ctc acc atg gtg gtg gac agt ggt tta aag gat gag ctt gtt ttg ggg	432
	Leu Thr Met Val Val Asp Ser Gly Leu Lys Asp Glu Leu Val Leu Gly	
	130 135 140	
	gat cct gat gca cct cgg ttt gtg ttg tgg aac agg aag ttg agg ccg	480
5	Asp Pro Asp Ala Pro Arg Phe Val Leu Trp Asn Arg Lys Leu Arg Pro	
	145 150 155 160	
	gtg ccc ggg aag ctg act gat ttg cct ttc ttt gac ttg atg agc att	528
	Val Pro Gly Lys Leu Thr Asp Leu Pro Phe Phe Asp Leu Met Ser Ile	
	165 170 175	
10	ggg ggc aaa atc agg gct ggc ttt ggt gcg ctt gga att cgg cct cct	576
	Gly Gly Lys Ile Arg Ala Gly Phe Gly Ala Leu Gly Ile Arg Pro Pro	
	180 185 190	
	cct cca ggt cat gag gaa tcg gtt gaa gag ttt gtt cgt cgg aac ctt	624
	Pro Pro Gly His Glu Glu Ser Val Glu Glu Phe Val Arg Arg Asn Leu	
15	195 200 205	
	ggg gat gag gtt ttt gaa cgg ttg ata gag cct ttt tgt tca ggg gtc	672
	Gly Asp Glu Val Phe Glu Arg Leu Ile Glu Pro Phe Cys Ser Gly Val	
	210 215 220	
	tat gca ggc gat cct tca aaa tta agt atg aaa gca gca ttc ggg aaa	720
20	Tyr Ala Gly Asp Pro Ser Lys Leu Ser Met Lys Ala Ala Phe Gly Lys	
	225 230 235 240	
	gtt tgg aag ctg gaa aaa aat ggt ggt agc att att ggt gga act ttc	768
	Val Trp Lys Leu Glu Lys Asn Gly Gly Ser Ile Ile Gly Gly Thr Phe	
	245 250 255	

Lys Ala Ile Gln Glu Arg Asn Gly Ala Ser Lys Pro Pro Arg Asp Pro

270

5 Arg Leu Pro Lys Pro Lys Gly Gln Thr Val Gly Ser Phe Arg Lys Gly

285

Leu Thr Met Leu Pro Asp Ala Ile Ser Ala Arg Leu Gly Asn Lys Val

300

Lys Leu Ser Trp Lys Leu Ser Ser Ile Ser Lys Leu Asp Ser Gly Glu

305 310 315 320

Tyr Ser Leu Thr Tyr Glu Thr Pro Glu Gly Val Val Ser Leu Gln Cys

15 325 330 335

aaa act gtt gtc ctg acc att cct tcc tat gtt gct agt aca ttg ctg 1056

Lys Thr Val Val Leu Thr Ile Pro Ser Tyr Val Ala Ser Thr Leu Leu

350

cgt cct ctg tct gct gct gct gca gat gca ctt tca aag ttt tat tac 1104

20 Arg Pro Leu Ser Ala Ala Ala Ala Asp Ala Leu Ser Lys Phe Tyr Tyr

365

cct cca gtt gct gca gtt tcc ata tcc tat cca aaa gaa gct att aga 1152

Pro Pro Val Ala Ala Val Ser Ile Ser Tyr Pro Lys Glu Ala Ile Arg

380

tca gaa tgc ttg ata gat ggt gag ttg aag ggg ttt ggt caa ttg cat 1200
 Ser Glu Cys Leu Ile Asp Gly Glu Leu Lys Gly Phe Gly Gln Leu His
 385 390 395 400
 cca cgt agc caa gga gtg gaa aca tta gga act ata tac agc tca tca 1248
 5 Pro Arg Ser Gln Gly Val Glu Thr Leu Gly Thr Ile Tyr Ser Ser Ser
 405 410 415
 cta ttc ccc aac cga gca cca cct gga agg gtt cta ctc ttg aat tac 1296
 Leu Phe Pro Asn Arg Ala Pro Pro Gly Arg Val Leu Leu Leu Asn Tyr
 420 425 430
 10 att gga gga gca act aat act gga att tta tgc aag acg gac agt gaa 1344
 Ile Gly Gly Ala Thr Asn Thr Gly Ile Leu Ser Lys Thr Asp Ser Glu
 435 440 445
 ctt gtg gaa aca gtt gat cga gat ttg agg aaa atc ctt ata aac cca 1392
 Leu Val Glu Thr Val Asp Arg Asp Leu Arg Lys Ile Leu Ile Asn Pro
 15 450 455 460
 aat gcc cag gat cca ttt gta gtg ggg gtg aga ctg tgg cct caa gct 1440
 Asn Ala Gln Asp Pro Phe Val Val Gly Val Arg Leu Trp Pro Gln Ala
 465 470 475 480
 att cca cag ttc tta gtt ggc cat ctt gat ctt cta gat gtt gct aaa 1488
 20 Ile Pro Gln Phe Leu Val Gly His Leu Asp Leu Leu Asp Val Ala Lys
 485 490 495
 gct tct atc aga aat act ggg ttt gaa ggg ctc ttc ctt ggg ggt aat 1536
 Ala Ser Ile Arg Asn Thr Gly Phe Glu Gly Leu Phe Leu Gly Gly Asn
 500 505 510

00607719.102700

Ala Arg Asp Arg Val Gly Gly Asn Ile Thr Thr Met Glu Arg Asp Gly

100

105

110

Tyr Leu Trp Glu Glu Gly Pro Asn Ser Phe Gln Pro Ser Asp Pro Met

115

120

125

5

Leu Thr Met Val Val Asp Ser Gly Leu Lys Asp Glu Leu Val Leu Gly

130

135

140

Asp Pro Asp Ala Pro Arg Phe Val Leu Trp Asn Arg Lys Leu Arg Pro

145

150

155

160

Val Pro Gly Lys Leu Thr Asp Leu Pro Phe Phe Asp Leu Met Ser Ile

10

165

170

175

Gly Gly Lys Ile Arg Ala Gly Phe Gly Ala Leu Gly Ile Arg Pro Pro

180

185

190

Pro Pro Gly His Glu Glu Ser Val Glu Glu Phe Val Arg Arg Asn Leu

195

200

205

15

Gly Asp Glu Val Phe Glu Arg Leu Ile Glu Pro Phe Cys Ser Gly Val

210

215

220

Tyr Ala Gly Asp Pro Ser Lys Leu Ser Met Lys Ala Ala Phe Gly Lys

225

230

235

240

Val Trp Lys Leu Glu Lys Asn Gly Gly Ser Ile Ile Gly Gly Thr Phe

20

245

250

255

Lys Ala Ile Gln Glu Arg Asn Gly Ala Ser Lys Pro Pro Arg Asp Pro

260

265

270

Arg Leu Pro Lys Pro Lys Gly Gln Thr Val Gly Ser Phe Arg Lys Gly

275

280

285

00697719.102700

Leu Thr Met Leu Pro Asp Ala Ile Ser Ala Arg Leu Gly Asn Lys Val
 290 295 300
 Lys Leu Ser Trp Lys Leu Ser Ser Ile Ser Lys Leu Asp Ser Gly Glu
 305 310 315 320
 5 Tyr Ser Leu Thr Tyr Glu Thr Pro Glu Gly Val Val Ser Leu Gln Cys
 325 330 335
 Lys Thr Val Val Leu Thr Ile Pro Ser Tyr Val Ala Ser Thr Leu Leu
 340 345 350
 Arg Pro Leu Ser Ala Ala Ala Ala Asp Ala Leu Ser Lys Phe Tyr Tyr
 10 355 360 365
 Pro Pro Val Ala Ala Val Ser Ile Ser Tyr Pro Lys Glu Ala Ile Arg
 370 375 380
 Ser Glu Cys Leu Ile Asp Gly Glu Leu Lys Gly Phe Gly Gln Leu His
 385 390 395 400
 15 Pro Arg Ser Gln Gly Val Glu Thr Leu Gly Thr Ile Tyr Ser Ser Ser
 405 410 415
 Leu Phe Pro Asn Arg Ala Pro Pro Gly Arg Val Leu Leu Leu Asn Tyr
 420 425 430
 Ile Gly Gly Ala Thr Asn Thr Gly Ile Leu Ser Lys Thr Asp Ser Glu
 20 435 440 445
 Leu Val Glu Thr Val Asp Arg Asp Leu Arg Lys Ile Leu Ile Asn Pro
 450 455 460
 Asn Ala Gln Asp Pro Phe Val Val Gly Val Arg Leu Trp Pro Gln Ala
 465 470 475 480

10

490

15

20

505

510

25

520

525

<210> 7

<211> 39

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<213> Designed oligonucleotide primer to amplify bchH gene

<400> 7

gacatctagt ctagacgacc atatgcacgg tgaagtctc 39

<210> 8

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to amplify bchH gene

5 <400> 8

acggaagctt ggtacctcac tcggcgga t 31

<210> 9

<211> 35

10 <212> DNA

<213> Artificial Sequence

<220>

15 <223> Designed oligonucleotide primer to amplify DNA fragment having
partial sequence of tobacco chlH gene

<400> 9

ccaatgtaat gctatggtac ctatgttatt cactc 35

20 <210> 10

<211> 34

<212> DNA

<213> Artificial Sequence

25 <220>

<223> Designed oligonucleotide primer to amplify DNA fragment having partial sequence of tobacco chlH gene

<400> 10

5 gagatcattc tttttgctgt cgacttatcg atcg 34

<210> 11

<211> 39

<212> DNA

10 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to amplify DNA fragment having partial sequence of soybean PPO gene

15

<400> 11

ggcggaggcg tcaccatggt ctgcacgcc caggcc 36

<210> 12

20 <211> 36

<212> DNA

<213> Artificial Sequence

<220>

25 <223> Designed oligonucleotide primer to amplify DNA fragment having

00597719.102700

partial sequence of soybean PPO gene

<400> 12

gcctgcagggt cgacaactgc tactatttgt acactc 36

5

<210> 13

<211> 33

<212> DNA

<213> Artificial Sequence

10

<220>

<223> Designed oligonucleotide primer to amplify DNA fragment having
partial sequence of soybean PPO gene

15

<400> 13

cacaggaag gtaccatggt ctgcatgcc cag 33

<210> 14

<211> 33

20

<212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to amplify DNA fragment having
partial sequence of soybean PPO gene

25

<400> 14

cetgcagctc gagagctcct actatttgta cac 33

5 <210> 15

<211> 28

<212> DNA

<213> Artificial Sequence

10 <220>

<223> Designed oligonucleotide primer to amplify Chlamydomonas PPO
gene

<400> 15

15 aatgatgttg acccagactc ctgggacc 28

<210> 16

<211> 27

<212> DNA

20 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to amplify Chlamydomonas PPO
gene

25

<400> 16

tactacacat cccagcaagc gccaatg 27

<210> 17

5 <211> 1838

<212> DNA

<213> Chlamydomonas reinhardtii CC407

<220>

10 <221> CDS

<222> (2)... (1693)

<400> 17

a atg atg ttg acc cag act cct ggg acc gcc acg gct tct agc cgg 46

15 Met Met Leu Thr Gln Thr Pro Gly Thr Ala Thr Ala Ser Ser Arg

1 5 10 15

cgg tcg cag atc cgc tcg gct gcg cac gtc tcc gcc aag gtc gcg cct 94

Arg Ser Gln Ile Arg Ser Ala Ala His Val Ser Ala Lys Val Ala Pro

20 25 30

20 cgg ccc acg cca ttc tcg gtc gcg agc ccc gcg acc gct gcg agc ccc 142

Arg Pro Thr Pro Phe Ser Val Ala Ser Pro Ala Thr Ala Ala Ser Pro

35 40 45

gcg acc gcg gcg gcc cgc cgc aca ctc cac cgc act gct gcg gcg gcc 190

Ala Thr Ala Ala Ala Arg Arg Thr Leu His Arg Thr Ala Ala Ala Ala

25 50 55 60

act ggt gct ccc acg gcg tcc gga gcc ggc gtc gcc aag acg etc gac 238
 Thr Gly Ala Pro Thr Ala Ser Gly Ala Gly Val Ala Lys Thr Leu Asp
 65 70 75
 aat gtg tat gac gtg atc gtg gtc ggt gga ggt ctc tcg ggc ctg gtg 286
 5 Asn Val Tyr Asp Val Ile Val Val Gly Gly Gly Leu Ser Gly Leu Val
 80 85 90 95
 acc ggc cag gcc ctg gcg gct cag cac aaa att cag aac ttc ctt gtt 334
 Thr Gly Gln Ala Leu Ala Ala Gln His Lys Ile Gln Asn Phe Leu Val
 100 105 110
 10 acg gag gct cgc gag cgc gtc ggc ggc aac att acg tcc atg tcg ggc 382
 Thr Glu Ala Arg Glu Arg Val Gly Gly Asn Ile Thr Ser Met Ser Gly
 115 120 125
 gat ggc tac gtg tgg gag gag ggc ccg aac agc ttc cag ccc aac gat 430
 Asp Gly Tyr Val Trp Glu Glu Gly Pro Asn Ser Phe Gln Pro Asn Asp
 15 130 135 140
 agc atg ctg cag att gcg gtg gac tct ggc tgc gag aag gac ctt gtg 478
 Ser Met Leu Gln Ile Ala Val Asp Ser Gly Cys Glu Lys Asp Leu Val
 145 150 155
 ttc ggt gac ccc acg gct ccc cgc ttc gtg tgg tgg gag ggc aag ctg 526
 20 Phe Gly Asp Pro Thr Ala Pro Arg Phe Val Trp Trp Glu Gly Lys Leu
 160 165 170 175
 cgc ccc gtg ccc tcg ggc ctg gac gcc ttc acc ttc gac etc atg tcc 574
 Arg Pro Val Pro Ser Gly Leu Asp Ala Phe Thr Phe Asp Leu Met Ser
 180 185 190

atc ccc ggc aag atc cgc gcc ggg ctg ggc gcc atc ggc ctc atc aac 622
 Ile Pro Gly Lys Ile Arg Ala Gly Leu Gly Ala Ile Gly Leu Ile Asn
 195 200 205
 gga gcc atg ccc tcc ttc gag gag agt gtg gag cag ttc atc cgc cgc 670
 5 Gly Ala Met Pro Ser Phe Glu Glu Ser Val Glu Gln Phe Ile Arg Arg
 210 215 220
 aac ctg ggc gat gag gtg ttc ttc cgc ctg atc gag ccc ttc tgc tcc 718
 Asn Leu Gly Asp Glu Val Phe Phe Arg Leu Ile Glu Pro Phe Cys Ser
 225 230 235
 10 ggc gtg tac gcg ggc gac ccc tcc aag ctg tcc atg aag gcg gcc ttc 766
 Gly Val Tyr Ala Gly Asp Pro Ser Lys Leu Ser Met Lys Ala Ala Phe
 240 245 250 255
 aac agg atc tgg att ctg gag aag aac ggc ggc agc ctg gtg gga ggt 814
 Asn Arg Ile Trp Ile Leu Glu Lys Asn Gly Gly Ser Leu Val Gly Gly
 15 260 265 270
 gcc atc aag ctg ttc cag gaa cgc cag tcc aac ccg gcc ccg ccg cgg 862
 Ala Ile Lys Leu Phe Gln Glu Arg Gln Ser Asn Pro Ala Pro Pro Arg
 275 280 285
 gac ccg cgc ctg ccg ccc aag ccc aag ggc cag acg gtg ggc tgc ttc 910
 20 Asp Pro Arg Leu Pro Pro Lys Pro Lys Gly Gln Thr Val Gly Ser Phe
 290 295 300
 cgc aag ggc ctg aag atg ctg ccg gac gcc att gag cgc aac atc ccc 958
 Arg Lys Gly Leu Lys Met Leu Pro Asp Ala Ile Glu Arg Asn Ile Pro
 305 310 315

gac aag atc cgc gtg aac tgg aag ctg gtg tct ctg ggc cgc gag gcg 1006
 Asp Lys Ile Arg Val Asn Trp Lys Leu Val Ser Leu Gly Arg Glu Ala
 320 325 330 335
 gac ggg cgg tac ggg ctg gtg tac gac acg ccc gag ggc cgt gtc aag 1054
 5 Asp Gly Arg Tyr Gly Leu Val Tyr Asp Thr Pro Glu Gly Arg Val Lys
 340 345 350
 gtg ttt gcc cgc gcc gtg gct ctg acc gcg ccc agc tac gtg gtg gcg 1102
 Val Phe Ala Arg Ala Val Ala Leu Thr Ala Pro Ser Tyr Val Val Ala
 355 360 365
 10 gac ctg gtc aag gag cag gcg ccc gcc gcc gcc gag gcc ctg gcc tcc 1150
 Asp Leu Val Lys Glu Gln Ala Pro Ala Ala Ala Glu Ala Leu Gly Ser
 370 375 380
 ttc gac tac ccg ccg gtg ggc gcc gtg acg ctg tgg tac ccg ctg agc 1198
 Phe Asp Tyr Pro Pro Val Gly Ala Val Thr Leu Ser Tyr Pro Leu Ser
 15 385 390 395
 gcc gtg cgg gag gag cgc aag gcc tgg gac ggg tcc gtg ccg gcc ttc 1246
 Ala Val Arg Glu Glu Arg Lys Ala Ser Asp Gly Ser Val Pro Gly Phe
 400 405 410 415
 ggt cag ctg cac ccg cgc acg cag ggc atc acc act ctg gcc acc atc 1294
 20 Gly Gln Leu His Pro Arg Thr Gln Gly Ile Thr Thr Leu Gly Thr Ile
 420 425 430
 tac agc tcc agc ctg ttc ccc gcc cgc gcg ccc gag gcc cac atg ctg 1342
 Tyr Ser Ser Ser Leu Phe Pro Gly Arg Ala Pro Glu Gly His Met Leu
 435 440 445

ctg ctc aac tac atc ggc ggc acc acc aac cgc ggc atc gtc aac cag 1390
 Leu Leu Asn Tyr Ile Gly Gly Thr Thr Asn Arg Gly Ile Val Asn Gln
 450 455 460
 acc acc gag cag ctg gtg gag cag gtg gac aag gac ctg cgc aac atg 1438
 5 Thr Thr Glu Gln Leu Val Glu Gln Val Asp Lys Asp Leu Arg Asn Met
 465 470 475
 gtc atc aag ccc gac gcg ccc aag ccc cgt gtg gtg ggc gtg cgc gtg 1486
 Val Ile Lys Pro Asp Ala Pro Lys Pro Arg Val Val Gly Val Arg Val
 480 485 490 495
 10 tgg ccg cgc gcc atc ccg cag ttc aac ctg ggc cac ctg gag cag ctg 1534
 Trp Pro Arg Ala Ile Pro Gln Phe Asn Leu Gly His Leu Glu Gln Leu
 500 505 510
 gac aag gcg cgc aag gcg ctg gac gcg gcg ggc ctg cag gcc gtg cac 1582
 Asp Lys Ala Arg Lys Ala Leu Asp Ala Ala Gly Leu Gln Gly Val His
 15 515 520 525
 ctg ggg ggc aac tac gtc agc ggt gtg gcc ctg ggc aag gtg gtg gag 1630
 Leu Gly Gly Asn Tyr Val Ser Gly Val Ala Leu Gly Lys Val Val Glu
 530 535 540
 cac ggc tac gag tcc gca gcc aac ctg gcc aag agc gtg tcc aag gcc 1678
 20 His Gly Tyr Glu Ser Ala Ala Asn Leu Ala Lys Ser Val Ser Lys Ala
 545 550 555
 gca gtc aag gcc taa gcgctgcag cagtagcagc agcagcatcg ggctgtagct 1733
 Ala Val Lys Ala
 560 563
 25 ggtaaatgcc gcagtggcac cggcagcagc aattggcaag cacttggggc aagcggagtg 1793

gaggcgaggg gggggctacc attggcgctt gctgggatgt gtagt

1838

<210> 18

<211> 563

5 <212> PRT

<213> Chlamydomonas reinhardtii CC407

<400> 18

10 Met Met Leu Thr Gln Thr Pro Gly Thr Ala Thr Ala Ser Ser Arg
 1 5 10 15
 Arg Ser Gln Ile Arg Ser Ala Ala His Val Ser Ala Lys Val Ala Pro
 20 25 30
 Arg Pro Thr Pro Phe Ser Val Ala Ser Pro Ala Thr Ala Ala Ser Pro
 15 35 40 45
 Ala Thr Ala Ala Ala Arg Arg Thr Leu His Arg Thr Ala Ala Ala Ala
 50 55 60
 Thr Gly Ala Pro Thr Ala Ser Gly Ala Gly Val Ala Lys Thr Leu Asp
 65 70 75
 20 Asn Val Tyr Asp Val Ile Val Val Gly Gly Gly Leu Ser Gly Leu Val
 80 85 90 95
 Thr Gly Gln Ala Leu Ala Ala Gln His Lys Ile Gln Asn Phe Leu Val
 100 105 110
 Thr Glu Ala Arg Glu Arg Val Gly Gly Asn Ile Thr Ser Met Ser Gly
 25 115 120 125

Asp Gly Tyr Val Trp Glu Glu Gly Pro Asn Ser Phe Gln Pro Asn Asp
 130 135 140
 Ser Met Leu Gln Ile Ala Val Asp Ser Gly Cys Glu Lys Asp Leu Val
 145 150 155
 5 Phe Gly Asp Pro Thr Ala Pro Arg Phe Val Trp Trp Glu Gly Lys Leu
 160 165 170 175
 Arg Pro Val Pro Ser Gly Leu Asp Ala Phe Thr Phe Asp Leu Met Ser
 180 185 190
 Ile Pro Gly Lys Ile Arg Ala Gly Leu Gly Ala Ile Gly Leu Ile Asn
 10 195 200 205
 Gly Ala Met Pro Ser Phe Glu Glu Ser Val Glu Gln Phe Ile Arg Arg
 210 215 220
 Asn Leu Gly Asp Glu Val Phe Phe Arg Leu Ile Glu Pro Phe Cys Ser
 225 230 235
 15 Gly Val Tyr Ala Gly Asp Pro Ser Lys Leu Ser Met Lys Ala Ala Phe
 240 245 250 255
 Asn Arg Ile Trp Ile Leu Glu Lys Asn Gly Gly Ser Leu Val Gly Gly
 260 265 270
 Ala Ile Lys Leu Phe Gln Glu Arg Gln Ser Asn Pro Ala Pro Pro Arg
 20 275 280 285
 Asp Pro Arg Leu Pro Pro Lys Pro Lys Gly Gln Thr Val Gly Ser Phe
 290 295 300
 Arg Lys Gly Leu Lys Met Leu Pro Asp Ala Ile Glu Arg Asn Ile Pro
 305 310 315

Asp Lys Ile Arg Val Asn Trp Lys Leu Val Ser Leu Gly Arg Glu Ala
 320 325 330 335
 Asp Gly Arg Tyr Gly Leu Val Tyr Asp Thr Pro Glu Gly Arg Val Lys
 340 345 350
 5 Val Phe Ala Arg Ala Val Ala Leu Thr Ala Pro Ser Tyr Val Val Ala
 355 360 365
 Asp Leu Val Lys Glu Gln Ala Pro Ala Ala Ala Glu Ala Leu Gly Ser
 370 375 380
 Phe Asp Tyr Pro Pro Val Gly Ala Val Thr Leu Ser Tyr Pro Leu Ser
 10 385 390 395
 Ala Val Arg Glu Glu Arg Lys Ala Ser Asp Gly Ser Val Pro Gly Phe
 400 405 410 415
 Gly Gln Leu His Pro Arg Thr Gln Gly Ile Thr Thr Leu Gly Thr Ile
 420 425 430
 15 Tyr Ser Ser Ser Leu Phe Pro Gly Arg Ala Pro Glu Gly His Met Leu
 435 440 445
 Leu Leu Asn Tyr Ile Gly Gly Thr Thr Asn Arg Gly Ile Val Asn Gln
 450 455 460
 Thr Thr Glu Gln Leu Val Glu Gln Val Asp Lys Asp Leu Arg Asn Met
 20 465 470 475
 Val Ile Lys Pro Asp Ala Pro Lys Pro Arg Val Val Gly Val Arg Val
 480 485 490 495
 Trp Pro Arg Ala Ile Pro Gln Phe Asn Leu Gly His Leu Glu Gln Leu
 500 505 510

Asp Lys Ala Arg Lys Ala Leu Asp Ala Ala Gly Leu Gln Gly Val His

515

520

525

Leu Gly Gly Asn Tyr Val Ser Gly Val Ala Leu Gly Lys Val Val Glu

530

535

540

5 His Gly Tyr Glu Ser Ala Ala Asn Leu Ala Lys Ser Val Ser Lys Ala

545

550

555

Ala Val Lys Ala

560

563

10 <210> 19

<211> 32

<212> DNA

<213> Artificial Sequence

15 <220>

<223> Designed oligonucleotide primer to amplify DNA fragment having
partial sequence of Chlamydomonas PPO gene

<400> 19

20 ggtcggtgga ggggatccga tgctggtgac cg 32

<210> 20

<211> 32

<212> DNA

25 <213> Artificial Sequence

0022016142390

<220>

<223> Designed oligonucleotide primer to amplify DNA fragment having partial sequence of Chlamydomonas PPO gene

5

<400> 20

gctactgctg cgagctctta ggccttgact gc 32

<210> 21

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to amplify DNA fragment having partial sequence of cucumber ferrochelataase gene

<400> 21

getttagaat cggatcctat ggcagtggat gac 33

<210> 22

<211> 36

<212> DNA

<213> Artificial Sequence

25

<220>

<223> Designed oligonucleotide primer to amplify DNA fragment having partial sequence of cucumber ferrochelatase gene

5 <400> 22

ggtgaacttc tatttgagct ctcaggtaaa tataag 36

<210> 23

<211> 25

10 <212> DNA

<213> Artificial Sequence

<220>

15 <223> Designed oligonucleotide primer to amplify Esherichia coli hemF gene

<400> 23

gctgaaggcg tgatcagtta ttcc 25

20 <210> 24

<211> 24

<212> DNA

<213> Artificial Sequence

25 <220>

<223> Designed oligonucleotide primer to amplify Esherichia coli hemF
gene

<400> 24

5 catcagcctg cagtgcgaaa agtg 24

<210> 25

<211> 26

<212> DNA

10 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to amplify Esherichia coli hemF
gene

15

<400> 25

cgaaaaaggg atccgttatg aaaccc 26

<210> 26

20

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

25

<223> Designed oligonucleotide primer to amplify Esherichia coli hemF

gene

<400> 26

gctgtttttcc gagctcccg t cac 23

5

<210> 27

<211> 22

<212> DNA

<213> Artificial Sequence

10

<220>

<223> Designed oligonucleotides to synthesize genes encoding random peptides comprising 5 amino acids

15

<400> 27

tggccnnknn knnknnknk gc 22

<210> 28

<211> 29

20

<212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotides to synthesize genes encoding random peptides comprising 5 amino acids

25

<400> 28

ggccgcmnnm nnnnnnnnnn nggccagct 29

5 <210> 29

<211> 22

<212> DNA

<213> Artificial Sequence

10 <220>

<223> Designed oligonucleotide to synthesize the gene encoding the peptide HASYS

<400> 29

15 tggcccatgc tagttagtcg gc 22

<210> 30

<211> 29

<212> DNA

20 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide to synthesize the gene encoding the peptide HASYS

25

<400> 30

tggcgccgac taactagcat gggccagct 29

<210> 31

5 <211> 22

<212> DNA

<213> Artificial Sequence

<220>

10 <223> Designed oligonucleotide to synthesize the gene encoding the peptide RASSL

<400> 31

tgccccgggc gtcgtcgttg gc 22

15

<210> 32

<211> 29

<212> DNA

<213> Artificial Sequence

20

<220>

<223> Designed oligonucleotide to synthesize the gene encoding the peptide RASSL

25

<400> 32

ggccgccaac gacgacgcc gggccagct 29

<210> 33

<211> 26

5 <212> DNA

<213> Artificial Sequence

<220>

10 <223> Designed oligonucleotide to synthesize the gene encoding the
peptide MGHASYS

<400> 33

catgggtcac gettcttact cctaag 26

15 <210> 34

<211> 26

<212> DNA

<213> Artificial Sequence

20 <220>

<223> Designed oligonucleotide to synthesize the gene encoding the
peptide MGHASYS

<400> 34

25 aattcttagg agtaagaagc gtgacc 26

<210> 35

<211> 26

<212> DNA

5 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide to synthesize the gene encoding the
peptide MGRASSL

10

<400> 35

catgggtcgt gcttcttccc tgtaag 26

<210> 36

15

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

20

<223> Designed oligonucleotide to synthesize the gene encoding the
peptide MGRASSL

<400> 36

aattcttaca gggaagaagc acgacc 26

25

<211> 23

<212> DNA

<213> Artificial Sequence

5 <220>

<223> Designed oligonucleotide to synthesize the gene encoding the peptide MGYAGF

<400> 39

10 catgggttac gctggcttct aag 23

<210> 40

<211> 23

<212> DNA

15 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide to synthesize the gene encoding the peptide MGYAGF

20

<400> 40

aattcttaga agccagcgta acc 23

<210> 41

25 <211> 34

<212> DNA

<213> Artificial Sequence

<220>

5 <223> Designed oligonucleotide to synthesize the gene encoding the
peptide MG(HASYS)4

<400> 41

catgggtcac gcttcttact cccatgcac ttac 34

<210> 42

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide to synthesize the gene encoding the
peptide MG(HASYS)4

<400> 42

gtgggagtaa gatgcatggg agtaagaagc gtgacc 36

<210> 43

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide to synthesize the gene encoding the
5 peptide MG (HASYS)4

<400> 43

tcccacgctt cttactccca tgcattctac tcctaag 37

10 <210> 44

<211> 35

<212> DNA

<213> Artificial Sequence

15 <220>

<223> Designed oligonucleotide to synthesize the gene encoding the
peptide MG (HASYS)4

<400> 44

20 aattcttagg agtaagatgc atgggagtaa gaagc 35

<210> 45

<211> 30

<212> DNA

25 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide to synthesize the gene encoding the
peptide MG(HASYS)8

5

<400> 45

tcccaagctt cttactccca tgcatttac 30

<210> 46

10 <211> 30

<212> DNA

<213> Artificial Sequence

<220>

15 <223> Designed oligonucleotide to synthesize the gene encoding the
peptide MG(HASYS)8

<400> 46

gtgggagtaa gatgcatggg agtaagaagc 30

20

<210> 47

<211> 34

<212> DNA

<213> Artificial Sequence

25

<220>

<223> Designed oligonucleotide to synthesize the gene encoding the peptide MG(RASSL)4

5 <400> 47

catgggtcgt gctttctccc tgcgcgcac ttcc 34

<210> 48

<211> 36

10 <212> DNA

<213> Artificial Sequence

<220>

15 <223> Designed oligonucleotide to synthesize the gene encoding the peptide MG(RASSL)4

<400> 48

acgcagggaa gatgcgcgca gggagaagaac acgacc 36

20 <210> 49

<211> 37

<212> DNA

<213> Artificial Sequence

25 <220>

<223> Designed oligonucleotide to synthesize the gene encoding the peptide MG(RASSL)4

<400> 49

5 ctgcgtgctt cttccctgcg cgcattcttc ctgtaag 37

<210> 50

<211> 35

<212> DNA

10 <213> Artificial Sequence

<220>

<223> Designed oligonucleotide to synthesize the gene encoding the peptide MG(RASSL)4

15

<400> 50

aattcttaca gggaagatgc ggcagaggaa gaagc 35

<210> 51

20 <211> 30

<212> DNA

<213> Artificial Sequence

<220>

25 <223> Designed oligonucleotide to synthesize the gene encoding the

peptide MG(RASSL)8

<400> 51

ctgcgtgctt cttccctgag cgcattctcc 30

5

<210> 52

<211> 30

<212> DNA

<213> Artificial Sequence

10

<220>

<223> Designed oligonucleotide to synthesize the gene encoding the peptide MG(RASSL)8

15

<400> 52

acgcagggaa gatgcgcgca gggaagaagc 30

<210> 53

<211> 5

20

<212> PRT

<213> Artificial Sequence

<220>

<223> Protoporphyrin IX binding protein HASYS

25

<400> 53

His Ala Ser Tyr Ser

1 5

5 <210> 54

<211> 7

<212> PRT

<213> Artificial Sequence

10 <220>

<223> Protoporphyrin IX binding protein MGHASYS

<400> 54

Met Gly His Ala Ser Tyr Ser

15 1 5

<210> 55

<211> 5

<212> PRT

20 <213> Artificial Sequence

<220>

<223> Protoporphyrin IX binding protein RASSL

25 <400> 55

0022036722560

Arg Ala Ser Ser Leu

1 5

<210> 56

5 <211> 7

<212> PRT

<213> Artificial Sequence

<220>

10 <223> Protoporphyrin IX binding protein MGRASSL

<400> 56

Met Gly Arg Ala Ser Ser Leu

1 5

15

<210> 57

<211> 4

<212> PRT

<213> Artificial Sequence

20

<220>

<223> H₂TMpyP binding protein YAGY.

<400> 57

25 Tyr Ala Gly Tyr 4

<210> 58

<211> 6

<212> PRT

5 <213> Artificial Sequence

<220>

<223> H₂TMpyP binding protein MGYAGY

10 <400> 58

Met Gly Tyr Ala Gly Tyr

1

5

<210> 59

15 <211> 4

<212> PRT

<213> Artificial Sequence

<220>

20 <223> H₂TMpyP binding protein YAGF

<400> 59

Tyr Ala Gly Phe

1

25

<210> 60

<211> 6

<212> PRT

<213> Artificial Sequence

5

<220>

<223> H₂TMpyP binding protein MGYAGF

<400> 60

10 Met Gly Tyr Ala Gly Phe

1

5

<210> 61

<211> 22

15 <212> PRT

<213> Artificial Sequence

<220>

<223> Protoporphyrin IX binding protein MG(HASYS)₄

20

<400> 61

Met Gly His Ala Ser Tyr Ser His Ala Ser Tyr Ser His Ala Ser Tyr

1

5

10

15

Ser His Ala Ser Tyr Ser

25

20

<210> 62

<211> 42

<212> PRT

5 <213> Artificial Sequence

<220>

<223> Protoporphyrin IX binding protein MG(HASYS)₈

10 <400> 62

Met Gly His Ala Ser Tyr Ser His Ala Ser Tyr Ser His Ala Ser Tyr

1 5 10 15

Ser His Ala Ser Tyr Ser His Ala Ser Tyr Ser His Ala Ser Tyr Ser

20 25 30

15 His Ala Ser Tyr Ser His Ala Ser Tyr Ser

35 40

<210> 63

<211> 22

20 <212> PRT

<213> Artificial Sequence

<220>

<223> Protoporphyrin IX binding protein MG(RASSL)₄

25

<400> 63

Met Gly Arg Ala Ser Ser Leu Arg Ala Ser Ser Leu Arg Ala Ser Ser

1 5 10 15

Leu Arg Ala Ser Ser Leu

5 20

<210> 64

<211> 42

<212> PRT

10 <213> Artificial Sequence

<220>

<223> Protoporphyrin IX binding protein MG(RASSL)₈.

15 <400> 64

Met Gly Arg Ala Ser Ser Leu Arg Ala Ser Ser Leu Arg Ala Ser Ser

1 5 10 15

Leu Arg Ala Ser Ser Leu Arg Ala Ser Ser Leu Arg Ala Ser Ser Leu

20 25 30

20 Arg Ala Ser Ser Leu Arg Ala Ser Ser Leu

35 40

<210> 65

<211> 36

25 <212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to introduce mutation into
5 arabidopsis PPO gene

<400> 65

tggtcagggtg tttatgttgg tgatccttca aaactg 36

10 <210> 66

<211> 32

<212> DNA

<213> Artificial Sequence

15 <220>

<223> Designed oligonucleotide primer to amplify herbicide resistant
arabidopsis PPO(A220V) gene

<400> 66

ccatgcggaa gcttatggag ttatctcttc tc 32

<210> 67

<211> 34

<212> DNA

25 <213> Artificial Sequence

<220>

<213> Designed oligonucleotide primer to amplify herbicide resistant
arabidopsis PP0(A220V) gene

5

<400> 67

gggagattta atgtcgacca ttacttgta agcg 34

<210> 68

10

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

15

<223> Designed oligonucleotide primer to amplify Arabidopsis
chloroplast ferrochelataase gene

<400> 68

gatcggttct gaaatttgga tccatgcagg c 31

20

<210> 69

<211> 31

<212> DNA

<213> Artificial Sequence

25

<220>

<223> Designed oligonucleotide primer to amplify Arabidopsis
chloroplast ferrochelatase gene

5 <400> 69

cacaaaacca acgagctcct ataggttcg g 31

<210> 70

<211> 30

10 <212> DNA

<213> Artificial Sequence

<220>

15 <223> Designed oligonucleotide primer to amplify soybean
coproporphyrinogen III oxidase gene

<400> 70

gaatcggatc cgaagcatga tgcattgtgc 30

20 <210> 71

<211> 30

<212> DNA

<213> Artificial Sequence

25 <220>

<400> 71

<210> 72

<212> DNA

 $\langle 220 \rangle$

<400> 72

20 <210> 73

<211> 32

<212> DNA

<213> Artificial Sequence

25 <220>

<223>Designed oligonucleotide primer to amplify DNA fragment having nucleotide sequence encoding the *Petunia hybrida* EPSPS chloroplast transit peptide and the *Agrobacterium* sp. strain CP4 EPSPS gene

5 <400> 73

gagtcgacgg tcatacaggca gccttcgtat cg 32

<210> 74

<211> 1587

10 <212> DNA

<213> *Petunia hybrida* EPSPS chloroplast transit peptide and *Agrobacterium* sp. strain CP4 EPSPS

<220>

15 <221> CDS

<222> (1)... (1581)

<400> 74

atg gca caa att aac aac atg gct caa ggg ata caa acc ctt aat ccc 48

20 Met Ala Gln Ile Asn Asn Met Ala Gln Gly Ile Gln Thr Leu Asn Pro

1 5 10 15

aat tcc aat ttc cat aaa ccc caa gtt cct aaa tct tca agt ttt ctt 96

Asn Ser Asn Phe His Lys Pro Gln Val Pro Lys Ser Ser Ser Phe Leu

20

25

30

	gtt ttt gga tct aaa aaa ctg aaa aat tca gca aat tct atg ttg gtt	144
	Val Phe Gly Ser Lys Lys Leu Lys Asn Ser Ala Asn Ser Met Leu Val	
	35 40 45	
	ttg aaa aaa gat tca att ttt atg caa aag ttt tgt tcc ttt agg att	192
5	Leu Lys Lys Asp Ser Ile Phe Met Gln Lys Phe Cys Ser Phe Arg Ile	
	50 55 60	
	tca gca tca gtg gct aca gcc tgc atg ctt cac ggt gca agc agc cgg	240
	Ser Ala Ser Val Ala Thr Ala Cys Met Leu His Gly Ala Ser Ser Arg	
	65 70 75	
10	ccc gca acc gcc cgc aaa tcc tct ggc ctt tcc gga acc gtc cgc att	288
	Pro Ala Thr Ala Arg Lys Ser Ser Gly Leu Ser Gly Thr Val Arg Ile	
	80 85 90 95	
	ccc ggc gac aag tcg atc tcc cac cgg tcc ttc atg ttc ggc ggt ctc	336
	Pro Gly Asp Lys Ser Ile Ser His Arg Ser Phe Met Phe Gly Gly Leu	
15	100 105 110	
	gcg agc ggt gaa acg cgc atc acc ggc ctt ctg gaa ggc gag gac gtc	384
	Ala Ser Gly Glu Thr Arg Ile Thr Gly Leu Leu Glu Gly Glu Asp Val	
	115 120 125	
	atc aat acg ggc aag gcc atg cag gcc atg ggc gcc agg atc cgt aag	432
20	Ile Asn Thr Gly Lys Ala Met Gln Ala Met Gly Ala Arg Ile Arg Lys	
	130 135 140	
	gaa ggc gac acc tgg atc atc gat ggc gtc ggc aat ggc ggc ctc ctg	480
	Glu Gly Asp Thr Trp Ile Ile Asp Gly Val Gly Asn Gly Gly Leu Leu	
	145 150 155	

<212> DNA

<213> Artificial Sequence

<220>

5 <223> Designed oligonucleotide primer to amplify DNA fragment having
partial sequence of soybean PPO gene

<400> 75

cacaggaaag gtaccatggt ctgcatgcc cag 33

<210> 76

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Designed oligonucleotide primer to amplify DNA fragment having
partial sequence of soybean PPO gene

<400> 76

cctgcagctc gagagctcct actatttgta cac 33

What is claimed is:

1. A method for giving resistance to weed control compounds to plants which comprises the steps of:

introducing a gene encoding a protein having the following characteristics (a) to (c):

(a) having a specific affinity for a substance which is concerned with the weed control activity of a weed control compound,

(b) having substantially no capability of modifying a substance for which said protein has a specific affinity, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin, into a plant cell; and

expressing the gene.

2. The method according to claim 1, wherein the gene is introduced into the plant cell in the form that it is operably ligated to a promoter and a terminator both of which are functional in the plant cell.

3. The method according to claim 1 or 2, wherein the substance which is concerned with the weed control activity of the weed control compound is the weed control compound itself.

4. The method according to claim 1, wherein the substance which is concerned with the weed control activity

of a weed control compound is an endogenous substance in a plant.

5 5. The method according to claim 1, wherein the weed control compound is that inhibiting porphyrin biosynthesis of a plant.

6. The method according to claim 1, wherein the weed control compound is a protoporphyrinogen IX oxidase inhibitory-type herbicidal compound.

10 7. The method according to claim 5 or 6, wherein the substance which is concerned with the weed control activity of the weed control compound is protoporphyrin IX.

15 8. The method according to claim 5 or 6, wherein the protein is protoporphyrin IX binding subunit protein of magnesium chelatase, or a variant of said protein having a specific affinity for protoporphyrin IX.

9. The method according to claim 8, wherein the protein is magnesium chelatase derived from a photosynthetic microorganism.

20 10. The method according to claim 8, wherein the protein is magnesium chelatase derived from a plant.

11. The method according to claim 8, wherein the protein is magnesium chelatase derived from tobacco.

25 12. The method according to claim 5 or 6, wherein the protein comprises the amino acid sequence of SEQ

ID NO: 53.

13. The method according to claim 5 or 6,
wherein the protein has the amino acid sequence of SEQ ID
NO: 54.

5 14. The method according to claim 5 or 6,
wherein the protein comprises the amino acid sequence of SEQ
ID NO: 55.

0950719-102700
10 15. The method according to claim 5 or 6,
wherein the protein has the amino acid sequence of SEQ ID
NO: 56.

16. The method according to claim 5 or 6,
wherein the protein comprises the amino acid sequence of SEQ
ID NO: 57.

15 17. The method according to claim 5 or 6,
wherein the protein has the amino acid sequence of SEQ ID
NO: 58.

18. The method according to claim 5 or 6,
wherein the protein comprises the amino acid sequence of SEQ
ID NO: 59.

20 19. The method according to claim 5 or 6,
wherein the protein has of the amino acid sequence of SEQ ID
NO: 60.

20. The method according to claim 5 or 6,
wherein the protein is composed of 4 to 100 amino acids.

25 21. The method according to claim 5 or 6,

wherein the substance which is concerned with the weed control activity of the weed control compound is protoporphyrinogen IX.

22. The method according to claim 5 or 6,
5 wherein the protein is a variant of protoporphyrinogen IX oxidase having no capability of oxidizing protoporphyrinogen IX and having a specific affinity for a protoporphyrinogen IX.

23. The method according to claim 5 or 6,
10 wherein the protein is a variant of protoporphyrinogen IX oxidase having no capability of oxidizing protoporphyrinogen IX and having a specific affinity for a protoporphyrin IX oxidase inhibitory-type herbicidal compound.

24. The method according to claim 22 or 23,
15 wherein the protein is a variant of protoporphyrinogen IX oxidase derived from a plant.

25. The method according to claim 22 or 23,
wherein the protein is a variant of protoporphyrinogen IX
20 oxidase derived from soybean.

26. The method according to claim 22 or 23,
wherein the protein is a variant of protoporphyrinogen IX
oxidase derived from an algae.

27. The method according to claim 22 or 23,
25 wherein the protein is a variant of protoporphyrinogen IX

oxidase derived from *Chlamydomonas*.

28. A method for giving resistance to weed control compounds to plants which comprises the steps of:

introducing a gene encoding a protein having the following characteristics (a) to (c):

(a) having a specific affinity for protoporphyrin IX,

(b) having substantially no capability of modifying protoporphyrinogen IX, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin, into a plant cell; and

expressing the gene.

29. The method according to claim 28, wherein the gene is introduced in the plant cell in the form that it is operably ligated to a promoter and a terminator both of which are functional in the plant cell.

30. The method according to claim 28, wherein the weed control compound is that inhibiting porphyrin biosynthesis of a plant.

31. The method according to claim 28, wherein the weed control compound is a protoporphyrinogen IX oxidase inhibitory-type herbicidal compound.

32. The method according to claim 30 or 31, wherein the protein is magnesium chelatase or a variant of

said protein having a specific affinity for protoporphyrin IX.

33. The method according to claim 30 or 31, wherein the protein is ferrochelatase or a variant of said protein having a specific affinity for protoporphyrin IX.

34. The method according to claim 30 or 31, wherein the protein is ferrochelatase derived from a plant.

35. The method according to claim 30 or 31, wherein the protein is ferrochelatase derived from barley.

36. The method according to claim 30 or 31, wherein the protein is ferrochelatase derived from cucumber.

37. The method according to claim 30 or 31, wherein the protein is a peptide composed of 4 to 100 amino acids.

38. A method for giving resistance to weed control compounds to plants which comprises the steps of:

introducing a gene encoding a protein having the following characteristics (a) to (c):

(a) having a specific affinity for protoporphyrinogen IX,

(b) having the capability for modifying coproporphyrinogen III, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin; into a plant cell; and

expressing the gene.

39. The method according to claim 38, wherein the gene is introduced into the plant cell in the form that it is operably ligated to a promoter and a terminator both of which are functional in the plant cell.

40. The method according to claim 38, wherein the protein is coproporphyrinogen III oxidase or a variant of said protein having a specific affinity for protoporphyrinogen IX.

41. The method according to claim 38, wherein the protein is coproporphyrinogen III oxidase derived from a microorganism.

42 The method according to claim 38, wherein the protein is coproporphyrinogen III oxidase derived from *Escherichia coli*.

43. A weed control compound-resistant plant whose resistance is given by the method of claim 1 or 28.

44. A weed control compound-resistant plant whose resistance is given by the method of claim 38.

45. A method for protecting a plant which comprises applying the weed control compound to a growth area of the plant of claim 43.

46. A method for protecting a plant which comprises applying said weed control compound to a growth area of the plant of claim 44.

47. A method for selecting a plant which comprises applying a weed control compound to which the plant of claim 43 is resistant to a growth area of the plant of claim 43 and other plants, and selecting either
5 plant on the basis of difference in growth between the plants.

48. A method for selecting a plant which comprises applying a weed control compound to which the plant of claim 44 is resistant to a growth area of the
10 plant of claim 44 and other plants, and selecting either plant on the basis of difference in growth between the plants.

49. The method according to claim 47, wherein the plants are plant cells.


15 50. The method according to claim 48, wherein the plants are plant cells.

51. The method according to claim 1 or 2, wherein the weed control compound is a protoporphyrinogen IX oxidase inhibitory-type herbicidal compound selected
20 from the compounds of (1) to (3) below, and the substance which is concerned with the weed control activity of the weed control compound is protoporphyrin IX, protoporphyrinogen IX or a protoporphyrinogen IX oxidase inhibitory-type herbicidal compound:


25 (1) chlormethoxynil, bifenox, chlornitrofen,

acifluorfen and its ethyl ester, acifluorfen-sodium, oxyfluorfen, oxadiazon, 2-[4-chloro-2-fluoro-5-(prop-2-ynyloxy)phenyl]-2,3,4,5,6,7-hexahydro-1H-isoindol-1,3-dione, chlorophthalim, TNPP-ethyl, or N3-(1-phenylethyl)-2,6-dimethyl-5-propionylnicotinamide;


(2) a compound represented by the general formula: J-G (I), wherein G is a group represented by any one of the following general formulas G-1 to G-9 and J is a group represented by any one of the following general formulas J-1 to J-30:




G-1




G-2




G-3




G-4




G-5




G-6




G-7




G-8




G-9




J-1




J-2




J-3




J-4




J-5




J-6




J-7




J-8




J-9




J-10




J-11




J-12




J-13




J-14




J-15




J-16




J-17




J-18




J-19




J-20




J-21




J-22




J-23




J-24




J-25




J-26




J-27



J-28



J-29



J-30

wherein the dotted lines in the formulas J-5, J-6,
J-12 and J-24 represent that the left hand ring contains
only single bonds, or one bond in the ring is a double bond
between carbon atoms;

X is oxygen atom or sulfur atom;

Y is oxygen atom or sulfur atom;

R¹ is hydrogen atom or halogen atom;

R² is hydrogen atom, C₁-C₈alkyl group, C₁-C₈
haloalkyl group, halogen atom, OH group, OR²⁷ group, SH
group, S(O)_pR²⁷ group, COR²⁷ group, CO₂R²⁷ group, C(O)SR²⁷
group, C(O)NR²⁹R³⁰ group, CHO group, CR²⁷=NOR³⁶ group,
CH=CR³⁷CO₂R²⁷ group, CH₂CHR³⁷CO₂R²⁷ group, CO₂N=CR³¹R³² group,
nitro group, cyano group, NHSO₂R³³ group, NHSO₂NHR³³ group,
NR²⁷R³⁸ group, NH₂ group or phenyl group optionally

substituted with one or more and the same or different C_1 - C_4 alkyl groups;

p is 0, 1 or 2;

R^3 is C_1 - C_2 alkyl group, C_1 - C_2 haloalkyl group,
5 OCH₃ group, SCH₃ group, OCHF₂ group, halogen atom, cyano group or nitro group;

R^4 is hydrogen atom, C_1 - C_3 alkyl group, C_1 - C_3 haloalkyl group or halogen atom;

R^5 is hydrogen atom, C_1 - C_3 alkyl group, halogen
10 atom, C_1 - C_3 haloalkyl group, cyclopropyl group, vinyl group, C_2 alkynyl group, cyano group, C(O) R^{38} group, CO₂ R^{38} group, C(O)NR³⁸R³⁹ group, CR³⁴R³⁵CN group, CR³⁴R³⁵C(O) R^{38} group, CR³⁴R³⁵CO₂ R^{38} group, CR³⁴R³⁵C(O)NR³⁸R³⁹ group, CHR³⁴OH group, CHR³⁴OC(O) R^{38} group or OCHR³⁴OC(O)NR³⁸R³⁹ group, or, when G is
15 G-2 or G-6, R^4 and R^5 may form C=O group together with the carbon atom to which they are attached;

R^6 is C_1 - C_6 alkyl group, C_1 - C_6 haloalkyl group, C_2 - C_6 alkoxyalkyl group, C_3 - C_6 alkenyl group or C_3 - C_6 alkynyl group;

20 X^1 is single bond, oxygen atom, sulfur atom, NH group, N(C_1 - C_3 alkyl) group, N(C_1 - C_3 haloalkyl) group or N(allyl) group;

R^7 is hydrogen atom, C_1 - C_6 alkyl group, C_1 - C_6 haloalkyl group, halogen atom, S(O)₂(C_1 - C_6 alkyl) group or
25 C(=O) R^{40} group;

R^{14} is C_1-C_6 alkyl group, C_1-C_6 alkylthio group, C_1-C_6 haloalkyl group or $N(CH_3)_2$ group;

W is nitrogen atom or CR^{15} ;

R^{15} is hydrogen atom, C_1-C_6 alkyl group, halogen atom, or phenyl group optionally substituted with C_1-C_6 alkyl group, one or two halogen atoms, C_1-C_6 alkoxy group or CF_3 group;

each Q is independently oxygen atom or sulfur atom;

10 Q^1 is oxygen atom or sulfur atom;

Z^1 is $CR^{16}R^{17}$ group, oxygen atom, sulfur atom, $S(O)$ group, $S(O)_2$ group or $N(C_1-C_6\text{alkyl})$ group;

each R^{16} is independently hydrogen atom, halogen atom, hydroxyl group, C_1-C_6 alkoxy group, C_1-C_6 haloalkyl group, C_1-C_6 haloalkoxy group, C_2-C_6 alkylcarbonyloxy group or C_2-C_6 haloalkylcarbonyloxy group;

each R^{17} is independently hydrogen atom, hydroxyl group or halogen atom;

20 R^{18} is C_1-C_6 alkyl group, halogen atom or C_1-C_6 haloalkyl group;

R^{19} and R^{20} are independently hydrogen atom, C_1-C_6 alkyl group, or C_1-C_6 haloalkyl group;

Z^2 is oxygen atom, sulfur atom, NR^9 group or CR^9R^{10} group;

25 R^{21} and R^{22} are independently C_1-C_6 alkyl group, C_1-

C₆ haloalkyl group, C₃-C₆ alkenyl group, C₃-C₆ haloalkenyl group, C₃-C₆ alkynyl group or C₃-C₆ haloalkynyl group;

R²³ is hydrogen atom, halogen atom or cyano group;


5 R²⁴ is C₁-C₆ alkylsulfonyl group, C₁-C₆ alkyl group, C₁-C₆ haloalkyl group, C₃-C₆ alkenyl group, C₃-C₆ alkynyl group, C₁-C₆ alkoxy group, C₁-C₆ haloalkoxy group or halogen atom;

10 R²⁵ is C₁-C₆ alkyl group, C₁-C₆ haloalkyl group, C₃-C₆ alkenyl group or C₃-C₆ alkynyl group;


R²⁶ is C₁-C₆ alkyl group, C₁-C₆ haloalkyl group or phenyl group optionally substituted with C₁-C₆ alkyl, one or two halogen atoms, one or two nitro groups, C₁-C₆ alkoxy group or CF₃ group;

15 W¹ is nitrogen atom or CH group;

T is a group represented by any one of the following general formulas T-1, T-2 and T-3;



T-1



T-2



T-3

20 (wherein E¹, E², E³, E⁴, E⁵, E⁶, E⁷, E⁸, E⁹, E¹⁰, E¹¹ and E¹² are independently hydrogen atom or C₁-C₃ alkyl group);

R²⁷ is C₁-C₈ alkyl group, C₃-C₈ cycloalkyl group,

C_3-C_8 alkenyl group, C_3-C_8 alkynyl group, C_1-C_8 haloalkyl group,
 C_2-C_8 alkoxyalkyl group, C_2-C_8 alkylthioalkyl group, C_2-C_8
 alkylsulfinylalkyl group, C_2-C_8 alkylsulfonylalkyl group,
 C_1-C_8 alkylsulfonyl group, phenylsulfonyl group whose phenyl
 5 ring may be substituted with at least one substituent
 selected from the group consisting of halogen atom and C_1-C_4
 alkyl group, C_4-C_8 alkoxyalkoxyalkyl group, C_4-C_8
 cycloalkylalkyl group, C_6-C_8 cycloalkoxyalkyl group, C_4-C_8
 alkenyloxyalkyl group, C_4-C_8 alkynyloxyalkyl group, C_3-C_8
 10 haloalkoxyalkyl group, C_4-C_8 haloalkenyloxyalkyl group, C_4-C_8
 haloalkynyloxyalkyl group, C_6-C_8 cycloalkylthioalkyl group,
 C_4-C_8 alkenylthioalkyl group, C_4-C_8 alkynylthioalkyl group,
 C_1-C_4 alkyl group substituted with phenoxy group whose ring
 is substituted with at least one substituent selected from
 15 the group consisting of halogen atom, C_1-C_3 alkyl group and
 C_1-C_3 haloalkyl group, benzyloxy group whose ring is
 substituted with at least one substituent selected from the
 group consisting of halogen atom, C_1-C_3 alkyl group and C_1-C_3
 haloalkyl group, C_4-C_8 trialkylsilylalkyl group, C_3-C_8
 20 cyanoalkyl group, C_3-C_8 halocycloalkyl group, C_3-C_8
 haloalkenyl group, C_5-C_8 alkoxyalkenyl group, C_5-C_8
 haloalkoxyalkenyl group, C_5-C_8 alkylthioalkenyl group, C_3-C_8
 haloalkynyl group, C_5-C_8 alkoxyalkynyl group, C_5-C_8
 haloalkoxyalkynyl group, C_5-C_8 alkylthioalkynyl group, C_2-C_8
 25 alkylcarbonyl group, benzyl group whose ring is substituted

with at least one substituent selected from the group consisting of halogen atom, C₁-C₃ alkyl group and C₁-C₃ haloalkyl group, CHR³⁴COR²⁸ group, CHR³⁴COOR²⁸ group, CHR³⁴P(O)(OR²⁸)₂ group, CHR³⁴P(S)(OR²⁸)₂ group, CHR³⁴C(O)NR²⁹R³⁰ group or CHR³⁴C(O)NH₂ group;

R²⁸ is C₁-C₆ alkyl group, C₂-C₆ alkenyl group, C₃-C₆ alkynyl group or tetrahydrofuranyl group;

R²⁹ and R³¹ are independently hydrogen atom or C₁-C₄ alkyl group;

R³⁰ and R³² are independently C₁-C₄ alkyl group or phenyl group whose ring may be substituted with at least one substituent selected from the group consisting of halogen atom, C₁-C₃ alkyl group and C₁-C₃ haloalkyl group; or,

R²⁹ and R³⁰ together may form -(CH₂)₅-, -(CH₂)₄- or -CH₂CH₂OCH₂CH₂-, or the ring thus formed may be substituted with at least one substituent selected from the group consisting of C₁-C₃ alkyl group, phenyl group and benzyl group; or,

R³¹ and R³² may from C₃-C₆ cycloalkyl group together with the carbon atom to which they are attached;

R³³ is C₁-C₄ alkyl group, C₁-C₄ haloalkyl group or C₃-C₆ alkenyl group;

R³⁴ and R³⁵ are independently hydrogen atom or C₁-C₄ alkyl group;

R³⁶ is hydrogen atom, C₁-C₆ alkyl group, C₃-C₆

alkenyl group or C₃-C₆ alkynyl group;

R³⁷ is hydrogen atom, C₁-C₄ alkyl group or halogen atom;

R³⁸ is hydrogen atom, C₁-C₆ alkyl group, C₃-C₆ cycloalkyl group, C₃-C₆ alkenyl group, C₃-C₆ alkynyl group, C₂-C₆ alkoxyalkyl group, C₁-C₆ haloalkyl group, phenyl group whose ring may be substituted with at least one substituent selected from the group consisting of halogen atom, C₁-C₄ alkyl group and C₁-C₄ alkoxy group, -CH₂CO₂(C₁-C₄ alkyl) group or -CH(CH₃)CO₂(C₁-C₄ alkyl) group;


R³⁹ is hydrogen atom, C₁-C₂ alkyl group or C(O)O(C₁-C₄ alkyl) group;

R⁴⁰ is hydrogen atom, C₁-C₆ alkyl group, C₁-C₆ alkoxy group or NH(C₁-C₆ alkyl) group;

R⁴¹ is C₁-C₆ alkyl group, C₁-C₆ haloalkyl group, C₁-C₆ alkoxy group, NH(C₁-C₆ alkyl) group, phenyl group whose ring may be substituted with one substituent selected from the group consisting of R⁴² group, benzyl group and C₂-C₆ dialkylamino group; and

R⁴² is C₁-C₆ alkyl group, one or two halogen atoms, C₁-C₆ alkoxy group or CF₃ group;

(3) a compound of the formula (II):



or nipilacrofen,

wherein R^{43} is C_1-C_4 alkyl group;

R^{44} is C_1-C_4 alkyl group, C_1-C_4 alkylthio group, C_1-C_4 alkoxy group, C_1-C_4 haloalkyl group, C_1-C_4 haloalkylthio group or C_1-C_4 haloalkoxy group;









R^{43} and R^{44} together may form $-(CH_2)_3-$ or $-(CH_2)_4-$;

R^{45} is hydrogen atom or halogen atom;

R^{46} is hydrogen atom or C_1-C_4 alkyl group;

R^{47} is hydrogen atom, nitro group, cyano group, $-COOR^{49}$ group, $-C(=X)NR^{50}R^{51}$ group or $-C(=X^2)R^{52}$ group;

R^{48} is hydrogen atom, halogen atom, cyano group, C_1-C_4 alkyl group optionally substituted with at least one substituent selected from the group consisting of halogen atom and hydroxyl group, C_1-C_4 alkoxy group, phenyl group optionally substituted with at least one substituent selected from the group consisting of halogen atom, nitro group, cyano group, C_1-C_4 alkyl group, C_1-C_4 alkoxy group and halo- C_1-C_4 alkyl group, pyrrolyl group, C_2-C_8 alkyl group, C_3-C_8 alkenyl group, C_3-C_8 alkynyl group, C_3-C_8 alkoxy group, a group selected from the group consisting of C_2-C_8 alkyl group, C_3-C_8 alkenyl group, C_3-C_8 alkynyl group and C_3-C_8 alkoxy group into which at least one oxygen atom is inserted, or any one of groups represented by the following formulas:



wherein R^{49} , R^{50} and R^{52} are, the same or different, hydrogen atom or $\text{C}_1\text{--C}_4$ alkyl group;

R^{50} and R^{51} may form saturated alicyclic 5 or 6 membered ring together with the nitrogen atom to which they are attached;

R^{52} is hydrogen atom, $\text{C}_1\text{--C}_4$ alkyl group or $\text{C}_1\text{--C}_4$ alkyl group substituted with at least one halogen atom;

R⁵³ is hydrogen atom, C₁-C₆ alkyl group optionally substituted with at least one halogen atom, C₂-C₆ alkenyl group optionally substituted with at least one halogen atom, C₃-C₆ alkynyl group optionally substituted with at least one
 5 halogen atom, phenyl group optionally substituted with at least one halogen atom, C₃-C₈ cycloalkyl group, cyanomethyl group, or R⁶³CO- group;

R⁵⁴ is hydrogen atom, C₁-C₆ alkyl group optionally substituted with at least one halogen atom, C₂-C₆ alkenyl
 10 group optionally substituted with at least one halogen atom, C₃-C₆ alkynyl group optionally substituted with at least one halogen atom, phenyl group optionally substituted with halogen atom, C₃-C₈ cycloalkyl group, cyanomethyl group, C₁-C₄ alkoxy-C₁-C₆ alkyl group, di-C₁-C₄ alkylamino-C₁-C₄ alkyl
 15 group, tetrahydrofurfurylmethyl group, C₃-C₆ alkynyloxy-C₁-C₄ alkyl group, benzyl whose ring may be substituted with substituent selected from the group consisting of halogen atom, nitro group, cyano group, C₁-C₄ alkyl group, C₁-C₄ alkoxy group and halo-C₁-C₄ alkyl group, -C(=X²)R⁶³ group, -
 20 (CH₂)_a-(O)_a-R⁷⁰ group, -(CH₂)_a-O-(CH₂)_b-R⁷⁰ group, -(CH₂)_a-X²-R⁷⁶ group;

R⁵³ and R⁵⁴ together with the nitrogen atom to which they are attached may form saturated alicyclic 3, 5 or 6 membered ring or aromatic 5 or 6 membered ring
 25 which a carbon atom may be optionally replaced with oxygen

atom;

R^{55} is hydrogen atom, C_1-C_4 alkyl group, C_2-C_6 alkenyl group or C_3-C_6 alkynyl group, or R^{55} and R^{56} together may form $-(CH_2)_a-$;

5 R^{56} and R^{57} are independently C_1-C_4 alkyl group optionally substituted with at least one halogen atom, C_2-C_6 alkenyl group optionally substituted with at least one halogen atom, C_3-C_6 alkynyl optionally substituted with at least one halogen atom or phenyl group optionally substituted with at least one halogen atom, hydrogen atom, 10 C_3-C_6 cycloalkyl group, $-XR^{60}$ group or $-NR^{61}R^{62}$ group;

R^{58} is hydrogen atom, C_1-C_6 alkyl group, C_2-C_6 alkenyl group, C_3-C_6 alkynyl group, C_1-C_4 alkylcarbonyl group, cyano- C_1-C_3 alkyl group, C_1-C_4 alkoxy carbonyl- C_1-C_4 alkyl 15 group, di- C_1-C_4 alkoxy carbonyl- C_1-C_4 alkyl group, benzyl group, C_1-C_4 alkoxy- C_1-C_4 alkynyl group, $-(CH_2)_a-R^{75}$ group, $-(CH_2)_a-X^2-R^{72}$ group, $-(CH_2)_a-X^2-(CH_2)_b-R^{72}$ group or $-(CH_2)_a-X^2-(CH_2)_b-R^{72}$ group;

R^{59} is hydrogen atom, C_1-C_4 alkyl group, C_2-C_6 20 alkenyl group, C_3-C_6 alkynyl group, cyano- C_1-C_3 alkyl group, C_1-C_4 alkylcarbonyl- C_1-C_3 alkyl group or phenyl group;

R^{60} is C_1-C_4 alkyl group optionally substituted with at least one halogen atom;

R^{61} and R^{62} are, the same or different, hydrogen 25 atom or C_1-C_4 alkyl group;

R^{63} is C_1-C_4 alkyl group optionally substituted with at least one halogen atom, C_1-C_4 alkoxy- C_1-C_4 alkyl group, C_1-C_4 alkylthio- C_1-C_4 alkyl group, C_3-C_6 cycloalkyl group, phenyl group whose ring may be substituted with one
 5 substituent selected from the group consisting of halogen atom, nitro group, cyano group, C_1-C_4 alkyl group, C_1-C_4 alkoxy group and halo- C_1-C_4 alkyl group, $-NR^{73}R^{74}$ group or $-(CH_2)_a-(O)_d-R^{75}$ group;

R^{64} is C_1-C_4 alkoxycarbonyl group or carboxyl
 10 group;

R^{65} is chloromethyl group, cyanomethyl group, C_3-C_6 cycloalkyl group into which at least one oxygen atom may be inserted, or C_1-C_4 alkoxycarbonyl- C_1-C_4 alkyl group;

R^{66} is hydroxyl group or $-NR^{67}R^{68}$ group;

A is $-NR^{67}R^{68}$ group or $-S(O)_2-R^{69}$ group;

R^{67} and R^{68} are, the same or different, hydrogen atom or C_1-C_4 alkyl group;

R^{69} is C_1-C_4 alkyl group or C_1-C_4 haloalkyl group;

R^{70} is hydrogen atom, hydroxyl group, halogen
 20 atom, C_1-C_4 alkyl group optionally substituted with at least one C_1-C_4 alkoxy group, C_3-C_6 cycloalkyl group into which at least one oxygen atom may be inserted, C_3-C_6 cycloalkyl group optionally substituted with one or two methyl groups, furyl group, thienyl group or $-C(=O)R^{71}$ group;

R^{71} and R^{72} are, the same or different, C_1-C_4 alkyl
 25

group or C₁-C₄ alkoxy group;

R⁷³ and R⁷⁴ are, the same or different, C₁-C₄ alkyl group or phenyl group;

R⁷⁵ is C₃-C₆ cycloalkyl into which at least one oxygen atom may be inserted, C₃-C₆ cycloalkyl group optionally substituted with one or two methyl groups, furyl group, thienyl group or -C(=O)R⁷¹ group;

R⁷⁶ is C₁-C₄ alkyl group;

a, b and c is independently 1, 2 or 3;

d is 0 or 1;

e is 2 or 3;

f is 1 or 2; and

X² is oxygen atom or sulfur atom.

52. The method according to claim 1, additionally comprising the steps of:

introducing into the plant cell, a second gene selected from a gene encoding a protein substantially having protoporphyrinogen oxidase activity, a gene encoding a protein substantially having 5-enolpyruvylshikamate-3-phosphate synthase activity and a gene encoding a protein substantially having glyphosate oxidoreductase activity; and

expressing said second gene.

53. A plant cell having:

a gene encoding a protein having the following

characteristics (a) to (c):

(a) having a specific affinity for a substance which is concerned with the weed control activity of a weed control compound,

5 (b) having substantially no capability of modifying a substance for which said protein has a specific affinity, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin; and

10 at least one altered form of an enzymatic activity which gives a resistance to a weed control compound in an amount inhibiting a naturally occurring form of said enzymatic activity, wherein said altered form of an enzymatic activity is a form of enzymatic activity selected
15 from a protoporphyrinogen oxidase activity, 5-enolpyruvylshikamate-3-phosphate synthase activity and glyphosate oxidoreductase activity.

54. A plant cell having:

20 a gene encoding a protein having the following characteristics (a) to (c):

(a) having a specific affinity for a substance which is concerned with the weed control activity of a weed control compound,

25 (b) having substantially no capability of modifying a substance for which said protein has a specific

00507719-102700

affinity, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin; and

an altered protoporphyrinogen oxidase activity

5 which gives a resistance to a weed control compound in an amount inhibiting a natural occurring protoporphyrinogen oxidase activity.

55. A plant cell having:

6697719.102700
10 a gene encoding a protein having the following characteristics (a) to (c):

(a) having a specific affinity for a substance which is concerned with the weed control activity of a weed control compound,

15 (b) having substantially no capability of modifying a substance for which said protein has a specific affinity, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin; and

20 an altered 5-enolpyruvylshikamate-3-phosphate synthase activity which gives a resistance to a weed control compound in an amount inhibiting a natural occurring 5-enolpyruvylshikamate-3-phosphate synthase activity.

56. A plant cell having:

25 a gene encoding a protein having the following

characteristics (a) to (c):

(a) having a specific affinity for a substance which is concerned with the weed control activity of a weed control compound,

5 (b) having substantially no capability of modifying a substance for which said protein has a specific affinity, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin; and

10 an altered glyphosate oxidoreductase activity which gives a resistance to a weed control compound in an amount inhibiting a natural occurring glyphosate oxidoreductase activity.

15 57. The plant cell according to claim 53, wherein said altered form of an enzymatic activity is conferred by a second gene selected from a gene encoding a protein substantially having a protoporphyrinogen oxidase activity, a gene encoding a protein substantially having 5-enolpyruvylshikimate-3-phosphate synthase activity and a
20 gene encoding a protein substantially having glyphosate oxidoreductase activity.

58. The plant cell according to claim 57, wherein the gene encoding a protein having the following characteristics (a) to (c):

25 (a) having a specific affinity for a substance

which is concerned with the weed control activity of a weed control compound,

(b) having substantially no capability of modifying a substance for which said protein has a specific affinity, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin; and the second gene are introduced into the plant cell in the form in that both of said genes are operably ligated to a promoter and a terminator both of which are functional in said plant cell.

59. The plant cell according to claim 57, wherein the protein substantially having a protoporphyrinogen IX oxidase activity is protoporphyrinogen IX oxidase, the protein substantially having a 5-enol-pyruvylshikamate-3-phosphate synthase activity is 5-enolpyruvylshikamate-3-phosphate synthase and the protein substantially having glyphosate oxidoreductase activity is glyphosate oxidoreductase.

60. The plant cell according to claim 53, wherein the plant cell is derived from dicotyledones or monocotyledones.

61. A plant comprising the plant cell of claim 54.

62. A plant comprising the plant cell of claim

55.

63. A plant comprising the plant cell of claim

56.

64. A method for protecting a plant which

5 comprises applying a protoporphyrinogen IX oxidase
inhibitory-type compound to a growth area of the plant of
claim 61.

65. A method for protecting a plant which

10 comprises applying a protoporphyrinogen IX oxidase
inhibitory-type compound and a compound inhibiting 5-
enolpyruvylshikamate-3-phosphate synthase to a growth area
of the plant of claim 62.

66. A method for protecting a plant which

15 comprises applying a protoporphyrinogen IX oxidase
inhibitory-type compound and a compound inhibiting 5-
enolpyruvylshikamate-3-phosphate synthase to a growth area
of the plant of claim 63.

67. A method for selecting a plant which

20 comprises applying a protoporphyrinogen IX oxidase
inhibitory-type compound to a growth area of the plant of
claim 61 and other plants, and selecting either plant on
the basis of difference in growth between the plants.

68. A method for selecting a plant which

25 comprises applying a protoporphyrinogen IX oxidase
inhibitory-type compound and a compound inhibiting 5-

enolpyruvylshikamate-3-phosphate synthase to a growth area of the plant of claim 62 and other plants, and selecting either plant on the basis of difference in growth between the plants.

- 5 69. A method for selecting a plant which comprises applying a protoporphyrinogen IX oxidase inhibitory-type compound and a compound inhibiting 5-enolpyruvylshikamate-3-phosphate synthase to a growth area of the plant of claim 63 and other plants, and selecting
0 either plant on the basis of difference in growth between the plants.

Abstract of the disclosure:

Weed control compound-resistant plants are produced by introducing a gene encoding a protein having the following characteristics (a) to (c):

(a) having a specific affinity for a substance which is concerned with the weed control activity of a weed control compound,

(b) having substantially no capability of modifying a substance for which said protein has a specific affinity, and

(c) being substantially free from framework regions of variable regions in an immunoglobulin, into a plant cell, and expressing the gene.

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11




Fig. 12




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17




Fig. 18




Fig. 19




Fig. 20




Fig. 21




Fig. 22




Fig. 23




Fig. 24




Fig. 25




Fig. 26




Fig. 27




Fig. 28




Fig. 29




Fig. 30




Fig. 31




Fig. 32




Fig. 33




Fig. 34




Fig. 35




Fig. 36




Fig. 37




Fig. 38




Fig. 39




Fig. 40




Fig. 41





Fig. 42



BIRCH, STEWART, KOLASCH & BIRCH, LLP

P.O. Box 747 • Falls Church, Virginia 22040-0747

Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

ATTORNEY DOCKET NO.
20-4764P

PLEASE NOTE:
YOU MUST
COMPLETE THE
FOLLOWING: ↓

COMBINED DECLARATION AND POWER OF ATTORNEY FOR PATENT AND DESIGN APPLICATIONS

As a below named inventor, I hereby declare that: my residence, post office address and citizenship are as stated next to my name; that I verily believe that I am the original, first and sole inventor (if only one inventor is named below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Insert Title: → METHOD FOR GIVING RESISTANCE TO WEED CONTROL COMPOUNDS TO
PLANTS

Fill in Appropriate
Information —
For Use
Without
Specification
Attached: →

the specification of which is attached hereto. If not attached hereto,

the specification was filed on _____ as
United States Application Number _____;
and amended on _____ (if applicable); and/or
the specification was filed on _____ as PCT
International Application Number _____; and was
amended under PCT Article 19 on _____ (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I do not know and do not believe the same was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to this application, that the same was not in public use or on sale in the United States of America more than one year prior to this application, that the invention has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months (six months for designs) prior to this application, and that no application for patent or inventor's certificate on this invention has been filed in any country foreign to the United States of America prior to this application by me or my legal representatives or assigns, except as follows.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 (a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)

Priority Claimed

120553/1998	Japan	04/30/1998	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
(Number)	(Country)	(Month / Day / Year Filed)		
281127/1998	Japan	10/02/1998	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
(Number)	(Country)	(Month / Day / Year Filed)		
330981/1998	Japan	11/20/1998	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
(Number)	(Country)	(Month / Day / Year Filed)		
054730/1999	Japan	03/02/1999	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
(Number)	(Country)	(Month / Day / Year Filed)		

Insert Priority
Information:
(if appropriate) →

Insert Provisional
Application(s): →
(if any)

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

_____	_____
(Application Number)	(Filing Date)
_____	_____
(Application Number)	(Filing Date)

All Foreign Applications, if any, for any Patent or Inventor's Certificate Filed More than 12 Months (6 Months for Designs) Prior to the Filing Date of This Application:

Country	Application Number	Date of Filing (Month / Day / Year)
_____	_____	_____
_____	_____	_____

Insert Requested
Information:
(if appropriate) →

I hereby claim the benefit under Title 35, United States Code, §120 of any United States and/or PCT application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States and/or PCT application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

09/30/2357	04/30/1999	Pending
(Application Number)	(Filing Date)	(Status — patented, pending, abandoned)
_____	_____	_____
(Application Number)	(Filing Date)	(Status — patented, pending, abandoned)

Insert Prior U.S.
Application(s): →
(if any)

I hereby appoint the following attorneys to prosecute this application and/or an international application based on this application and to transact all business in the Patent and Trademark Office connected therewith and in connection with the resulting patent based on instructions received from the entity who first sent the application papers to the attorneys identified below, unless the inventor(s) or assignee provides said attorneys with a written notice to the contrary:

Raymond C. Stewart (Reg. No. 21,066)
 Joseph A. Kolasch (Reg. No. 22,463)
 Bernard L. Sweeney (Reg. No. 24,448)
 Charles Gorenstein (Reg. No. 29,271)
 Leonard R. Svensson (Reg. No. 30,330)
 Andrew D. Meikle (Reg. No. 32,868)
 Joe McKinney Muncy (Reg. No. 32,334)
 Donald J. Daley (Reg. No. 34,313)
 John A. Castellano (Reg. No. 35,094)

Terrell C. Birch (Reg. No. 19,382)
 James M. Slattery (Reg. No. 28,380)
 Michael K. Mutter (Reg. No. 29,680)
 Gerald M. Murphy, Jr. (Reg. No. 28,977)
 Terry L. Clark (Reg. No. 32,644)
 Marc S. Weiner (Reg. No. 32,181)
 C. Joseph Faraci (Reg. No. 32,350)
 John W. Bailey (Reg. No. 32,881)

Send Correspondence to: **BIRCH, STEWART, KOLASCH & BIRCH, LLP**
 P.O. Box 747 • Falls Church, Virginia 22040-0747
 Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

PLEASE NOTE:
 YOU MUST
 COMPLETE
 THE
 FOLLOWING:
 ↓

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of First or
 Sole Inventor,
 Inventor Name of
 Inventor
 Insert Date This
 Document is Signed
 →
 Insert Residence
 Inventor Citizenship
 →
 Insert Post Office
 Address
 →
 Full Name of Second
 Inventor, if any;
 see above
 →
 Full Name of Third
 Inventor, if any
 see above
 →
 Full Name of Fourth
 Inventor, if any
 see above
 →
 Full Name of Fifth
 Inventor, if any
 see above
 →

GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Hiroki	NAKAJIMA	<i>Hiroki Nakajima</i>	10/3/00
Residence (City, State & Country)			CITIZENSHIP
Nishinomiya-shi, Hyogo-ken, Japan			Japan
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
2-29-301, Higashiyamadaai, Nishinomiya-shi, Hyogo-ken, Japan			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Akitsu	NAGASAWA	<i>Akitsu Nagasawa</i>	10/3/00
Residence (City, State & Country)			CITIZENSHIP
Kobe-shi, Hyogo-ken, Japan			Japan
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
5-14-8, Sumiyoshiyamate, Higashinada-ku, Kobe-shi, Hyogo-ken, Japan			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)			CITIZENSHIP
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)			CITIZENSHIP
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)			CITIZENSHIP
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			

* DATE OF SIGNATURE